

THE INDEX OF A GEODESIC IN A RANDERS SPACE AND SOME REMARKS ABOUT THE LACK OF REGULARITY OF THE ENERGY FUNCTIONAL OF A FINSLER METRIC

ERASMO CAPONIO

ABSTRACT. In a series of papers ([2, 3, 4]) the relations existing between the metric properties of Randers spaces and the conformal geometry of stationary Lorentzian manifolds were discovered and investigated. These relations were called in [4] *Stationary-to-Randers Correspondence (SRC)*. In this paper we focus on one aspect of SRC, the equality between the index of a geodesic in a Randers space and that of its lightlike lift in the associated conformal stationary spacetime. Moreover we make some remarks about regularity of the energy functional of a Finsler metric on the infinite dimensional manifold of H^1 curves connecting two points, in connection with infinite dimensional techniques in Morse Theory.

1. INTRODUCTION

Let S be a manifold of dimension n and $R = \sqrt{h} + \omega$ be a Randers metric on S . To (S, R) we associate a one-dimensional higher manifold $M = S \times \mathbb{R}$ endowed with the bilinear symmetric tensor

$$g = h - (\omega - dt)^2.$$

The condition on the norm of ω ensuring that R is a positive definite function on TS , i.e. $(\omega_p(v))^2 < h_p(v, v)$ for all $v \in T_pS$ and for all $p \in S$, makes g a non-degenerate symmetric bilinear form of index 1, that is a Lorentzian metric on $S \times \mathbb{R}$.

Let t be the natural coordinate on \mathbb{R} . The vector field $\partial_t = \frac{\partial}{\partial t}$ on $S \times \mathbb{R}$ is timelike at any point (i.e. $g_p(\partial_t, \partial_t) < 0$, for all $p \in M$) and it is a Killing

2000 *Mathematics Subject Classification.* 53C22, 53C50, 53C60, 58B20.

Key words and phrases. Stationary Lorentzian manifolds, lightlike geodesics, Morse index, Finsler metric, Randers space.

Supported by M.I.U.R. Research project PRIN07 “Metodi Variazionali e Topologici nello Studio di Fenomeni Nonlineari”.

vector field for g . A Lorentzian manifold admitting a timelike Killing vector field is called *stationary* (see for instance [12, p. 119]) and whenever the timelike Killing vector field is irrotational is said *static*.

For any fixed $p \in S$, the function $R(p, \cdot): T_p S \rightarrow [0, +\infty)$ arises as the non-negative solution of the equation in the variable τ

$$(1) \quad h_p(v, v) - (\omega_p(v) - \tau)^2 = 0.$$

Eq. (1) and $\tau \geq 0$ are the conditions that a future pointing lightlike vector $(v, \tau) \in T_p S \times \mathbb{R}$ has to satisfy by definition.

We recall that a Lorentzian manifold (M, g) is said *time-oriented* if it admits a smooth timelike vector field Y . In particular a stationary Lorentzian manifold is time-oriented by one of its timelike Killing vector field. A vector $v \in T_p M$ is said *future pointing* (resp. *past pointing*) if $g_p(v, Y) < 0$ (resp. $g_p(v, Y) > 0$) and *lightlike* if $g_p(v, v) = 0$. Analogously, a smooth curve $\gamma: [a, b] \rightarrow M$ is said future pointing, past pointing, lightlike iff its velocity vector field is future pointing, past pointing, lightlike. Observe that if (v, τ) is past pointing and lightlike then τ is equal to the non-positive solution of (1) and $-\tau$ is equal to the Randers metric obtained reversing R , that is $-\tau = R(p, -v)$.

In analogy with a terminology used for static spacetimes (cf. [9, p. 360]), a stationary Lorentzian manifold (M, g) is said *standard* if it is isometric to a product manifold $S \times \mathbb{R}$ endowed with the metric

$$g_0 + w \otimes dt + dt \otimes w - \beta dt^2,$$

where g_0 , w and β are respectively a Riemannian metric, a one-form and a positive function on S . The conditions defining future pointing lightlike vectors on (M, g) define now the non-negative function on TS

$$R = \sqrt{g_0/\beta + (w/\beta)^2} + w/\beta.$$

Whatever the one-form w is, the norm of w/β with respect to the Riemannian metric

$$(2) \quad h = g_0/\beta + (w/\beta)^2$$

is less than 1 and thus R is a Randers metric.

Since Eq. (1) is invariant under conformal transformations of the metric g , the same Randers metric R is associated to the conformal class of g . Conversely, a Randers space (S, R) individuates the conformal standard stationary Lorentzian manifold $(S \times \mathbb{R}, h - (\omega - dt)^2)$.

The bijection between Randers spaces and conformal standard stationary Lorentzian manifolds has been called in [4] *Stationary-to-Randers correspondence (SRC)* and it has been used in [2] and in [4] to study the causal structure of a conformal standard stationary Lorentzian manifold.

One of the basic observation about SRC is that there is a one-to-one correspondence between lightlike geodesics of the conformal standard stationary Lorentzian manifold and the geodesics of the associated Randers space. Going

into more details, we mention that lightlike geodesics on a Lorentzian manifold are invariant under conformal changes of the metric in the sense that if $\gamma: [0, 1] \rightarrow M$ is a lightlike geodesic of (M, g) then γ is a pregeodesic of λg for any positive function λ , i. e. there exists a reparametrization $\sigma: [0, 1] \rightarrow [0, 1]$ such that $\gamma \circ \sigma$ is a lightlike geodesic of $(M, \lambda g)$ (see for example [8, p. 14]). We consider now a conformal standard stationary Lorentzian manifold $(S \times \mathbb{R}, g)$ and we take, as representative of the class, the metric $h - (\omega - dt)^2$, where h is equal to (2) and $\omega = w/\beta$. If $z(s) = (x(s), t(s))$ is a future pointing lightlike geodesic of $(S \times \mathbb{R}, h - (\omega - dt)^2)$ then (see [2, Theorem 4.5]) x is a geodesic of the Randers space (S, R) , $R = \sqrt{h} + \omega$, parametrized with $h(\dot{x}, \dot{x}) = \text{const}$. The fact that x has to be parametrized with constant Riemannian speed can be seen recalling that $g(\dot{z}, \dot{z}) = 0$ and, since ∂_t is a Killing vector field, $g(\dot{z}, \partial_t) = \omega(\dot{x}) - \dot{t} = \text{const}$. thus also $h(\dot{x}, \dot{x})$ has to be constant.

The other way round, a geodesic $x = x(s)$ in (S, R) can be lifted to a future pointing lightlike curve on $S \times \mathbb{R}$ by taking

$$(3) \quad t = t(s) = t_0 + \int_{s_0}^s R(x, \dot{x}).$$

If x is parametrized with constant Riemannian speed, its future pointing lightlike lift is a lightlike geodesic of $(S \times \mathbb{R}, h - (\omega - dt)^2)$.

The same relation holds between geodesics of the reversed metric $\tilde{R}(x, v) = R(x, -v)$ and past pointing lightlike geodesic of $(S \times \mathbb{R}, h - (\omega - dt)^2)$.

In Section 2 of this note, we focus on one aspect of SRC that is the equality between the index of a geodesic in the Randers space (S, R) and the index of its future pointing lightlike lift in $(S \times \mathbb{R}, h - (\omega - dt)^2)$.

An immediate consequence of this equality (which holds also for a geodesic of the reversed Randers metric \tilde{R} and the corresponding past pointing lightlike geodesic of $(S \times \mathbb{R}, h - (\omega - dt)^2)$), is that the index of a lightlike geodesic is a conformal invariant for standard stationary Lorentzian manifolds. This gives an alternative proof to a well known fact which holds for any conformal Lorentzian manifold (see for example [8, Theorem 2.36]).

Another consequence of this equality is that the Morse theory for future pointing lightlike geodesic connecting a point $\tilde{p} = (p, t_0)$ to an integral line of the timelike Killing vector field ∂_t passing through the point $\tilde{q} = (q, t_0)$, can be reduced to the Morse theory for geodesics connecting the points p and q in the associated Randers space.

Although Morse theory for geodesics connecting two points on a Finsler manifold (M, F) can be developed by using finite dimensional approximations of the path space by broken geodesics (see [10]), infinite dimensional techniques in Morse theory can be adapted to work in the Sobolev manifold $\Omega_{p,q}(M)$ of the H^1 curves connecting the points p and q . The main problem in regard to this approach is the lack of twice Frechet differentiability of the energy functional E of a Finsler metric at any critical point with respect to the H^1 -topology.

Anyway E has enough regularity to get a version of the Morse Lemma which allows us to compute the critical groups and to obtain the Morse relations (see [3]). In Section 3 we illustrate what is the problem in trying to prove that E is twice Frechet differentiable with respect to the H^1 -topology and we will extend to the Finsler case a recent argument by A. Abbondandolo and M. Schwartz [1]. In fact, in [1] the authors prove that a smooth time dependent Lagrangian $L: [0, 1] \times TM \rightarrow \mathbb{R}$, which is subquadratic in the velocities and whose action functional is twice Frechet differentiable at a regular curve on the Sobolev manifold $\Omega(M)$ of all the H^1 curves on M , must be a polynomial of degree at most two in the velocity variables along the curve. This fact can be seen as an infinite dimensional version of the well known property that if the square of a Finsler metric is C^2 on the whole TM then actually it is the square of the norm of a Riemannian metric.

2. THE EQUALITY BETWEEN THE INDEXES

Let M be a Lorentzian or a Finsler manifold and let γ be a geodesic on M . By $\mu(\gamma)$ we denote the *index* of γ , that is the number of conjugate points along γ counted with their multiplicity. The equality between $\mu(x)$, where x is a geodesic of the Randers space (S, R) , and $\mu(z)$, where z is the future pointing lightlike lift of x in $(S \times \mathbb{R}, g = h - (\omega - dt)^2)$, can be carried out by comparing the Jacobi equation of x in (S, R) with the Jacobi equation of z in $(S \times \mathbb{R}, h - (\omega - dt)^2)$, as done in [3, Theorem 13].

Here we give a different proof based on a comparison of the Morse index of the energy functional of the Randers metric at x and the Morse index at z of the functional introduced by Uhlenbeck in [11]:

$$J(\sigma) = \int_0^1 \left(g(\dot{\sigma}, \dot{\sigma}) + \left(\frac{dP(\sigma)}{ds} \right)^2 \right) ds.$$

Here σ belongs to the set of piecewise differentiable curves on $S \times \mathbb{R}$, satisfying the constraint $g(\dot{\sigma}, \dot{\sigma}) = 0$ and the boundary conditions $\sigma(0) = \tilde{p} \in S \times \mathbb{R}$, $\sigma(1) \in l(\mathbb{R})$, where $l = l(s)$ is an integral line of the Killing vector field ∂_t ($\tilde{p} \notin l(\mathbb{R})$) and $P: S \times \mathbb{R} \rightarrow \mathbb{R}$ is the natural projection on \mathbb{R} .

The critical point of J are the lightlike geodesics connecting \tilde{p} to $l(\mathbb{R})$. Moreover J admits second variation at any critical point. A critical point is non degenerate if and only if its endpoints are non-conjugate. The Morse index of a critical point is finite and it is equal to $\mu(z)$ (see [11, Lemma 4.2]). Using these properties of J we can prove the following

Theorem 2.1. *Let $(S \times \mathbb{R}, h - (\omega - dt)^2)$ be the conformal standard stationary spacetime associated by SRC to (S, R) and $z(s) = (x(s), t(s)): [0, 1] \rightarrow S \times \mathbb{R}$ be the future pointing lightlike geodesic associated to the geodesic $x(s)$ in (S, R) . Then the points $x(0)$ and $x(1)$ are non-conjugate along x in (S, R) if and only if the points $z(0)$ and $z(1)$ are non-conjugate along z in $(S \times \mathbb{R}, h - (\omega - dt)^2)$.*

Moreover

$$\mu(z) = \mu(x).$$

Proof. Consider the energy functional of the Randers metric R

$$E(\gamma) = \frac{1}{2} \int_0^1 R^2(\gamma, \dot{\gamma}) ds.$$

Since the Morse index of E at the geodesic x is equal to $\mu(x)$ (see [6]) and the Morse index of J at z is equal to $\mu(z)$, it is enough to prove the equality for the Morse indexes. To this end, we will show that the set \mathcal{W}_x of continuous piecewise smooth vector field along x vanishing at $x(0)$ and $x(1)$ is isomorphic to the set of admissible variations \mathcal{U}_z for J which is given by the continuous piecewise smooth vector fields U along z , vanishing at $z(0)$ and $z(1)$ and such that $g(\dot{z}, U) = 0$ (see [11]). Let us denote by $\mathcal{P}_{x(0),x(1)}(S)$ and $\mathcal{L}_{z(0),l}(S \times \mathbb{R})$ respectively the set of the continuous, piecewise smooth curves on S , parametrized on the interval $[0, 1]$ and connecting $x(0)$ to $x(1)$ and the set of the continuous, piecewise smooth, future pointing, lightlike curves on $S \times \mathbb{R}$, parametrized on $[0, 1]$ and connecting $z(0)$ to $l(\mathbb{R})$. Consider the map

$$\Psi(\gamma)(s) = \left(\gamma(s), t_0 + \int_0^s R(\gamma, \dot{\gamma}) d\nu \right).$$

Recalling that the future pointing lightlike lift of a curve γ in S has t component in $S \times \mathbb{R}$ given by (3), we immediately see that Ψ maps $\mathcal{P}_{x(0),x(1)}(S)$ to $\mathcal{L}_{z(0),l}(S \times \mathbb{R})$.

We are going to show that the isomorphism between \mathcal{W}_x and \mathcal{U}_z is given by $\Psi'(x)$ where, for each $W \in \mathcal{W}_x$, $\Psi'(x)[W]$ is the vector field along z belonging to \mathcal{U}_z defined as $\frac{\partial}{\partial r}(\Psi \circ \varphi_0)(r, s)|_{r=0}$ where $\varphi_0 = \varphi_0(r, s): (-\varepsilon, \varepsilon) \times [0, 1] \rightarrow S$ is the variation of the geodesic x defined by W . Observe that, since x is a critical point of the length functional $x \mapsto \int_0^1 R(x, \dot{x}) ds$, for any $W \in \mathcal{W}_x$ there holds

$$(\Psi'(x)[W])(0) = (\Psi'(x)[W])(1) = 0.$$

Let I be the functional defined in the same way as J

$$I(\sigma) = \int_0^1 (g(\dot{\sigma}, \dot{\sigma}) + \left(\frac{dP(\sigma)}{ds}\right)^2) ds$$

but now σ varies on the set of the continuous, piecewise smooth, future pointing curves, non necessarily lightlike, connecting $z(0)$ to $l(\mathbb{R})$. For any future pointing lightlike curve $\sigma(s) = (\gamma(s), \tau(s))$ we have

$$J(\sigma) = I(\Psi(\gamma)) = 2E(\gamma).$$

Moreover, for any geodesic x of (S, R) and for any $W \in \mathcal{W}_x$, we have

$$(4) \quad (\Psi'(x)[W])(s) = \left(W(s), \int_0^s (R_x(x, \dot{x})[W] + R_v(x, \dot{x})[\dot{W}]) ds \right);$$

hence $\Psi'(x)$ is an injective map (in the right-hand side of (4), we have used the expression $R_x(x, \dot{x})[W] + R_v(x, \dot{x})[\dot{W}]$ which is meaningful only in local coordinates).

Let $U(s) = (W(s), u(s)) \in \mathcal{U}_z$. We are going to show that $\Psi'(x)[W] = U$ and hence $\Psi'(x)$ is also surjective.

As $U \in \mathcal{U}_z$, we have

$$g(U, \dot{z}) = 0 \iff h(W, \dot{x}) - (\omega(W) - u)(\omega(\dot{x}) - \dot{t}) = 0.$$

Since z is lightlike and future pointing $\omega(\dot{x}) - \dot{t} = -\sqrt{h(\dot{x}, \dot{x})}$ and thus

$$u = \frac{h(W, \dot{x})}{\sqrt{h(\dot{x}, \dot{x})}} + \omega(W)$$

Since x is a critical point of E and $W(0) = 0$, integrating by part the t component of the vector field $\Psi'(x)[W]$ in (4) and using the Euler-Lagrange equation satisfied by x , we deduce that such a component is equal to

$$R_v(x, \dot{x})[W] = \frac{h(W, \dot{x})}{\sqrt{h(\dot{x}, \dot{x})}} + \omega(W) = u.$$

Now let $\varphi = \varphi(r, s): (-\varepsilon, \varepsilon) \times [0, 1] \rightarrow S \times \mathbb{R}$ be a variation defined by the admissible variational vector field $U = (W, u)$, and

$$\varphi_0 = \varphi_0(r, s): (-\varepsilon, \varepsilon) \times [0, 1] \rightarrow S$$

be the one defined by W , we have that

$$\begin{aligned} J''(z)(U, U) &= \frac{d^2}{dr^2} J(\varphi(r, \cdot))|_{r=0} \\ &= \frac{d^2}{dr^2} I(\Psi(\varphi_0(r, \cdot)))|_{r=0} = 2 \frac{d^2}{dr^2} E(\varphi_0(r, \cdot))|_{r=0} = 2E''(x)(W, W). \end{aligned}$$

By polarization, the above equality gives the thesis. □

3. THE LACK OF TWICE DIFFERENTIABILITY OF THE ENERGY FUNCTIONAL WITH RESPECT TO THE H^1 -TOPOLOGY

Let (M, F) be a Finsler manifold and $p, q \in M$. Let $\Omega(M)$ be the Sobolev manifold of the absolutely continuous curves $\gamma: [0, 1] \rightarrow M$, whose square of the norm of the velocity vector field is integrable with respect to a fixed (and then to any) auxiliary Riemannian metric α on M . Let us denote by $\Omega_{p,q}(M)$ the submanifold of the curves in $\Omega(M)$, such that $\gamma(0) = p, \gamma(1) = q$ (see [5]). Let us consider the energy functional of F on $\Omega_{p,q}(M)$:

$$E: \Omega_{p,q}(M) \rightarrow \mathbb{R}, \quad E(\gamma) = \frac{1}{2} \int_0^1 F^2(\gamma, \dot{\gamma}) ds$$

It is well known that E is $C^{1,1}$ on $\Omega_{p,q}(M)$, [7].

We are going to show that if E is twice differentiable on $\Omega_{p,q}(M)$ at a regular curve γ then F^2 is the square of the norm of a Riemannian metric along the curve.

By *regular curve* we mean a curve $\gamma \in \Omega_{p,q}(M)$ such that $\dot{\gamma} \neq 0$ a. e. in $[0, 1]$.

Remark 3.1. We point out that in [1] the authors consider a time-dependent Lagrangian $L: [0, 1] \times TM \rightarrow \mathbb{R}$, $L = L(t, q, v)$, which is C^2 on TM and which satisfies the following conditions: there exists a continuous positive function $C = C(q)$ such that for any $(t, q, v) \in [0, 1] \times TM$:

$$\begin{aligned} \|\partial_{vv}L(t, q, v)\| &\leq C(q), \\ \|\partial_{vq}L(t, q, v)\| &\leq C(q)(1 + \sqrt{\alpha(v, v)}), \\ \|\partial_{qq}L(t, q, v)\| &\leq C(q)(1 + \alpha(v, v)). \end{aligned}$$

They prove that if the action functional of L

$$\gamma: \Omega(M) \rightarrow \int_0^1 L(t, \gamma(t), \dot{\gamma}(t))dt,$$

is twice differentiable in $\Omega(M)$ at a curve γ , then the map

$$v \in T_{\gamma(t)}M \mapsto L(t, \gamma(t), v)$$

is a polynomial of degree at most two. Thus, in particular, the subquadratic and strongly convex in the velocities, time-independent, C^2 Lagrangians whose action functional is twice differentiable at any curve in $\Omega(M)$ are all and only of the type

$$L(q, v) = h_q(v, v) + \omega_q(v) + V(q),$$

where h , ω and V are respectively a Riemannian metric, a one-form and a function on M . Clearly, the square of a Finsler metric satisfies the growth conditions above but it is only a $C^{1,1}$ function on TM (it is C^2 on $TM \setminus 0$). Anyway, as we show below, the proof in [1] does not involve existence and continuity of the derivatives $\partial_{vv}L(t, q, v)$ for $v = 0$ and then it extends also to the Finsler case. Another difference from [1] is that we consider the manifold $\Omega_{p,q}(M)$ and not $\Omega(M)$.

Before going into the details of the proof, we would like to point out what is the problem in trying to prove that E is twice differentiable in $\Omega_{p,q}(M)$ at a regular curve. To fix ideas, we assume that F is defined on an open subset U of \mathbb{R}^n , $F: TU \rightarrow \mathbb{R}$, $U \subset \mathbb{R}^n$. Arguing as in [1, Proposition 3.1] gives that E is twice Gateaux differentiable in $\Omega_{p,q}(U)$ at any regular curve x and its second Gateaux differential is equal to

$$\begin{aligned} D^2\tilde{E}(x)[\xi, \eta] &= \frac{1}{2} \int_0^1 (\partial_{qq}F^2(x, \dot{x})[\xi, \eta] + \partial_{vq}F^2(x, \dot{x})[\dot{\xi}, \eta])ds \\ &\quad + \frac{1}{2} \int_0^1 (\partial_{qv}F^2(x, \dot{x})[\xi, \dot{\eta}])ds + \partial_{vv}F^2(x, \dot{x})[\dot{\xi}, \dot{\eta}]ds. \end{aligned}$$

The problem is the continuity of the map

$$x \in \Omega_{p,q}(U) \mapsto \int_0^1 \partial_{vv} F^2(x, \dot{x})[\cdot, \cdot] ds,$$

where the target space is the space of bounded bilinear operators on $H_0^1([0, 1], U)$. Namely, we can prove that if $x_n \rightarrow x$ in $\Omega_{p,q}(U)$ then

$$\int_0^1 \partial_{vv} F^2(x_n, \dot{x}_n)[\dot{\xi}, \dot{\eta}] ds \rightarrow \int_0^1 \partial_{vv} F^2(x, \dot{x})[\dot{\xi}, \dot{\eta}] ds,$$

as $n \rightarrow +\infty$, but we cannot prove that the convergence is uniform with respect to ξ and η in the unit ball of $H_0^1([0, 1], U)$, unless $\partial_{vv} F^2$ is independent from v (and then F^2 is the square of the norm of a Riemannian metric). In fact, we have the following

Proposition 3.2. *If the energy functional of a Finsler metric F is twice differentiable at a regular curve $\gamma \in \Omega_{p,q}(M)$ then for a. e. $s \in [0, 1]$ the function*

$$v \in T_{\gamma(s)}M \mapsto F^2(\gamma(s), v)$$

is a quadratic positive definite form.

Proof. For simplicity and without loss of generality, we prove the statement in the case where M is an open subset of \mathbb{R}^n . Since $\dot{\gamma} \neq 0$ a. e. on $[0, 1]$, the thesis is equivalent to the fact that for almost every $s \in [0, 1]$, there holds

$$\partial_v F^2(\gamma(s), \dot{\gamma}(s) + v) - \partial_v F^2(\gamma(s), \dot{\gamma}(s)) - \partial_{vv} F^2(\gamma(s), \dot{\gamma}(s))[v] = 0,$$

for all $v \in \mathbb{R}^n$. By contradiction, we assume that there is a set of positive measure $J \subset [0, 1]$ and two non-zero vectors $v, w \in \mathbb{R}^n$, and a positive number c such that

$$(5) \quad \left(\partial_v F^2(\gamma(s), \dot{\gamma}(s) + v) - \partial_v F^2(\gamma(s), \dot{\gamma}(s)) - \partial_{vv} F^2(\gamma(s), \dot{\gamma}(s))[v] \right) \cdot w > c,$$

For every $\epsilon > 0$ smaller than the measure of J , choose a subset $J_\epsilon \subset J$ of measure ϵ , in such a way that $J_\epsilon \subset J_{\epsilon'}$ if $\epsilon < \epsilon'$. Define the following functions

$$\eta_\epsilon(s) = v \int_0^s (\chi_\epsilon(t) - \epsilon) dt, \quad \xi_\epsilon(s) = w \int_0^s (\chi_\epsilon(t) - \epsilon) dt,$$

where χ_ϵ is the characteristic function of J_ϵ . Observe that, for any ϵ , the functions $\eta_\epsilon, \xi_\epsilon$ belong to $T_\gamma \Omega_{p,q}(M) = H_0^1([0, 1], \mathbb{R}^n)$ and

$$\|\eta_\epsilon\|_{H_0^1} = |v|(\epsilon - \epsilon^2)^{1/2} \quad \|\xi_\epsilon\|_{H_0^1} = |w|(\epsilon - \epsilon^2)^{1/2}.$$

We can repeat the proof of Proposition 3.2 in [1] taking care only that the derivatives of η and ξ here are given by $v(\chi_\epsilon - \epsilon)$ and $w(\chi_\epsilon - \epsilon)$ and the terms involving integrals of the type

$$\int_0^1 (\partial_v F^2(\gamma + \eta_\epsilon, \dot{\gamma} + \dot{\eta}_\epsilon) - \partial_v F^2(\gamma, \dot{\gamma} + \dot{\eta}_\epsilon) - \partial_{qv} F^2(\gamma, \dot{\gamma} + \dot{\eta}_\epsilon)[\eta_\epsilon]) \cdot (\epsilon w) ds$$

belong to $o(\epsilon)$, as $\epsilon \rightarrow 0$ (such terms do not appear in [1] because the functions playing the role of η and ξ , not having to belong to $H_0^1([0, 1], \mathbb{R}^n)$, are defined as $\eta(s) = v \int_0^s \chi_\epsilon(t)dt$ and $\xi(s) = w \int_0^s \chi_\epsilon(t)dt$).

We point out that the non existence of the derivatives $\partial_{vv}F^2(q, v)$ for $v = 0$ does not affect that part of the proof since only the smoothness of $\partial_v F^2(q, v)$ with respect to q is used.

Thus, as in [1], we can deduce

$$(6) \quad \int_0^1 (\partial_v F^2(\gamma, \dot{\gamma} + \dot{\eta}_\epsilon) - \partial_v F^2(\gamma, \dot{\gamma}) - \partial_{vv} F^2(\gamma, \dot{\gamma})[\dot{\eta}_\epsilon]) \cdot \dot{\xi}_\epsilon ds = o(\epsilon),$$

as $\epsilon \rightarrow 0$. The left-hand side of (6) is equal to

$$(7) \quad \begin{aligned} & \int_0^1 (\partial_v F^2(\gamma, \dot{\gamma} + \dot{\eta}_\epsilon) - \partial_v F^2(\gamma, \dot{\gamma}) - \partial_{vv} F^2(\gamma, \dot{\gamma})[\dot{\eta}_\epsilon]) \cdot (\chi_\epsilon w - \epsilon w) ds \\ &= \int_{J_\epsilon} (\partial_v F^2(\gamma, \dot{\gamma} + (1 - \epsilon)v) - \partial_v F^2(\gamma, \dot{\gamma}) - \partial_{vv} F^2(\gamma, \dot{\gamma})[(1 - \epsilon)v]) \cdot w ds \\ & \quad + \int_0^1 (\partial_v F^2(\gamma, \dot{\gamma} + \dot{\eta}_\epsilon) - \partial_v F^2(\gamma, \dot{\gamma}) - \partial_{vv} F^2(\gamma, \dot{\gamma})[\dot{\eta}_\epsilon]) \cdot (\epsilon w) ds \end{aligned}$$

Since $\dot{\eta}_\epsilon \rightarrow 0$ a. e. as $\epsilon \rightarrow 0$, by the Lebesgue's dominated convergence theorem, the absolute value of the second integral in the right-hand side of (7) is less than $\epsilon|w|o(1)$.

Therefore, putting together (6) and (7), we have

$$(8) \quad \int_{J_\epsilon} (\partial_v F^2(\gamma, \dot{\gamma} + (1 - \epsilon)v) - \partial_v F^2(\gamma, \dot{\gamma}) - \partial_{vv} F^2(\gamma, \dot{\gamma})[(1 - \epsilon)v]) \cdot w ds = o(\epsilon),$$

as $\epsilon \rightarrow 0$. By (5) and the continuity of the map

$$v \in \mathbb{R}^n \mapsto \left(\partial_v F^2(\gamma(s), \dot{\gamma}(s) + v) - \partial_v F^2(\gamma(s), \dot{\gamma}(s)) - \partial_{vv} F^2(\gamma(s), \dot{\gamma}(s))[v] \right) \cdot w$$

the integral in (8) is larger than $c\epsilon$, for ϵ small enough, giving a contradiction. \square

REFERENCES

[1] A. Abbondandolo and M. Schwarz. A smooth pseudo-gradient for the Lagrangian action functional. *Adv. Nonlinear Stud.*, 9(4):597–623, 2009.
 [2] E. Caponio, M. A. Javaloyes, and A. Masiello. On the energy functional on Finsler manifolds and applications to stationary spacetimes. arXiv:math/0702323v3, 2008. *Math. Ann.* to appear.
 [3] E. Caponio, M. Á. Javaloyes, and A. Masiello. Morse theory of causal geodesics in a stationary spacetime via Morse theory of geodesics of a Finsler metric. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 27(3):857–876, 2010.
 [4] E. Caponio, M. A. Javaloyes, and M. Sánchez. On the interplay between lorentzian causality and finsler metrics of randers type. arXiv:0903.3501, 2009.
 [5] W. Klingenberg. *Riemannian geometry*. Walter de Gruyter & Co., Berlin, 1982.
 [6] M. Matsumoto. *Foundations of Finsler geometry and special Finsler spaces*. Kaiseisha Press, Shigaken, 1986.

- [7] F. Mercuri. The critical points theory for the closed geodesics problem. *Math. Z.*, 156(3):231–245, 1977.
- [8] E. Minguzzi and M. Sánchez. The causal hierarchy of spacetimes. In *Recent developments in pseudo-Riemannian geometry*, ESI Lect. Math. Phys., pages 299–358. Eur. Math. Soc., Zürich, 2008.
- [9] B. O’Neill. *Semi-Riemannian geometry*, volume 103 of *Pure and Applied Mathematics*. Academic Press Inc., New York, 1983.
- [10] Z. Shen. *Lectures on Finsler geometry*. World Scientific Publishing Co., Singapore, 2001.
- [11] K. Uhlenbeck. A Morse theory for geodesics on a Lorentz manifold. *Topology*, 14:69–90, 1975.
- [12] R. M. Wald. *General relativity*. University of Chicago Press, Chicago, 1984.

DIPARTIMENTO DI MATEMATICA,
POLITECNICO DI BARI,
VIA ORABONA 4, 70125, BARI, ITALY
E-mail address: `caponio@poliba.it`