

THE ISOMORPHISM PROBLEM FOR SEMIGROUP C^* -ALGEBRAS
OF RIGHT-ANGLED ARTIN MONOIDS

SØREN EILERS, XIN LI, AND EFREN RUIZ

Received: February 2, 2016

Communicated by Joachim Cuntz

ABSTRACT. Semigroup C^* -algebras for right-angled Artin monoids were introduced and studied by Crisp and Laca. In the paper at hand, we are able to present the complete answer to their question of when such C^* -algebras are isomorphic. The answer to this question is presented both in terms of properties of the graph defining the Artin monoids as well as in terms of classification by K -theory, and is obtained using recent results from classification of non-simple C^* -algebras. Moreover, we are able to answer another natural question: Which of these semigroup C^* -algebras for right-angled Artin monoids are isomorphic to graph algebras? We give a complete answer, and note the consequence that many of the C^* -algebras under study are semiprojective.

2010 Mathematics Subject Classification: Primary: 46L05. Secondary: 20F36, 20M05, 46L35.

1 INTRODUCTION

Semigroup C^* -algebras for right-angled Artin monoids were introduced and studied by Crisp and Laca in [CL02] and [CL07]. In [CL07], the authors ask how to classify these semigroup C^* -algebras up to $*$ -isomorphism. We now present the complete answer to their question.

The Artin monoids studied here are given by countable, symmetric and antireflexive graphs $\Gamma = (V, E)$ as

$$A_{\Gamma}^{\dagger} := \langle \{\sigma_v : v \in V\} \mid \sigma_v \sigma_w = \sigma_w \sigma_v \text{ if } (v, w) \in E \rangle^{\dagger}.$$

The corresponding right-angled Artin groups, defined by the same generators and relations, are special cases of Artin groups, which form an important class of examples of groups. We refer the reader to [CL02, CL07] and the references therein for more details.

Semigroup C^* -algebras of left cancellative semigroups, generated by the left regular representation of the semigroup, have been studied for a long time. Recently, there has been a renewed interest in this topic (see [Li12, Li13] and the references therein). By [CL02], the semigroup C^* -algebras $C^*(A_\Gamma^+)$ attached to right-angled Artin monoids are given as the universal C^* -algebras for

$$\left\langle \{s_v: v \in V\} \left| \begin{array}{l} [s_v, s_w] = [s_v, s_w^*] = 0 \text{ if } (v, w) \in E \\ s_v^* s_w = \delta_{v,w} \text{ if } (v, w) \notin E \end{array} \right. \right\rangle$$

We answer the question of when two graphs Γ, Λ produce C^* -algebras that are isomorphic. Although we emphasize that our results cover the full range of such graphs, it is instructive to state our main results in the case of finite graphs. This is a specialization of the combination of Theorems 4.2 and 5.2.

THEOREM 1.1 *Let Γ and Λ be finite undirected graphs with no loops. The following are equivalent*

1. $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$
2. (a) $t(\Gamma) = t(\Lambda)$
 (b) $N_k(\Gamma) + N_{-k}(\Gamma) = N_k(\Lambda) + N_{-k}(\Lambda)$ for all $k \in \mathbb{Z}$
 (c) $N_0(\Gamma) > 0$ or

$$\sum_{k>0} N_{-k}(\Gamma) \equiv \sum_{k>0} N_{-k}(\Lambda) \pmod{2}$$

3. $[FK_+(C^*(A_\Gamma^+)), [1_{C^*(A_\Gamma^+)}]] \cong [FK_+(C^*(A_\Lambda^+)), [1_{C^*(A_\Lambda^+)}]]$

In this result, the invariant mentioned in (3) is the standard ordered filtered K -theory – implicitly containing the primitive ideal space – which has been conjectured in [ERR10] to be a complete invariant for a large and important class of C^* -algebras. This conjecture is still open, but has been confirmed in a multitude of situations partially overlapping with the case at hand. But the main strength of our result is in fact that the *ad hoc* invariant of (2) is extremely easy to compute for Γ and Λ . Indeed, as we shall detail below, the numbers $t(\Gamma)$ and $N_k(\Gamma)$ are obtained by dividing Γ into co-irreducible components and then counting how many of these are singletons, yielding $t(\Gamma)$, and counting how many of the remaining co-irreducible components have Euler characteristic k , yielding $N_k(\Gamma)$. In Figure 1 this process has been completed for all 34 graphs with five vertices, and we conclude that they define 18 different C^* -algebras. When the number of vertices increase, it is possible for two graphs to have different sets of invariants, yet define the same C^* -algebra. For instance, defining




Figure 1: Invariants for all graphs with 5 vertices. Any quantity not mentioned equals zero.

a graph Γ' with 10 vertices having its co-irreducible components chosen among those given in Figure 1 so that

$$N_{-1}(\Gamma') = 2$$

and Λ' similarly defined so that

$$N_1(\Lambda') = 2,$$

then $C^*(A_{\Gamma'}^+)$ and $C^*(A_{\Lambda'}^+)$ will be isomorphic. Similarly, we may define Γ'' and Λ'' with 15 vertices each so that

$$\begin{aligned} N_{-1}(\Gamma'') &= 1, N_0(\Gamma'') = 1, N_1(\Gamma'') = 1 \\ N_{-1}(\Lambda'') &= 2, N_0(\Lambda'') = 1 \end{aligned}$$

obtaining that $C^*(A_{\Gamma''}^+) \cong C^*(A_{\Lambda''}^+)$.

In the general case of possibly infinite graphs, an additional quantity $o(\Gamma)$ must be introduced to count those co-irreducible components which have an infinite number of vertices, and to address the possibility of having an infinite number of co-irreducible components, but the necessary condition replacing (2) in this general case is not much more complicated than the one given above.

We note that the C^* -algebras associated via semigroups to undirected and loop-free graphs are not always graph C^* -algebras in the usual sense, not only because graph C^* -algebras are defined using directed graphs. We provide a complete description of when $C^*(A_{\Gamma}^+)$ is in fact a graph C^* -algebra, and note that there is a rather complicated relation between Γ and the G_{Γ} when in fact $C^*(A_{\Gamma}^+) \cong C^*(G_{\Gamma})$. In this case, G_{Γ} is not unique.

Our results have surprising consequences for the issue of *stable relations* (cf. [Lor97]) among sets of isometries of separable Hilbert spaces, subject to commutativity or orthogonality relations as given by the graph Γ , or, which is nearly the same, for the issue of *semiprojectivity* (cf. [Bla85]) of the C^* -algebras $C^*(A_{\Gamma}^+)$. Indeed, it is easy to see by spectral theory that $C^*(A_{\Gamma}^+)$ is semiprojective when Γ is a finite graph with no edges, corresponding to a family of isometries having orthogonal range projections. Similarly, it follows e.g. from considering the celebrated Voiculescu matrices ([Voi83],[EL91]) that when Γ is a complete graph with more than one vertex, $C^*(A_{\Gamma}^+)$ cannot have this property. It is a question attracting a lot of attention (see e.g. [Bla04]) to what extent it is possible to obtain stable relations for commuting sets of stable relations, or to what extent tensor products of semiprojective C^* -algebras can themselves be semiprojective. In fact, it was only established recently ([End15]) that $\mathcal{O}_3 \otimes \mathcal{O}_3$ is semiprojective, as a consequence of Enders' sweeping solution to the semiprojectivity problem for Kirchberg algebras. In our setting, because we have found that many settings in which some isometries are required to be orthogonal, and others to commute, give the same C^* -algebras as the ones where all are required to be orthogonal, we immediately see that many such settings — for instance the first 12 listed in Figure 1 — provide for stable

relations. Involving the notion of graph algebras as outlined above, we will show in Theorem 6.9 semiprojectivity and nonsemiprojectivity for many of the C^* -algebras under study, and it follows from our results that exactly those C^* -algebras arising from the graphs in Figure 1 in the non-shaded entries are semiprojective. We have not been able to resolve the issue completely as Enders’ methods do not apply directly, the first open case having six vertices and two co-irreducible components each with Euler characteristic -2 .

All authors acknowledge support by the Danish National Research Foundation through the Centre for Symmetry and Deformation (DNRF92), and thank the Department of Mathematical Sciences at the University of Copenhagen, where the initial phases of this work were carried out, for providing excellent facilities. The first named author gratefully acknowledges support from the VILLUM Foundation, and the third named author gratefully acknowledges support from the Simons Foundation (#279369 to Efren Ruiz).

2 PRELIMINARIES

2.1 SEMIGROUP C^* -ALGEBRAS FOR RIGHT-ANGLED ARTIN MONOIDS

Let Γ be a countable graph. $\Gamma = (V, E)$ is given by a countable set of vertices V and a set of edges E . We only consider unoriented edges, and given two vertices, there is at most one edge joining these two vertices. In other words, we can think of E as a symmetric subset of $V \times V$ not containing elements of the diagonal.

Given such a graph $\Gamma = (V, E)$, the right-angled Artin group A_Γ is defined as follows:

$$A_\Gamma := \langle \{\sigma_v : v \in V\} \mid \sigma_v \sigma_w = \sigma_w \sigma_v \text{ if } (v, w) \in E \rangle.$$

Similarly, the right-angled Artin monoid A_Γ^+ is defined as follows:

$$A_\Gamma^+ := \langle \{\sigma_v : v \in V\} \mid \sigma_v \sigma_w = \sigma_w \sigma_v \text{ if } (v, w) \in E \rangle^+.$$

It turns out that the canonical semigroup homomorphism $A_\Gamma^+ \rightarrow A_\Gamma$ is injective, see [Par02]. Moreover, it is shown in [CL02] that $A_\Gamma^+ \subseteq A_\Gamma$ is quasi-lattice ordered. This means that for every g in A_Γ , either $(gA_\Gamma^+) \cap A_\Gamma^+ = \emptyset$ or there exists $p \in A_\Gamma^+$ with $(gA_\Gamma^+) \cap A_\Gamma^+ = pA_\Gamma^+$.

The (left) reduced semigroup C^* -algebra of A_Γ^+ is given by

$$C_\lambda^*(A_\Gamma^+) = C^* \langle \{S_v : v \in V\} \rangle \subseteq \mathcal{L}(\ell^2(A_\Gamma^+)),$$

where $S_v e_x = e_{\sigma_v x}$ with $\{e_x\}$ the canonical orthonormal basis. The full semigroup C^* -algebra of A_Γ^+ is defined as

$$C^*(A_\Gamma^+) = C_{\text{univ}}^* \left\langle \{s_v : v \in V\} \mid \begin{array}{l} [s_v, s_w] = [s_v, s_w^*] = 0 \text{ if } (v, w) \in E \\ s_v^* s_w = \delta_{v,w} \text{ if } (v, w) \notin E \end{array} \right\rangle$$

The canonical homomorphism $C^*(A_\Gamma^+) \rightarrow C_\lambda^*(A_\Gamma^+)$ is an isomorphism by [CL02]. Hence we do not distinguish between reduced and full versions and simply write $C^*(A_\Gamma^+)$ for the semigroup C^* -algebra of A_Γ^+ .

2.2 CO-IRREDUCIBLE COMPONENTS

The graph Γ is called co-reducible if there exist non-empty subsets V_1 and V_2 of V with $V = V_1 \sqcup V_2$ such that $V_1 \times V_2 \subseteq E$. Γ is called co-irreducible if Γ is not co-reducible. In general, we can always decompose Γ into co-irreducible components. This means that there exist co-irreducible graphs $\Gamma_i = (V_i, E_i)$ such that $A_\Gamma^+ \cong \bigoplus_i A_{\Gamma_i}^+$ (and also $A_\Gamma \cong \bigoplus_i A_{\Gamma_i}$). As explained in [CL07], these co-irreducible components are found by looking at the opposite graph of Γ . For the semigroup C^* -algebra, we get $C^*(A_\Gamma^+) \cong \bigotimes_i C^*(A_{\Gamma_i}^+)$. Note that if there are (necessarily countably) infinitely many co-irreducible components, the tensor product is defined as an inductive limit of finite tensor products with respect to the canonical unital embeddings as tensor factors.

It is shown in [CL07] that for a co-irreducible graph $\Gamma = (V, E)$ with $1 < |V| < \infty$, $C^*(A_\Gamma^+)$ has a unique non-trivial ideal isomorphic to the compact operators. It is easy to see the compact operators in the description of $C^*(A_\Gamma^+)$ as a concrete C^* -algebra on $\ell^2(A_\Gamma^+)$: We just have to observe that $1 - \bigvee_{v \in V} S_v S_v^*$ is the orthogonal projection onto the one-dimensional subspace of $\ell^2(A_\Gamma^+)$ corresponding to the identity element of A_Γ^+ . This projection then generates the ideal of compact operators. The corresponding quotient $C_Q^*(A_\Gamma^+)$ is a (unital) Kirchberg algebra satisfying the UCT. However, if our co-irreducible graph has infinitely many vertices, then $C^*(A_\Gamma^+)$ itself is a (unital) Kirchberg algebra satisfying the UCT. That we obtain UCT Kirchberg algebras follows also from [Li13, Corollary 7.23]. The case where Γ consists of only one vertex is easy to understand; in that case, $C^*(A_\Gamma^+)$ is canonically isomorphic to the Toeplitz algebra \mathcal{T} .

2.3 PRIMITIVE IDEAL SPACE

We can now describe the primitive ideal space of $C^*(A_\Gamma^+)$ for arbitrary Γ . Let $\Gamma_i = (V_i, E_i)$ be the co-irreducible components of Γ . Then by [Bla77, Theorem 4.9], we have an identification

$$\text{Prim}(C^*(A_\Gamma^+)) \cong \prod_i \text{Prim}(C^*(A_{\Gamma_i}^+)).$$

Under this identification, an element (I_i) of the space on the right hand side corresponds to the primitive ideal I of $C^*(A_\Gamma^+)$ which is generated by $\{\bigotimes_j J_{ij}\}_i$, where $J_{ij} = C^*(A_{\Gamma_j}^+)$ if $j \neq i$ and $J_{ii} = I_i$. Since each of the Γ_i is co-irreducible, the primitive ideal space $\text{Prim}(C^*(A_{\Gamma_i}^+))$ is easy to describe because of the results summarized above:

- If Γ_i just consists of one point, then $\text{Prim}(C^*(A_{\Gamma_i}^+))$ is homeomorphic to the primitive ideal space of the Toeplitz algebra. This means that as a set, $\text{Prim}(C^*(A_{\Gamma_i}^+))$ is the disjoint union of a point and a circle, and the non-empty open sets are given by unions of the point and open subsets in the usual topology of the circle.

- If Γ_i has more than one, but finitely many vertices, $\text{Prim}(C^*(A_{\Gamma_i}^+))$ consists of two points, one of which is closed (the corresponding primitive ideal is the ideal of compact operators) and the other one is dense.
- If Γ_i has infinitely many vertices, then $\text{Prim}(C^*(A_{\Gamma_i}^+))$ consists of only one point.

2.4 K-THEORY

K-theory for $C^*(A_{\Gamma}^+)$ and the quotients $C_Q^*(A_{\Gamma}^+)$ has been computed in [Iva10] in an ad hoc way, and can also be computed using [CEL13]. Let us explain the computation via the latter route. First of all, we need the Euler characteristic of a graph Γ . We view Γ as a simplicial complex by defining for every $n = 0, 1, 2, \dots$ the set of n -simplices by

$$K_n := \{\{v_0, \dots, v_n\} \subseteq V : (v_i, v_j) \in E \text{ for all } i, j \in \{0, \dots, n\}, i \neq j\}.$$

Then we set for a graph Γ with finitely many vertices $\chi(\Gamma) := 1 - \sum_{n=0}^{\infty} (-1)^n |K_n|$.

REMARK 2.1 *It is easy to see that there are co-irreducible graphs attaining any integer as its Euler characteristic. Indeed, letting Γ_{-m} denote the graph with $m + 1$ vertices and no edges, we clearly have*

$$\chi(\Gamma_{-m}) = -m.$$

Systematically generating positive characteristics is harder; one option is to let Γ_{n^2-1} denote the graph with $2n + 2$ vertices obtained by deleting one edge from the complete bipartite graph $K_{n+1, n+1}$ and note that

$$\chi(\Gamma_{n^2-1}) = n^2 - 1$$

To obtain positive characteristics in $\{(n - 1)^2, \dots, n^2 - 1\}$ one may simply add a suitable number of isolated vertices to Γ_{n^2-1} .

Now, by [CEL13, Theorem 5.2], we know that we always have $K_*(C^*(A_{\Gamma}^+)) \cong K_*(\mathbb{C})$, and $K_0(C^*(A_{\Gamma}^+)) \cong \mathbb{Z}$ is generated by the class of the unit [1]. Here we use that right-angled Artin groups satisfy the Baum-Connes conjecture with coefficients because these groups have the Haagerup property (see [NR97], and also [AD]). To compute K-theory for the quotient $C_Q^*(A_{\Gamma}^+)$ in the case that Γ has (more than one and) finitely many vertices, we consider the short exact sequence $0 \rightarrow \mathcal{K} \rightarrow C^*(A_{\Gamma}^+) \rightarrow C_Q^*(A_{\Gamma}^+) \rightarrow 0$ and its six term exact sequence in K-theory:

$$\begin{array}{ccccc}
 K_0(\mathcal{K}) & \longrightarrow & K_0(C^*(A_{\Gamma}^+)) & \longrightarrow & K_0(C_Q^*(A_{\Gamma}^+)) \\
 \uparrow & & & & \downarrow \\
 K_1(C_Q^*(A_{\Gamma}^+)) & \longleftarrow & K_1(C^*(A_{\Gamma}^+)) & \longleftarrow & K_1(\mathcal{K})
 \end{array} \tag{1}$$

Since both $K_1(\mathcal{K})$ and $K_1(C^*(A_\Gamma^+))$ vanish, all we have to do is to compute the homomorphism $K_0(\mathcal{K}) \cong \mathbb{Z} \rightarrow \mathbb{Z} \cong K_0(C^*(A_\Gamma^+))$. $K_0(\mathcal{K}) \cong \mathbb{Z}$ is generated by the class of any minimal projection. So we can take $e = 1 - \bigvee_{v \in V} S_v S_v^*$. It is easy to see that in K_0 , $[e] \in K_0(\mathcal{K})$ is sent to $\chi(\Gamma)[1] \in K_0(C^*(A_\Gamma^+))$. Therefore, by exactness of (1), we conclude that $K_0(C_Q^*(A_\Gamma^+)) \cong \mathbb{Z}/|\chi(\Gamma)|\mathbb{Z}$ and

$$K_1(C_Q^*(A_\Gamma^+)) \cong \begin{cases} \{0\} & \text{if } \chi(\Gamma) \neq 0 \\ \mathbb{Z} & \text{if } \chi(\Gamma) = 0 \end{cases} .$$

3 EXTENSION ALGEBRAS

We now discuss the C*-algebras associated to co-irreducible graphs and see how they are all isomorphic to either the Toeplitz algebra, the Cuntz algebra \mathcal{O}_∞ , or an *extension algebra* as specified below.

THEOREM 3.1 *Consider the following properties for a unital C*-algebra A:*

- (1) *A contains \mathcal{K} as an ideal, and A/\mathcal{K} is a Kirchberg algebra satisfying the UCT,*
- (2) *$K_0(A) = \mathbb{Z}$ with $[1_A] = 1$.*

For each $k \in \mathbb{Z} \setminus \{0\}$ there is a unique C-algebra satisfying (1), (2) and*

- (3) *The six-term exact sequence for \mathcal{K} and A is given by*

$$\begin{array}{ccccc} \mathbb{Z} & \xrightarrow{k} & \mathbb{Z} & \longrightarrow & \mathbb{Z}/k\mathbb{Z} \\ \uparrow & & & & \downarrow \\ 0 & \longleftarrow & 0 & \longleftarrow & 0 \end{array}$$

There is also a unique C-algebra satisfying (1), (2) and*

- (3') *The six-term exact sequence for \mathcal{K} and A is given by*

$$\begin{array}{ccccc} \mathbb{Z} & \xrightarrow{0} & \mathbb{Z} & \xlongequal{\quad} & \mathbb{Z} \\ \parallel & & & & \downarrow \\ \mathbb{Z} & \longleftarrow & 0 & \longleftarrow & 0 \end{array}$$

Proof: Note that \mathcal{K} is an essential ideal of A (i.e., every nonzero ideal of A has a nontrivial intersection with \mathcal{K}) since A is unital and A/\mathcal{K} is simple. Uniqueness follows from [ERR, Corollary 4.22]. For existence, we note that when Γ is a finite and co-irreducible graph with $|\Gamma| > 1$ and $\chi(\Gamma) = k$, all properties are met as noted in Section 2. □

When specifying the map $K_0(\mathcal{K}) \rightarrow K_0(A)$ above we let the unit of the leftmost copy of \mathbb{Z} denote the class of a minimal projection of \mathbb{Z} .

DEFINITION 3.2 *The unique C^* -algebras satisfying (1),(2) and (3) are denoted $E_{|k|+1}^{\text{sgn}(k)}$. The unique C^* -algebra satisfying (1),(2) and (3') is denoted E_1^0 . The quotient E_1^0/\mathcal{K} is denoted \mathcal{O}_1 .*

Our notation has been chosen to fit the notation E_n^k for the extension algebras of \mathcal{O}_n studied in [FL07]. With our name \mathcal{O}_1 for the appropriately chosen Kirchberg algebra, we have

$$0 \longrightarrow \mathcal{K} \xrightarrow{\iota} E_n^k \xrightarrow{\pi} \mathcal{O}_n \longrightarrow 0$$

for any $k \in \{-1, 0, 1\}$ and $n \in \mathbb{N}$, provided $k = 0$ precisely when $n = 1$.

LEMMA 3.3 *$E_n^k \cong E_{n'}^{k'}$, only when $n = n'$ and $k = k'$. $E_n^k \otimes \mathcal{K} \cong E_{n'}^{k'} \otimes \mathcal{K}$ precisely when $n = n'$.*

Proof: Since the six-term exact sequences are as specified in (3) or (3') of Theorem 3.1, stable isomorphism can only occur if $n = n'$, and hence we only need to check that for $n > 0$, we have $E_n^1 \not\cong E_n^{-1}$, yet $E_n^1 \otimes \mathcal{K} \cong E_n^{-1} \otimes \mathcal{K}$.

We note that the only two options for an isomorphism among the six-term exact sequences in this case are given as

$$\begin{array}{ccccc} \mathbb{Z} & \xrightarrow{n} & \mathbb{Z} & \longrightarrow & \mathbb{Z}/n\mathbb{Z} \\ \downarrow \pm 1 & & \downarrow \mp 1 & & \downarrow \\ \mathbb{Z} & \xrightarrow{-n} & \mathbb{Z} & \longrightarrow & \mathbb{Z}/n\mathbb{Z} \end{array}$$

and that we must choose +1 as the left most isomorphism to preserve the positive cone of $K_0(\mathcal{K})$. Thus, an isomorphism is ruled out as it would fail to send the class of the unit of E_n^1 to the unit of E_n^{-1} , but an isomorphism after stabilization is guaranteed by, e.g., [ERR09]. \square

The following result follows directly from § 2.2, § 2.4 and Theorem 3.1.

THEOREM 3.4 *When Γ is a co-irreducible graph, $C^*(A_\Gamma^\pm)$ is one of the C^* -algebras $\mathcal{T}, E_n^k, \mathcal{O}_\infty$ according to*

1. *If $|\Gamma| = 1$, $C^*(A_\Gamma^\pm) \cong \mathcal{T}$*
2. *If $1 < |\Gamma| < \infty$, $C^*(A_\Gamma^\pm) \cong E_{1+|\chi(\Gamma)|}^{\text{sgn } \chi(\Gamma)}$*
3. *If $|\Gamma| = \infty$, $C^*(A_\Gamma^\pm) \cong \mathcal{O}_\infty$*

We note that by the information already noted on the ideal structures in combination with Lemma 3.3, the C^* -algebras appearing are not mutually isomorphic, and hence we have a complete classification by the cardinality of Γ and the Euler characteristic in the co-irreducible case.

In preparation for the general case we now study isomorphisms between various tensor products amongst the relevant extension algebras and some of their quotients. For this, we will need:

THEOREM 3.5 *Let $A_i, i = 1, 2$, be unital C^* -algebras whose proper ideals are precisely given by $(0), I_i, J_i$ and $I_i \oplus J_i$. We assume that I_i and J_i are UCT Kirchberg algebras, and the quotients $A_i/(I_i \oplus J_i)$ are also UCT Kirchberg algebras.*

Let $\alpha_I : K_(I_1) \cong K_*(I_2), \alpha_J : K_*(J_1) \cong K_*(J_2), \alpha_{I \oplus J} : K_*(I_1 \oplus J_1) \cong K_*(I_2 \oplus J_2), \beta : K_*(A_1) \cong K_*(A_2), \gamma_I : K_*(A/I_1) \cong K_*(A/I_2), \gamma_J : K_*(A/J_1) \cong K_*(A/J_2)$, and $\gamma_{I \oplus J} : K_*(A_1/(I_1 \oplus J_1)) \cong K_*(A_2/(I_2 \oplus J_2))$ be isomorphisms, with β preserving the K_0 -classes of the units and $\alpha_{I \oplus J} = \alpha_I \oplus \alpha_J$ (under the canonical isomorphism $K_*(I_i \oplus J_i) \cong K_*(I_i) \oplus K_*(J_i)$).*

Furthermore, we assume that these isomorphism are compatible with the K-theoretic six term exact sequences attached to

$$\begin{aligned} 0 \rightarrow I_i \rightarrow A_i \rightarrow A_i/I_i \rightarrow 0, \quad 0 \rightarrow J_i \rightarrow A_i \rightarrow A_i/J_i \rightarrow 0, \\ 0 \rightarrow I_i \oplus J_i \rightarrow A_i \rightarrow A_i/(I_i \oplus J_i) \rightarrow 0, \quad 0 \rightarrow J_i \rightarrow A_i/I_i \rightarrow A_i/(I_i \oplus J_i) \rightarrow 0 \end{aligned}$$

and

$$0 \rightarrow I_i \rightarrow A_i/J_i \rightarrow A_i/(I_i \oplus J_i) \rightarrow 0.$$

Then there exists an isomorphism $\varphi : A_1 \cong A_2$ which induces $\alpha_I, \alpha_J, \alpha_{I \oplus J}, \beta, \gamma_I, \gamma_J$ and $\gamma_{I \oplus J}$ in K-theory.

Proof: Combine [Kir00, Folgerung 4.3] and [BK, Theorem 1.3] with [RR07, Theorem 2.1] or [ERR, Theorem 3.3]. □

LEMMA 3.6 *For every $n \geq 2$, we have $\mathcal{O}_\infty \otimes E_n^{+1} \cong \mathcal{O}_\infty \otimes E_n^{-1}$.*

Proof: Both $\mathcal{O}_\infty \otimes E_n^{+1}$ and $\mathcal{O}_\infty \otimes E_n^{-1}$ are unital C^* -algebras with unique ideal isomorphic to $\mathcal{O}_\infty \otimes \mathcal{K}$ and corresponding quotient isomorphic to $\mathcal{O}_\infty \otimes \mathcal{O}_n \cong \mathcal{O}_n$. The K-theoretic six term exact sequences for $0 \rightarrow \mathcal{O}_\infty \otimes \mathcal{K} \rightarrow \mathcal{O}_\infty \otimes E_n^{+1} \rightarrow \mathcal{O}_n \rightarrow 0$ and $0 \rightarrow \mathcal{O}_\infty \otimes \mathcal{K} \rightarrow \mathcal{O}_\infty \otimes E_n^{-1} \rightarrow \mathcal{O}_n \rightarrow 0$ look as follows:

$$\begin{array}{ccccc} \mathbb{Z} & \longrightarrow & \mathbb{Z} & \longrightarrow & \mathbb{Z}/(n-1)\mathbb{Z} \\ \uparrow & & & & \downarrow \\ 0 & \longleftarrow & 0 & \longleftarrow & 0 \end{array}$$

where $\mathbb{Z} \cong K_0(\mathcal{O}_\infty \otimes \mathcal{K})$ is generated by $[1 \otimes e]$ for a minimal projection $e \in \mathcal{K}$ and the unit 1 of \mathcal{O}_∞ , and $\mathbb{Z} \cong K_0(\mathcal{O}_\infty \otimes E_n^{\pm 1})$ is generated by the class of the unit. The only difference is that for E_n^{+1} , the homomorphism $\mathbb{Z} \rightarrow \mathbb{Z}$ is given by $z[1 \otimes e] \mapsto (n-1)z[1]$, whereas for E_n^{-1} , the homomorphism $\mathbb{Z} \rightarrow \mathbb{Z}$ is given by $z[1 \otimes e] \mapsto -(n-1)z[1]$ (for $z \in \mathbb{Z}$). We now apply [RR07, Theorem 2.2] to $I_i = \mathcal{O}_\infty \otimes \mathcal{K}, A_1 = \mathcal{O}_\infty \otimes E_n^{+1}, A_2 = \mathcal{O}_\infty \otimes E_n^{-1}, Q_i = \mathcal{O}_n$ and the homomorphisms $\alpha = -\text{id}_{K_0(\mathcal{O}_\infty \otimes \mathcal{K})}, \beta : K_0(\mathcal{O}_\infty \otimes E_n^{+1}) \rightarrow K_0(\mathcal{O}_\infty \otimes E_n^{-1}), z[1] \mapsto z[1]$ (for $z \in \mathbb{Z}$), $\gamma = \text{id}_{K_0(\mathcal{O}_n)}$. It is then obvious that all the assumptions in [RR07] are satisfied, and we conclude that $\mathcal{O}_\infty \otimes E_n^{+1} \cong \mathcal{O}_\infty \otimes E_n^{-1}$. □

Now recall that we have introduced the extension algebra E_1^0 in Theorem 3.1. The C^* -algebra $E_1^0 \otimes E_n^{+1}$ ($n \geq 2$) contains the ideal $\mathcal{K} \otimes \mathcal{K} \cong \mathcal{K}$, and we

denote the corresponding quotient by Q^+ . Obviously, the primitive ideals of Q^+ are given by $\mathcal{K} \otimes \mathcal{O}_n$, $\mathcal{O}_1 \otimes \mathcal{K}$ and $\mathcal{K} \otimes \mathcal{O}_n \oplus \mathcal{O}_1 \otimes \mathcal{K}$. From the six term exact sequence in K-theory for $0 \rightarrow \mathcal{K} \rightarrow E_1^0 \otimes E_n^{+1} \rightarrow Q^+ \rightarrow 0$, we obtain $K_0(Q^+) \cong \mathbb{Z} \cong K_1(Q^+)$, where $K_0(Q^+)$ is generated by $[1_{Q^+}]$. All this also holds for the quotient Q^- of $E_1^0 \otimes E_n^{-1}$ by the ideal $\mathcal{K} \otimes \mathcal{K} \cong \mathcal{K}$.

LEMMA 3.7 *Q^+ and Q^- are isomorphic. Moreover, there exists an automorphism of Q^+ which induces $\text{id}_{\mathbb{Z}}$ on K_0 and $-\text{id}_{\mathbb{Z}}$ on K_1 .*

Proof: The first statement is an application of Theorem 3.5 to $A_1 = Q^+$, $A_2 = Q^-$, $I_1 = \mathcal{K} \otimes \mathcal{O}_n \triangleleft Q^+$, $J_1 = \mathcal{O}_1 \otimes \mathcal{K} \triangleleft Q^+$, $I_2 = \mathcal{K} \otimes \mathcal{O}_n \triangleleft Q^-$, $J_2 = \mathcal{O}_1 \otimes \mathcal{K} \triangleleft Q^-$. Namely, it is straightforward to check that it is possible to choose $\alpha_I, \alpha_J, \alpha_{I \oplus J}, \beta, \gamma_I, \gamma_J$, and $\gamma_{I \oplus J}$ with all the desired properties in Theorem 3.5.

The second statement follows in a similar way by applying Theorem 3.5 to $A_1 = A_2 = Q^+$, $I_1 = I_2 = \mathcal{K} \otimes \mathcal{O}_n \triangleleft Q^+$, $J_1 = J_2 = \mathcal{O}_1 \otimes \mathcal{K} \triangleleft Q^+$. □

LEMMA 3.8 *For every $n \geq 2$, we have $E_1^0 \otimes E_n^{+1} \cong E_1^0 \otimes E_n^{-1}$.*

Proof: By the previous lemma, we can identify Q^+ and Q^- (we use the same notation as in the previous lemma) so that we can view $E_1^0 \otimes E_n^{+1}$ and $E_1^0 \otimes E_n^{-1}$ as extensions of Q^+ :

$$0 \rightarrow \mathcal{K} \rightarrow E_1^0 \otimes E_n^{+1} \rightarrow Q^+ \rightarrow 0 \tag{2}$$

$$0 \rightarrow \mathcal{K} \rightarrow E_1^0 \otimes E_n^{-1} \rightarrow Q^+ \rightarrow 0. \tag{3}$$

Again by the previous lemma, we can choose the identification $Q^+ \cong Q^-$ in such a way that for a fixed choice of isomorphisms $K_1(Q^+) \cong \mathbb{Z}$, $K_0(\mathcal{K}) \cong \mathbb{Z}$, the index maps for both extensions (2) and (3) coincide. Now [BD96, Theorem 2] implies that (2) and (3) give the same class in $\text{Ext}_s(Q^+)$. The reason is that $\text{Ext}(K_0(Q^+), [1_{Q^+}])$ is the trivial group as $K_0(Q^+) \cong \mathbb{Z}$ and $[1_{Q^+}]$ is a generator of $K_0(Q^+) \cong \mathbb{Z}$. So the short exact sequence in [BD96, Theorem 2] tells us that two extensions of Q^+ by \mathcal{K} give the same class in $\text{Ext}_s(Q^+)$ if their index maps coincide. But this is the case for (2) and (3) by construction. Hence $E_1^0 \otimes E_n^{+1} \cong E_1^0 \otimes E_n^{-1}$ by [JT91, § 3.2]. □

For $m, n \geq 2$, the C^* -algebra $E_m^{+1} \otimes E_n^{-1}$ contains the ideal $\mathcal{K} \otimes \mathcal{K} \cong \mathcal{K}$, and we denote the corresponding quotient by Q^{+-} . Obviously, the primitive ideals of Q^{+-} are given by $\mathcal{K} \otimes \mathcal{O}_n$, $\mathcal{O}_m \otimes \mathcal{K}$ and $\mathcal{K} \otimes \mathcal{O}_n \oplus \mathcal{O}_m \otimes \mathcal{K}$. From the six term exact sequence in K-theory for $0 \rightarrow \mathcal{K} \rightarrow E_m^{+1} \otimes E_n^{-1} \rightarrow Q^{+-} \rightarrow 0$, we obtain $K_0(Q^{+-}) \cong \mathbb{Z}/(m-1)(n-1)\mathbb{Z}$, with the class of the unit being a generator, and $K_1(Q^{+-}) \cong \{0\}$. All this also holds for the quotient Q^{-+} of $E_m^{-1} \otimes E_n^{+1}$ by the ideal $\mathcal{K} \otimes \mathcal{K} \cong \mathcal{K}$.

LEMMA 3.9 *Q^{+-} and Q^{-+} are isomorphic.*

Proof: As Lemma 3.7, this is an application of Theorem 3.5 to $A_1 = Q^{+-}$, $A_2 = Q^{-+}$, $I_1 = \mathcal{K} \otimes \mathcal{O}_n \triangleleft Q^{+-}$, $J_1 = \mathcal{O}_m \otimes \mathcal{K} \triangleleft Q^{+-}$, $I_2 = \mathcal{K} \otimes \mathcal{O}_n \triangleleft Q^{-+}$,

$J_2 = \mathcal{O}_m \otimes \mathcal{K} \triangleleft Q^{-+}$. Namely, it is straightforward to check that it is possible to choose $\alpha_I, \alpha_J, \alpha_{I \oplus J}, \beta, \gamma_I, \gamma_J$, and $\gamma_{I \oplus J}$ with all the desired properties in Theorem 3.5. \square

LEMMA 3.10 *We have $E_m^{+1} \otimes E_n^{-1} \cong E_m^{-1} \otimes E_n^{+1}$.*

Proof: By the previous lemma, we can identify Q^{+-} and Q^{-+} (using the same notation as in the previous lemma) so that we can view $E_m^{+1} \otimes E_n^{-1}$ and $E_m^{-1} \otimes E_n^{+1}$ as extensions of Q^{+-} :

$$0 \rightarrow \mathcal{K} \rightarrow E_m^{+1} \otimes E_n^{-1} \rightarrow Q^{+-} \rightarrow 0 \tag{4}$$

$$0 \rightarrow \mathcal{K} \rightarrow E_m^{-1} \otimes E_n^{+1} \rightarrow Q^{+-} \rightarrow 0. \tag{5}$$

Since $\text{Hom}(K_1(Q^{+-}), \mathbb{Z}) = \{0\}$, [BD96, Theorem 2] yields $\text{Ext}((K_0(Q^{+-}), [1]), \mathbb{Z}) \cong \text{Ext}_s(Q^{+-})$. Hence (2) and (3) give the same class in $\text{Ext}_s(Q^{+-})$. The reason is that the exact sequences in K_0 for (4) and (5) clearly give rise to the same class in $\text{Ext}((K_0(Q^{+-}), [1]), \mathbb{Z})$. Hence $E_m^{+1} \otimes E_n^{-1} \cong E_m^{-1} \otimes E_n^{+1}$ by [JT91, § 3.2]. \square

In an entirely analogous way, we get

LEMMA 3.11 *For all $m, n \geq 2$, we have $E_m^{+1} \otimes E_n^{+1} \cong E_m^{-1} \otimes E_n^{-1}$.*

4 CLASSIFICATION OF SEMIGROUP C*-ALGEBRAS

We are now ready to address the general classification problem for C*-algebras of the form $C^*(A_\Gamma^+)$. We begin with notation:

DEFINITION 4.1 *Let Γ be a graph with co-irreducible components $\Gamma_i = (V_i, E_i)$. We set*

$$t(\Gamma) = \# \{ \Gamma_i : |V_i| = 1 \}$$

$$o(\Gamma) = \# \{ \Gamma_i : |V_i| = \infty \},$$

and for every $n \in \mathbb{Z}$

$$N_n(\Gamma) = \# \{ \Gamma_i : 1 < |V_i| < \infty, \chi(\Gamma_i) = n \}.$$

THEOREM 4.2 *Let Γ and Λ be two graphs. The semigroup C*-algebras $C^*(A_\Gamma^+)$ and $C^*(A_\Lambda^+)$ of the Artin monoids for Γ and Λ are stably isomorphic if and only if the following conditions hold:*

- (i) $t(\Gamma) = t(\Lambda)$;
- (ii) $N_{-n}(\Gamma) + N_n(\Gamma) = N_{-n}(\Lambda) + N_n(\Lambda)$ for any $n \in \mathbb{Z}$;
- (iii) $\sum_{n \in \mathbb{Z}} N_n(\Gamma) = \infty$ or $\min(o(\Gamma), 1) = \min(o(\Lambda), 1)$.

They are isomorphic if and only if further

(iv) If $\sum_{n \in \mathbb{Z}} N_n(\Gamma) < \infty$, $o(\Gamma) = 0$ and $N_0(\Gamma) = 0$, then

$$\sum_{n=1}^{\infty} N_{-n}(\Gamma) \equiv \sum_{n=1}^{\infty} N_{-n}(\Lambda) \pmod{2}$$

holds.

REMARK 4.3 Note that when (ii) holds, all the conditions in (iii) are symmetric in Γ and Λ . Similarly, when (ii) and (iii) hold, so are the conditions in (iv).

For the proof of Theorem 4.2, we need some preparation. Given a graph Γ with co-irreducible components $\Gamma_i = (V_i, E_i)$, let Γ' be the graph we get from Γ by removing all the co-irreducible components Γ_i with $|V_i| = 1$ and the corresponding edges. We then have a canonical isomorphism $C^*(A_{\Gamma}^+) \cong \bigotimes_{\{\Gamma_i: |V_i|>1\}} C^*(A_{\Gamma_i}^+)$.

LEMMA 4.4 *There is a primitive ideal I' of $C^*(A_{\Gamma}^+)$ such that $\text{Prim}(C^*(A_{\Gamma}^+)/I')$ does not continuously surject onto $\text{Prim}(\mathcal{T})$ and which is minimal among all the primitive ideals having this property, and we have*

$$C^*(A_{\Gamma}^+)/I' \cong C^*(A_{\Gamma'}^+).$$

Proof: Let I be a primitive ideal of $C^*(A_{\Gamma}^+)$. As seen in Section 2, we know that I is generated by $\left\{ \bigotimes_j J_{ij} \right\}_i$, where $J_{ij} = C^*(A_{\Gamma_j}^+)$ for $i \neq j$ and $J_{ii} = I_i$ for primitive ideals I_i of $C^*(A_{\Gamma_i}^+)$. It follows that $C^*(A_{\Gamma}^+)/I \cong \bigotimes_i C^*(A_{\Gamma_i}^+)/I_i$, and hence $\text{Prim}(C^*(A_{\Gamma}^+)/I) \cong \prod_i \text{Prim}(C^*(A_{\Gamma_i}^+)/I_i)$. We now claim that there exists a continuous surjection $\text{Prim}(C^*(A_{\Gamma}^+)/I) \rightarrow \text{Prim} \mathcal{T}$ if and only if there exists a co-irreducible component Γ_i of Γ with $|V_i| = 1$ and $I_i = (0)$. The direction “ \Leftarrow ” is obvious. For “ \Rightarrow ”, assume that for every co-irreducible component Γ_i of Γ with $|V_i| = 1$, I_i is a maximal ideal of $C^*(A_{\Gamma_i}^+)$ such that $C^*(A_{\Gamma_i}^+)/I_i \cong \mathbb{C}$. Then $\text{Prim}(C^*(A_{\Gamma}^+)/I) \cong \prod_k X_k$ where $X_k = \{x_k, y_k\}$ and the open subsets of X_k are given by \emptyset , $\{x_k\}$ and X_k . This means that $\overline{\{x_k\}} = X_k$ and $\overline{\{y_k\}} = \{y_k\}$. Furthermore, we know that $\text{Prim}(\mathcal{T}) = \{\bullet\} \sqcup \mathbb{T}$, where $\{\bullet\} = \text{Prim}(\mathcal{T})$. Let $f: \prod_k X_k \rightarrow \text{Prim}(\mathcal{T})$ be a continuous map. We want to show that f cannot be surjective. Let $y = (y_k)_k$ and $f(y) = z$. For arbitrary $x \in \prod_k X_k$, we always have $y \in \overline{\{x\}}$. As $f^{-1}(\overline{\{f(x)\}})$ is closed and contains x , it must also contain y . Hence $z = f(y)$ lies in $\overline{\{f(x)\}}$. This implies that $f(x) = z$ or $f(x) = \bullet$. But this holds for every x in $\prod_k X_k$. Hence the image of f contains at most 2 points, and thus f cannot be surjective. This shows our claim.

Therefore, a primitive ideal I' of $C^*(A_{\Gamma}^+)$ such that $\text{Prim}(C^*(A_{\Gamma}^+)/I')$ does not continuously surject onto $\text{Prim}(\mathcal{T})$ and which is minimal among all the primitive ideals with this property is generated by $\left\{ \bigotimes_j J_{ij} \right\}_i$, where for a co-irreducible component Γ_i with $|V_i| = 1$, $J_{ii} = I_i$ is a maximal ideal of $C^*(A_{\Gamma_i}^+)$ with $C^*(A_{\Gamma_i}^+)/I_i \cong \mathbb{C}$, and for a co-irreducible component Γ_i with $|V_i| > 1$, $J_{ii} = (0)$. We conclude that $C^*(A_{\Gamma}^+)/I' \cong C^*(A_{\Gamma'}^+)$. \square

LEMMA 4.5 *Let Γ and Λ be two graphs.*

- (1) *If $C^*(A_\Gamma^+)$ and $C^*(A_\Lambda^+)$ are isomorphic, then $t(\Gamma) = t(\Lambda)$ and $C^*(A_{\Gamma'}^+) \cong C^*(A_{\Lambda'}^+)$.*
- (2) *If $C^*(A_\Gamma^+) \otimes \mathcal{K}$ and $C^*(A_\Lambda^+) \otimes \mathcal{K}$ are isomorphic, then $t(\Gamma) = t(\Lambda)$ and $C^*(A_{\Gamma'}^+) \otimes \mathcal{K} \cong C^*(A_{\Lambda'}^+) \otimes \mathcal{K}$.*

Proof: We first prove (1). Since an isomorphism $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$ sends the primitive ideal I to a primitive ideal of $C^*(A_\Lambda^+)$ with the analogous property, we conclude that every isomorphism $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$ induces an isomorphism $C^*(A_{\Gamma'}^+) \cong C^*(A_{\Lambda'}^+)$. To prove that $t(\Gamma) = t(\Lambda)$, we observe that the primitive ideals of $C^*(A_\Gamma^+)$ which are contained in I are in one-to-one correspondence with the subsets of $\{\Gamma_i: |V_i| = 1\}$. Again, as an isomorphism $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$ sends the primitive ideal I to a primitive ideal of $C^*(A_\Lambda^+)$ with the analogous property, we conclude that the power sets of $\{\Gamma_i: |V_i| = 1\}$ and $\{\Lambda_j: |W_j| = 1\}$ have the same cardinality. Hence also $\{\Gamma_i: |V_i| = 1\}$ and $\{\Lambda_j: |W_j| = 1\}$ must have the same cardinality (which is either finite or countably infinite). This proves (1).

(2) is proved in a similar way as (1) using the observation that every primitive ideal of $B \otimes \mathcal{K}$ is of the form $I \otimes \mathcal{K}$, where I is a primitive ideal of B . \square

LEMMA 4.6 *Let A_i , $i = 1, 2, \dots$, be a countably infinite family of properly infinite unital C^* -algebras. Then $A = \bigotimes_{i=1}^{\infty} A_i$ is purely infinite.*

Proof: We have to show that every non-zero positive element a of A is properly infinite. By [KR02, Lemma 3.3], it suffices to find for every $\varepsilon > 0$ a properly infinite, positive element $b \in A$ with $\|a - b\| < \varepsilon$ and $b \preceq a$. Since $A = \bigotimes_{i=1}^{\infty} A_i$, there exists a (sufficiently large) natural number n and a positive element $x \in \bigotimes_{i=1}^n A_i$ with $\|a - x \otimes 1\| < \frac{\varepsilon}{2}$. By [KR02, Lemma 2.2], we have that $b := (x - \frac{\varepsilon}{2})_+ \otimes 1 = (x \otimes 1 - \frac{\varepsilon}{2})_+$ satisfies $b \preceq a$. Also, we have $\|b - a\| \leq \|b - x \otimes 1\| + \|x \otimes 1 - a\| < \varepsilon$. So it suffices to show that b is properly infinite. By construction, b is of the form $c \otimes 1$ for some positive element $c \in \bigotimes_{i=1}^n A_i$. Since the unit $1 \in A_{n+1}$ is properly infinite, we can find isometries s and t in A_{n+1} with $ss^* \perp tt^*$. So $b = c \otimes 1 = (c^{1/2} \otimes s)^*(c^{1/2} \otimes s) \approx (c^{1/2} \otimes s)(c^{1/2} \otimes s)^* = c \otimes ss^*$. Similarly, $b \approx c \otimes tt^*$. But since $(c \otimes ss^*)(c \otimes tt^*) = 0$, we conclude that $b \oplus b \approx (c \otimes ss^*) \oplus (c \otimes tt^*) \approx c \otimes (ss^* + tt^*) \leq c \otimes 1 = b$. \square

LEMMA 4.7 *Let Γ be a graph with (countably) infinitely many co-irreducible components $\Gamma_i = (V_i, E_i)$, $i = 1, 2, \dots$. Assume that $1 < |V_i| < \infty$ for all i . Then $C^*(A_\Gamma^+)$ is strongly purely infinite, i.e., $C^*(A_\Gamma^+) \cong C^*(A_\Gamma^+) \otimes \mathcal{O}_\infty$.*

Proof: By [CL07, Theorem 8.3], we know that $C^*(A_\Gamma^+)$ has the ideal property (the definition can be found in [PR07, Remark 2.1]). Moreover, we know that $C^*(A_\Gamma^+) \cong \bigotimes_{i=1}^{\infty} C^*(A_{\Gamma_i}^+)$, and each of the $C^*(A_{\Gamma_i}^+)$ is a properly infinite unital C^* -algebra. Hence by the previous lemma, we know that $C^*(A_\Gamma^+)$ is purely infinite. Therefore, [PR07, Proposition 2.14] tells us that $C^*(A_\Gamma^+)$ is strongly

purely infinite. And finally, if $C^*(A_\Gamma^+)$ is strongly purely infinite, then [KR02, Theorem 9.1] implies that $C^*(A_\Gamma^+) \cong C^*(A_\Gamma^+) \otimes \mathcal{O}_\infty$ because $C^*(A_\Gamma^+)$ is nuclear and unital. \square

Finally, we are ready for the proof of Theorem 4.2. *Proof:*[Proof of Theorem 4.2] Let us first of all show that if $C^*(A_\Gamma^+) \otimes \mathcal{K} \cong C^*(A_\Lambda^+) \otimes \mathcal{K}$ holds, then conditions (i), (ii) and (iii) must be satisfied. By Lemma 4.5 condition (i) holds and that $C^*(A_{\Gamma'}^+) \otimes \mathcal{K} \cong C^*(A_{\Lambda'}^+) \otimes \mathcal{K}$. Hence we may assume that all the co-irreducible components of Γ and Λ have more than one vertex.

To prove (ii), we observe that the minimal non-zero primitive ideals of $C^*(A_\Gamma^+)$ are of the form $I_i = \otimes_j J_{ij}$, where $J_{ij} = C^*(A_{\Gamma_j}^+)$ if $j \neq i$, and $J_{ii} = \mathcal{K} \triangleleft C^*(A_{\Gamma_i}^+)$ (Γ_i consists of only finitely many vertices). For the corresponding quotient, we get $C^*(A_\Gamma^+)/I_i \cong \otimes_j Q_{ij}$, where $Q_{ij} = C^*(A_{\Gamma_j}^+)$ if $j \neq i$, and $Q_{ii} = C^*(A_{\Gamma_i}^+)/\mathcal{K}$. Since $K_0(C^*(A_{\Gamma_i}^+)/\mathcal{K}) \cong \mathbb{Z}/|\chi(\Gamma_i)|\mathbb{Z}$, it follows that $K_0(C^*(A_\Gamma^+)/I_i) \cong \mathbb{Z}/|\chi(\Gamma_i)|\mathbb{Z}$. Hence, we have shown that $N_0(\Gamma)$ is the number of minimal non-zero primitive ideals $I \otimes \mathcal{K}$ of $C^*(A_\Gamma^+) \otimes \mathcal{K}$ with the property that $K_0(C^*(A_\Gamma^+)/I) \cong \mathbb{Z}$, and that for every $n = 1, 2, \dots$, $N_{-n}(\Gamma) + N_n(\Gamma)$ is the number of minimal non-zero primitive ideals $I \otimes \mathcal{K}$ of $C^*(A_\Gamma^+) \otimes \mathcal{K}$ with the property that $K_0(C^*(A_\Gamma^+)/I) \cong \mathbb{Z}/n\mathbb{Z}$. Since these descriptions are invariant under stable isomorphisms of C^* -algebras, we conclude that (ii) must hold.

Let us now prove (iii) under the assumption of stable isomorphism. If $\sum N_n(\Gamma) = \infty$ we are done, so suppose the contrary and note that in this case, $C^*(A_\Gamma^+)$ is strongly purely infinite if and only if $o(\Gamma) > 0$. The direction “ \Rightarrow ” is clear, since $o(\Gamma) > 0$ implies that $C^*(A_\Gamma^+)$ has \mathcal{O}_∞ as a tensor factor. To prove “ \Leftarrow ”, we observe that if $o(\Gamma) = 0$, then $C^*(A_\Gamma^+)$ contains the algebra of compact operators as an ideal, hence cannot be strongly purely infinite. As a consequence, $C^*(A_\Gamma^+) \otimes \mathcal{K} \cong C^*(A_\Lambda^+) \otimes \mathcal{K}$ implies that either both $o(\Gamma) > 0$ and $o(\Lambda) > 0$, or $o(\Gamma) = o(\Lambda) = 0$, as desired.

Finally, we assume that $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$ and that $\sum N_n(\Gamma) < \infty$, that $N_0(\Gamma) = 0$ and that $o(\Gamma) = 0$. The algebra \mathcal{K} of compact operators sits inside $C^*(A_\Gamma^+)$ as the (unique) minimal non-zero ideal. The inclusion $\mathcal{K} \hookrightarrow C^*(A_\Gamma^+)$ sends in K-theory the K_0 -class of a minimal projection to $(\prod_i \chi(\Gamma_i)) \cdot [1]$, where $(\prod_i \chi(\Gamma_i))$ is the product over all co-irreducible components of Γ (there are only finitely many by assumption) of the Euler characteristics. As $N_0(\Gamma) = 0$, $(\prod_i \chi(\Gamma_i))$ is a non-zero number, and it is positive if and only if $\sum_{n=1}^\infty N_{-n}(\Gamma) \equiv 0 \pmod{2}$. Since $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$, we must have $\sum_{n=1}^\infty N_{-n}(\Gamma) \equiv \sum_{n=1}^\infty N_{-n}(\Lambda) \pmod{2}$. Therefore, all in all, condition (iv) follows when the C^* -algebras are isomorphic.

In the opposite direction, we know from Sections 2 and 3 that

$$C^*(A_\Gamma^+) \cong \mathcal{T}^{\otimes t(\Gamma)} \otimes \mathcal{O}_\infty^{\otimes o(\Gamma)} \otimes \bigotimes_{n=0}^\infty \bigotimes_{\{i: |\chi(\Gamma_i)|=n\}} E_{1+n}^{\text{sgn}(\chi(\Gamma_i))}$$

and

$$C^*(A_\Lambda^+) \cong \mathcal{T}^{\otimes t(\Lambda)} \otimes \mathcal{O}_\infty^{\otimes o(\Lambda)} \otimes \bigotimes_{n=0}^\infty \bigotimes_{\{i: |\chi(\Lambda_i)|=n\}} E_{1+n}^{\text{sgn}(\chi(\Lambda_i))}$$

We note from the outset that whenever $o(\Gamma) > 0$ or $N_0(\Gamma) > 0$ then by repeated application of either Lemma 3.6 or Lemma 3.8 we may simplify these expressions to

$$C^*(A_\Gamma^+) \cong \mathcal{T}^{\otimes t(\Gamma)} \otimes \mathcal{O}_\infty^{\otimes o(\Gamma)} \otimes (E_1^0)^{\otimes N_0(\Gamma)} \otimes \bigotimes_{n=1}^{\infty} (E_{1+n}^{+1})^{\otimes (N_{-n}(\Gamma) + N_n(\Gamma))}. \quad (6)$$

Assume that (i), (ii) and (iii) hold. We begin by noting that in the case $\sum N_n(\Gamma) < \infty$ if either $o(\Gamma) > 0$ or $N_0(\Gamma) > 0$, we also have either $o(\Lambda) > 0$ or $N_0(\Lambda) > 0$, and we get $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$ by reducing to the form given in (6) and applying (i) and (ii).

When $\sum N_n(\Gamma) = \infty$ then we have by (ii) and Lemma 4.7 that both $C^*(A_\Gamma^+)$ and $C^*(A_\Lambda^+)$ are strongly purely infinite, and hence we have

$$\begin{aligned} C^*(A_\Gamma^+) &\cong \mathcal{T}^{\otimes t(\Gamma)} \otimes \mathcal{O}_\infty \otimes (E_1^0)^{\otimes N_0(\Gamma)} \otimes \bigotimes_{n=1}^{\infty} (E_{1+n}^{+1})^{\otimes (N_{-n}(\Gamma) + N_n(\Gamma))} \\ &= \mathcal{T}^{\otimes t(\Lambda)} \otimes \mathcal{O}_\infty \otimes (E_1^0)^{\otimes N_0(\Lambda)} \otimes \bigotimes_{n=1}^{\infty} (E_{1+n}^{+1})^{\otimes (N_{-n}(\Lambda) + N_n(\Lambda))} \\ &\cong C^*(A_\Lambda^+), \end{aligned}$$

since Lemma 3.6 may be applied as above.

It remains to treat the case that $o(\Gamma) = N_0(\Gamma) = 0$ and $\sum N_n(\Gamma) < \infty$. Again, by (ii) and (iii), we must have $o(\Lambda) = N_0(\Lambda) = 0$ and $\sum N_n(\Lambda) < \infty$ as well, and we get

$$\begin{aligned} C^*(A_\Gamma^+) \otimes \mathcal{K} &\cong \mathcal{T}^{\otimes t(\Gamma)} \otimes \bigotimes_{n=1}^{\infty} (E_{1+n}^{+1})^{\otimes (N_{-n}(\Gamma) + N_n(\Gamma))} \otimes \mathcal{K} \\ &= \mathcal{T}^{\otimes t(\Lambda)} \otimes \bigotimes_{n=1}^{\infty} (E_{1+n}^{+1})^{\otimes (N_{-n}(\Lambda) + N_n(\Lambda))} \otimes \mathcal{K} \\ &\cong C^*(A_\Lambda^+) \otimes \mathcal{K} \end{aligned}$$

this time appealing to the second half of Lemma 3.3.

Assuming further (iv), we now aim for exact isomorphism, noting that we have already established it when $o(\Gamma) > 0$, $N_0(\Gamma) > 0$ or $\sum N_n(\Gamma) = \infty$. We hence assume that $o(\Gamma) = N_0(\Gamma) = 0$ and note that also $o(\Lambda) = N_0(\Lambda) = 0$

Consider first the case where both $\sum_{n=1}^\infty N_{-n}(\Gamma)$ and $\sum_{n=1}^\infty N_{-n}(\Lambda)$ are even. We have

$$\begin{aligned} C^*(A_\Gamma^+) &\cong \mathcal{T}^{\otimes t(\Gamma)} \otimes \bigotimes_{n=1}^\infty (E_{1+n}^{+1})^{\otimes N_n(\Gamma)} \otimes \bigotimes_{n=1}^\infty (E_{1+n}^{-1})^{\otimes N_{-n}(\Gamma)} \\ &\cong \mathcal{T}^{\otimes t(\Gamma)} \otimes \bigotimes_{n=1}^\infty (E_{1+n}^{+1})^{\otimes N_n(\Gamma)} \otimes \bigotimes_{n=1}^\infty (E_{1+n}^{+1})^{\otimes N_{-n}(\Gamma)} \\ &\cong \mathcal{T}^{\otimes t(\Lambda)} \otimes \bigotimes_{n=1}^\infty (E_{1+n}^{+1})^{\otimes (N_{-n}(\Gamma) + N_n(\Gamma))} \\ &= \mathcal{T}^{\otimes t(\Lambda)} \otimes \bigotimes_{n=1}^\infty (E_{1+n}^{+1})^{\otimes (N_{-n}(\Lambda) + N_n(\Lambda))} \cong C^*(A_\Lambda^+) \end{aligned}$$

by Lemma 3.11. Now assume both $\sum_{n=1}^\infty N_{-n}(\Gamma)$ and $\sum_{n=1}^\infty N_{-n}(\Lambda)$ are odd. If there exists $\chi < 0$ such that there are co-irreducible components Γ_k and Λ_l with $\chi(\Gamma_k) = \chi = \chi(\Lambda_l)$, then we deduce from the previous case that

$$\begin{aligned} C^*(A_\Gamma^+) &\cong \left(\bigotimes_{\Gamma_i \neq \Gamma_k} C^*(A_{\Gamma_i}^+) \right) \otimes C^*(A_{\Gamma_k}^+) \\ &\cong \left(\bigotimes_{\Gamma_i \neq \Gamma_k} C^*(A_{\Gamma_i}^+) \right) \otimes E_{1+|\chi|}^{-1} \\ &\cong \left(\bigotimes_{\Lambda_j \neq \Lambda_l} C^*(A_{\Lambda_j}^+) \right) \otimes C^*(A_{\Lambda_l}^+) \cong C^*(A_\Lambda^+). \end{aligned}$$

If there exists no such χ , then by (ii) there must be $\chi < 0$, $\psi < 0$ and co-irreducible components Γ_{k_-} , Γ_{k_+} , Λ_{l_-} , Λ_{l_+} with $\chi(\Gamma_{k_-}) = \chi$, $\chi(\Lambda_{l_+}) = -\chi$, $\chi(\Gamma_{k_+}) = -\psi$ and $\chi(\Lambda_{l_-}) = \psi$. Hence

$$\begin{aligned} C^*(A_\Gamma^+) &\cong \left(\bigotimes_{\Gamma_i \neq \Gamma_{k_+}, \Gamma_{k_-}} C^*(A_{\Gamma_i}^+) \right) \otimes C^*(A_{\Gamma_{k_+}}^+) \otimes C^*(A_{\Gamma_{k_-}}^+) \\ &\cong \left(\bigotimes_{\Gamma_i \neq \Gamma_{k_+}, \Gamma_{k_-}} E_{1+|\chi(\Gamma_i)|}^{\text{sgn}(\chi(\Gamma_i))} \right) \otimes E_{1+|\psi|}^{+1} \otimes E_{1+|\chi|}^{-1} \\ &\cong \left(\bigotimes_{\Gamma_i \neq \Gamma_{k_+}, \Gamma_{k_-}} E_{1+|\chi(\Gamma_i)|}^{\text{sgn}(\chi(\Gamma_i))} \right) \otimes E_{1+|\psi|}^{-1} \otimes E_{1+|\chi|}^{+1} \\ &\cong \left(\bigotimes_{\Gamma_i \neq \Gamma_{k_+}, \Gamma_{k_-}} C^*(A_{\Gamma_i}^+) \right) \otimes C^*(A_{\Lambda_{l_-}}^+) \otimes C^*(A_{\Lambda_{l_+}}^+) \cong C^*(A_\Lambda^+). \end{aligned}$$

In the third step, we used Lemma 3.10, and in the fourth step, we used our argument in the previous case. \square

5 THE ISOMORPHISM PROBLEM FROM THE PERSPECTIVE OF CLASSIFICATION OF NON-SIMPLE C^* -ALGEBRAS

We give an interpretation of Theorem 4.2 from the point of view of classifying non-simple C^* -algebras.

We let $\mathcal{O}(\text{Prim}(A))$ denote the set of open subsets in $\text{Prim}(A)$, and $\mathbb{I}(A)$ the lattice of ideals.

A lattice map $\psi_A: \mathcal{O}(\text{Prim}(A)) \rightarrow \mathbb{I}(A)$ given by $\psi_A(U) = \bigcap_{\rho \notin U} \rho$ is then a lattice isomorphism which preserves arbitrary suprema and finite infima. We denote $\psi_A(U)$ by $A[U]$. For every C^* -algebra A , we denote the pair

$$\left(\text{Prim}(A), \{K_{\text{six}}^+(A/A[U]; A[V]/A[U])\}_{V, U \in \mathcal{O}(\text{Prim}(A))} \right)$$

by $\mathfrak{F}(A)$, where $K_{\text{six}}^+(B, J)$ denotes the standard six-term exact sequence associated to an ideal J of a C^* -algebra B , considering each K_0 -group as an ordered group.

An isomorphism from $\mathfrak{F}(A)$ to $\mathfrak{F}(B)$ thus consists of a homeomorphism

$$\phi: \text{Prim}(A) \rightarrow \text{Prim}(B)$$

and isomorphisms

$$\alpha_{U, V}: K_*(A[V]/A[U]) \rightarrow K_*(B[\phi(V)]/B[\phi(U)])$$

for each $U, V \in \mathcal{O}(\text{Prim}(A))$ with $U \subseteq V$, such that $(\alpha_{U, V}, \alpha_{X, U}, \alpha_{X, V})$ is an isomorphism from $K_{\text{six}}^+(A/A[U]; A[V]/A[U])$ to

$$K_{\text{six}}^+(B/B[\phi(U)]; B[\phi(V)]/B[\phi(U)])$$

in the sense that it makes all squares commute and is an order isomorphism on all even parts of the K -theory.

If A and B are unital, we write $(\mathfrak{F}(A), [1_A]) \cong (\mathfrak{F}(B), [1_B])$ if $\mathfrak{F}(A) \cong \mathfrak{F}(B)$ in such a way that the isomorphism $\alpha_{X, \emptyset}$ sends $[1_A]$ in $K_0(A)$ to $[1_B]$ in $K_0(B)$. Note that if $\phi: \text{Prim}(A) \rightarrow \text{Prim}(B)$ is a homeomorphism, there exists a lattice isomorphism from $\mathbb{I}(A)$ to $\mathbb{I}(B)$ given by $I \mapsto \psi_B(\phi(\psi_A^{-1}(I)))$. Hence, if A and B are separable and $\phi: \text{Prim}(A) \rightarrow \text{Prim}(B)$ is a homeomorphism, then for all $U \in \mathcal{O}(\text{Prim}(A))$, we have that $A[U]$ is a primitive ideal of A if and only if $B[\phi(U)]$ is a primitive ideal of B (because primitive ideals are precisely given by prime ideals for separable C^* -algebras).

The following easy observation is left to the reader.

LEMMA 5.1 *Let A and B be separable C^* -algebras. Let $U \in \mathcal{O}(\text{Prim}(A))$.*

(1) If $\mathfrak{F}(A) \cong \mathfrak{F}(B)$ via a homeomorphism $\phi: \text{Prim}(A) \rightarrow \text{Prim}(B)$, then

$$\mathfrak{F}(A/A[U]) \cong \mathfrak{F}(B/B[\phi(U)]).$$

(2) If A and B are unital C^* -algebras and $(\mathfrak{F}(A), [1_A]) \cong (\mathfrak{F}(B), [1_B])$ via a homeomorphism $\phi: \text{Prim}(A) \rightarrow \text{Prim}(B)$, then

$$(\mathfrak{F}(A/A[U]), [1_{A/A[U]}]) \cong (\mathfrak{F}(B/B[\phi(U)]), [1_{B/B[\phi(U)}]).$$

THEOREM 5.2 *Let Γ and Λ be two (countable) graphs, and let $\Gamma_i = (V_i, E_i)$ be the co-irreducible components of Γ , $\Lambda_j = (W_j, F_j)$ the co-irreducible components of Λ . Then $C^*(A_\Gamma^+) \cong C^*(A_\Lambda^+)$ if and only if $(\mathfrak{F}(C^*(A_\Gamma^+)), [1_{C^*(A_\Gamma^+)}]) \cong (\mathfrak{F}(C^*(A_\Lambda^+)), [1_{C^*(A_\Lambda^+)}])$.*

Proof: The direction “ \Rightarrow ” is obvious. To prove “ \Leftarrow ”, we show that

$$(\mathfrak{F}(C^*(A_\Gamma^+)), [1_{C^*(A_\Gamma^+)}]) \cong (\mathfrak{F}(C^*(A_\Lambda^+)), [1_{C^*(A_\Lambda^+)}])$$

implies (i), (ii), (iii) and (iv) from Theorem 4.2, using the notations from Lemma 4.5. The first step is to prove that $(\mathfrak{F}(C^*(A_\Gamma^+)), [1_{C^*(A_\Gamma^+)}]) \cong (\mathfrak{F}(C^*(A_\Lambda^+)), [1_{C^*(A_\Lambda^+)}])$ implies $t(\Gamma) = t(\Lambda)$ and $(\mathfrak{F}(C^*(A_{\Gamma'}^+)), [1_{C^*(A_{\Gamma'}^+)}]) \cong (\mathfrak{F}(C^*(A_{\Lambda'}^+)), [1_{C^*(A_{\Lambda'}^+)}])$. $t(\Gamma) = t(\Lambda)$ follows by Lemma 4.4, because we only use the primitive ideal space and the lattice structure of the set of ideals in this proof. To see that $(\mathfrak{F}(C^*(A_{\Gamma'}^+)), [1_{C^*(A_{\Gamma'}^+)}]) \cong (\mathfrak{F}(C^*(A_{\Lambda'}^+)), [1_{C^*(A_{\Lambda'}^+)}])$, let I' be a primitive ideal of $C^*(A_\Gamma^+)$ stipulated in Lemma 4.4, and let U be an open set of $\text{Prim}(C^*(A_\Gamma^+))$ such that $C^*(A_\Gamma^+)[U] = I'$. Then $C^*(A_\Lambda^+)[\phi(U)]$ is an ideal with the analogous property. In the proof of Lemma 4.5, we have seen that $C^*(A_\Gamma^+)/C^*(A_\Gamma^+)[U] \cong C^*(A_{\Gamma'}^+)$. Similarly, we have $C^*(A_\Lambda^+)/C^*(A_\Lambda^+)[\phi(U)] \cong C^*(A_{\Lambda'}^+)$. Therefore, (2) from the previous lemma tells us that $(\mathfrak{F}(C^*(A_{\Gamma'}^+)), [1_{C^*(A_{\Gamma'}^+)}]) \cong (\mathfrak{F}(C^*(A_{\Lambda'}^+)), [1_{C^*(A_{\Lambda'}^+)}])$, as desired.

In particular, this implies (i), and we may assume as in the proof of Theorem 4.2 that all the co-irreducible components of Γ and Λ have more than one vertex. Then (ii) follows in exactly the same way as in the proof of Theorem 4.2 because we only use primitive ideal spaces, lattice structures of sets of ideals and K_0 in this proof. All this can be extracted from the invariant \mathfrak{F} . Let us prove (iii). As we have seen in the proof of Theorem 4.2, $o(\Gamma) = 0$ implies that \mathcal{K} is an ideal of $C^*(A_\Gamma^+)$, whereas $o(\Gamma) > 0$ implies that $C^*(A_\Gamma^+)$ (and hence also every non-zero ideal) is strongly purely infinite. These two cases can be distinguished by the order on K_0 . Therefore, we see as in the proof of Theorem 4.2 that if $(\mathfrak{F}(C^*(A_\Gamma^+)), [1_{C^*(A_\Gamma^+)}]) \cong (\mathfrak{F}(C^*(A_\Lambda^+)), [1_{C^*(A_\Lambda^+)}])$, then either $o(\Gamma) > 0$ and $o(\Lambda) > 0$ or $o(\Gamma) = 0$ and $o(\Lambda) = 0$. The proof of (iv) then follows the proof of Theorem 4.2, where we only use lattice structures of sets of ideals, K_0 and the K_0 -classes of the units. All this can be extracted from the invariant \mathfrak{F} together with the position of the K_0 -class of the unit. \square

6 GRAPH ALGEBRAS AND THE SEMIPROJECTIVITY QUESTION

Apart from semigroup C^* -algebras we discussed above, there is another - more traditional - way of constructing a C^* -algebra out of a directed graph, possibly allowing for loops. Now we would like to discuss the overlap of these two constructions. In other words, we are interested in the question: Which semigroup C^* -algebras for right-angled Artin monoids are isomorphic to graph algebras? We can provide a complete answer to this question.

6.1 EXTENSIONS OF C^* -ALGEBRAS

We first establish some facts about absorbing extensions and the C^* -algebras associated to these extensions. To each injective Busby map $\tau: A \rightarrow \mathcal{Q}(B)$, where $\mathcal{Q}(B) = \mathcal{M}(B)/B$ with $\mathcal{M}(B)$ the multiplier algebra of B , associate as usual the extension

$$e : \begin{array}{ccccccc} 0 & \longrightarrow & B^C & \longrightarrow & E & \xrightarrow{\psi} & A & \longrightarrow & 0 \\ & & \parallel & & \downarrow & & \downarrow \tau & & \\ 0 & \longrightarrow & B^C & \longrightarrow & \mathcal{M}(B) & \xrightarrow{\pi} & \mathcal{Q}(B) & \longrightarrow & 0 \end{array}$$

with $E = \pi^{-1}(\tau(A))$ and $\psi(x) = \tau^{-1}(\pi(x))$. Note that ψ is a homomorphism since τ is injective.

We call τ (and e) *unital* if A is unital and τ is a unital homomorphism, or, equivalently, if E is a unital C^* -algebra. If $\tau = \pi \circ \alpha$ for some homomorphism $\alpha: A \rightarrow \mathcal{M}(B)$, then τ is called a *trivial extension*. If A is unital and $\tau = \pi \circ \alpha$ for some unital homomorphism $\alpha: A \rightarrow \mathcal{M}(B)$, then τ is called *strongly unital*. Not all unital trivial extensions are strongly unital.

Assume that B is stable. The sum $\tau \oplus \tau'$ of two extensions $\tau, \tau': A \rightarrow \mathcal{Q}(B)$ is defined as follows. Since B is stable, there exist isometries $s_1, s_2 \in \mathcal{M}(B)$ with $1_{\mathcal{M}(B)} = s_1 s_1^* + s_2 s_2^*$. Set

$$(\tau \oplus \tau')(a) = \pi(s_1)\tau(a)\pi(s_1^*) + \pi(s_2)\tau'(a)\pi(s_2^*)$$

for all $a \in A$.

Two extensions $\tau, \tau': A \rightarrow \mathcal{Q}(B)$ are said to be *unitarily equivalent*, denoted by $\tau \sim_u \tau'$, if there exists a unitary $u \in \mathcal{M}(B)$ such that $\pi(u)\tau(a)\pi(u)^* = \tau'(a)$ for all $a \in A$. Then two extensions $\tau_1, \tau_2: A \rightarrow \mathcal{Q}(B)$ define the same element in $\text{Ext}(A, B)$ if there exists a unitary $u \in \mathcal{M}(B)$ and there exist trivial extensions $\tau'_1, \tau'_2: A \rightarrow \mathcal{Q}(B)$ such that $\tau_1 \oplus \tau'_1 \sim_u \tau_2 \oplus \tau'_2$. If τ_1 and τ_2 are unital extensions, then τ'_1 and τ'_2 can be chosen to be unital extensions (see [Ror97, Section 5]).

For a C^* -algebra C , we let \tilde{C} be the unitization of C (adding a new unit if C is a unital C^* -algebra) and let $\iota_C: C \rightarrow \tilde{C}$ be the embedding of C into \tilde{C} as an ideal.

Recall that an ideal I of a C^* -algebra A is an *essential ideal* if every nonzero ideal of A has a nontrivial intersection with I . An extension $0 \rightarrow I \xrightarrow{\iota} A \rightarrow$

$B \rightarrow 0$ is *essential* if $\iota(I)$ is an essential ideal of A . It is a well-known fact that an extension $0 \rightarrow I \rightarrow A \rightarrow B \rightarrow 0$ is an essential extension if and only if the Busby invariant of the extension is injective. We now prove in the following proposition that every essential extension $0 \rightarrow B \rightarrow E \rightarrow A \rightarrow 0$ with A a non-unital, separable, nuclear C^* -algebra and B a C^* -algebra that is isomorphic to either \mathcal{K} or a nuclear, purely infinite simple C^* -algebra is absorbing.

Before proving the proposition, we show that any absorbing extension must be an essential extension. Hence, the assumption that the extension is essential is necessary. Note that if τ or τ' is injective, then the sum $\tau \oplus \tau'$ is injective. Since B is stable, there exists a unital embedding from \mathcal{O}_2 to $\mathcal{M}(B)$ which induces a unital embedding from \mathcal{O}_2 to $\mathcal{Q}(B)$. Nuclearity of A gives us an embedding of A into \mathcal{O}_2 , thus the composition gives a trivial essential extension $\tau_0: A \rightarrow \mathcal{Q}(B)$. Therefore, an absorbing extension τ is an essential extension since τ is unitarily equivalent to $\tau \oplus \tau_0$.

PROPOSITION 6.1 *Let A be a non-unital, separable, nuclear C^* -algebra and let B be a separable C^* -algebra that is isomorphic to either \mathcal{K} or a nuclear, purely infinite simple C^* -algebra. If $\tau: A \rightarrow \mathcal{Q}(B)$ is an essential extension, then for every trivial extension $\tau_0: A \rightarrow \mathcal{Q}(B)$ we have that $\tau \sim_u \tau \oplus \tau_0$. Consequently, if $e_i: 0 \rightarrow B \rightarrow E_i \rightarrow A \rightarrow 0$ is an essential extension for $i = 1, 2$ and $[\tau_{e_1}] = [\tau_{e_2}]$ in $\text{Ext}(A, B)$, then $E_1 \cong E_2$.*

Proof: Let $\alpha_0: A \rightarrow \mathcal{M}(B)$ be a homomorphism with $\tau_0 = \pi \circ \alpha_0$. Extend τ and α_0 to the unitization of \tilde{A} , and denote these extensions by $\tilde{\tau}: \tilde{A} \rightarrow \mathcal{Q}(B)$ and $\tilde{\alpha}_0: \tilde{A} \rightarrow \mathcal{M}(B)$ respectively.

We claim that $\tilde{\tau}$ is injective. Let $y \in \ker(\tilde{\tau})$. Then $\tau(yx) = \tilde{\tau}(y)\tilde{\tau}(\iota_A(x)) = 0$ and $\tau(xy) = \tilde{\tau}(\iota_A(x))\tilde{\tau}(y) = 0$ for all $x \in A$. Since τ is injective, we have that $yx = xy = 0$ for all $x \in A$. Since A is non-unital, A is an essential ideal of \tilde{A} . Hence, $y = 0$. Thus, proving our claim.

Set $E = \pi^{-1}(\tilde{\tau}(\tilde{A})) \subseteq \mathcal{M}(B)$. Since $\tilde{\tau}$ is injective, we may define a surjective homomorphism $\psi: E \rightarrow \tilde{A}$ by $\psi(x) = \tilde{\tau}^{-1}(\pi(x))$. Define $\eta: E \rightarrow \mathcal{M}(B)$ by $\eta(x) = \tilde{\alpha}_0 \circ \psi(x)$. Then η is a unital homomorphism such that $\eta(E \cap B) = \{0\}$. Let s_1 and s_2 be isometries such that $1_{\mathcal{M}(B)} = s_1s_1^* + s_2s_2^*$. By [Arv77, Corollary 2] and [Kir, Proposition 7], there exists a unitary $u \in \mathcal{M}(B)$ such that $u(s_1xs_1^* + s_2\eta(x)s_2^*)u^* - x \in B$ for all $x \in E$.

We claim u implements a unitary equivalence between τ and $\tau \oplus \tau_0$. Let $a \in A$. Choose $x \in E$ such that $\pi(x) = (\tilde{\tau} \circ \iota_A)(a)$. Note that

$$\pi \circ \eta(x) = \pi \circ \tilde{\alpha}_0 \circ \psi(x) = (\pi \circ \tilde{\alpha}_0)(\tilde{\tau}^{-1}(\pi(x))) = (\pi \circ \tilde{\alpha}_0 \circ \iota_A)(a).$$

Then

$$\begin{aligned}
& \pi(u) (\tau(a) \oplus \tau_0(a)) \pi(u)^* \\
&= \pi(u) (\pi(s_1) \tilde{\tau}(\iota_A(a)) \pi(s_1^*) + \pi(s_2) (\pi \circ \tilde{\alpha}_0 \circ \iota_A)(a) \pi(s_2^*)) \pi(u)^* \\
&= \pi(u (s_1 x s_1^* + s_2 \eta(x) s_2^*) u^*) \\
&= \pi(x) \\
&= (\tilde{\tau} \circ \iota_A)(a) \\
&= \tau(a).
\end{aligned}$$

Hence, $\tau \oplus \tau_0 \sim_u \tau$, proving the first part of the proposition.

Suppose $e_i : 0 \rightarrow B \rightarrow E_i \rightarrow A \rightarrow 0$ is an essential extension for $i = 1, 2$ and $[\tau_{e_1}] = [\tau_{e_2}]$ in $\text{Ext}(A, B)$. By the discussion before the proposition, there exist trivial extensions $\tau'_1, \tau'_2 : A \rightarrow \mathcal{Q}(B)$ such that $\tau_{e_1} \oplus \tau'_1 \sim_u \tau_{e_2} \oplus \tau'_2$. By the first part of the proposition, we have that $\tau_{e_1} \sim_u \tau_{e_1} \oplus \tau'_1$ and $\tau_{e_2} \oplus \tau'_2 \sim_u \tau'_{e_2}$. Therefore, $\tau_{e_1} \sim_u \tau_{e_2}$. By [JT91, § 3.2], $E_1 \cong E_2$. \square

6.2 CORNERS OF GRAPH ALGEBRAS

We also need some results involving corners of graph algebras. The general case will be worked out in [AGR]. For the convenience of the reader, we will prove the case that will suit our purposes (see Proposition 6.2).

Recall that if $E = (E^0, E^1, r, s)$ is a graph, the C^* -algebra $C^*(E)$ associated to E is the universal C^* -algebra generated by $\{p_v : v \in E^0\} \sqcup \{s_e : e \in E^1\}$ subject to the relations

- (i) $p_v p_w = \delta_{v,w} p_v$ for all $v, w \in E^0$;
- (ii) $s_e^* s_f = \delta_{e,f} p_{r(e)}$ for all $e, f \in E^1$;
- (iii) $s_e s_e^* \leq p_{s(e)}$ for all $e \in E^1$; and
- (iv) $p_v = \sum_{e \in s^{-1}(v)} s_e s_e^*$ for all $v \in E^0$ with $0 < |s^{-1}(v)| < \infty$.

A *loop* in E is a path $\alpha = e_1 \cdots e_n$ with $s(e_1) = s(e_n)$ and we say that $s(e_1)$ is the *base point* of α . A *simple loop* in E is a loop $\alpha = e_1 \cdots e_n$ such that $s(e_i) \neq s(e_j)$ for $i \neq j$. We say that E satisfies *Condition (K)* if every vertex is either the base point of at least two simple loops or is not the base point of a loop. It is well-known that if A is a Cuntz-Krieger algebra, then A is isomorphic to $C^*(E)$, where E is a finite graph with no sinks. If, in addition, A is purely infinite, then E will also satisfy Condition (K).

PROPOSITION 6.2 *Let E be a graph with finitely many vertices. Suppose there exists a vertex w in E such that*

- (i) $\{w\}$ is a hereditary and saturated subset of E^0 ;
- (ii) $|\{e \in E^1 : s(e) = w\}|$ is either equal to 0 or ∞ ;

- (iii) for every $v \in E^0 \setminus \{w\}$, there are finitely many edges from v to w and there exists at least one $v \in E^0 \setminus \{w\}$ such that there exists an edge from v to w ; and
- (iv) every vertex $v \in E^0 \setminus \{w\}$ emits finitely many edges and is the base point of at least two loops of length one.

Then for every full projection $p \in C^*(E) \otimes \mathcal{K}$, we have that $p(C^*(E) \otimes \mathcal{K})p$ is isomorphic to a graph algebra. Consequently, if A is a unital C^* -algebra such that $A \otimes \mathcal{K} \cong C^*(E) \otimes \mathcal{K}$, then A is isomorphic to a graph algebra.

Proof: Let $\{e_{ij}\}$ be a system of matrix units for \mathcal{K} . Throughout the proof, if p is a projection in $C^*(E)$ and $n \in \mathbb{N}$, then set $np = \overbrace{p \oplus \cdots \oplus p}^n$ in $C^*(E) \otimes \mathcal{K}$. Let $\{p_v, s_e : v \in E^0, e \in E^1\}$ be a Cuntz-Krieger E -family generating $C^*(E)$. Since the only vertex in E that is a singular vertex, i.e., emits no edges or infinitely many edges, is w , by [HL+14, Theorem 3.4 and Corollary 3.5],

$$p \sim \left(\bigoplus_{v \in S} n_v p_v \right) \oplus n_1 \left(p_w - \sum_{e \in T_1} s_e s_e^* \right) \oplus \cdots \oplus n_k \left(p_w - \sum_{e \in T_k} s_e s_e^* \right), \quad (7)$$

where $n_v > 0$ for all $v \in S$, $n_i \geq 0$ for all i , $S \subseteq E^0 \setminus \{w\}$, and T_i is a finite (possibly empty) subset of $s^{-1}(w)$ for all i . Arguing as in [AR15, Lemma 4.6], we have that the projection on the right hand side of (7) is Murray-von Neumann equivalent to

$$q = \bigoplus_{v \in E^0} m_v p_v$$

where $m_v > 0$ for all $v \in E^0$. We use the fact that if S is a finite subset of $s^{-1}(w)$, then

$$|S|p_w \oplus \left(p_w - \sum_{e \in S} s_e s_e^* \right) \sim \left(\sum_{e \in S} s_e s_e^* \right) \oplus \left(p_w - \sum_{e \in S} s_e s_e^* \right) \sim p_w$$

and the fact that if $v_0 \in E^0 \setminus \{w\}$ with $s^{-1}(v_0) \cap r^{-1}(w) \neq \emptyset$, then for any n , we have that $p_{v_0} \sim np_w \oplus p_{v_0} \oplus \left(\bigoplus_{v \in E^0} m'_v p_v \right)$ for $m'_v \geq 0$. Now, arguing as in [AR15, Proposition 4.7], we have that $q(C^*(E) \otimes \mathcal{K})q$ is isomorphic to a graph algebra. Since $p \sim q$, we have that $p(C^*(E) \otimes \mathcal{K})p \cong q(C^*(E) \otimes \mathcal{K})q$. Therefore, $p(C^*(E) \otimes \mathcal{K})p$ is isomorphic to a graph algebra.

For the last part of the proposition note that $A \cong p(C^*(E) \otimes \mathcal{K})p$, where p is the projection given by the image of $1_A \otimes e_{11}$ under some isomorphism from $A \otimes \mathcal{K}$ to $C^*(E) \otimes \mathcal{K}$. Since $1_A \otimes e_{11}$ is full in $A \otimes \mathcal{K}$, we have that p is full in $C^*(E) \otimes \mathcal{K}$. □

6.3 SEMIGROUP C*-ALGEBRAS AND GRAPH ALGEBRAS

We now determine when a C*-algebra associated to an Artin monoid is isomorphic to a graph algebra. To do this, we need to determine when an extension of two graph algebras is isomorphic to a graph algebra. In spite of substantial effort the extension problem for graph algebras has not been completely resolved even for the single non-trivial ideal case. Moreover, the results in the literature are not sufficient for our purposes. The following ad hoc result will give us what we need.

LEMMA 6.3 *For each i , let A_i be a separable, nuclear C*-algebra with an essential ideal I_i such that I_i is isomorphic to either \mathcal{K} or a purely infinite simple C*-algebra with trivial K_1 group, A_i/I_i satisfies the Universal Coefficient Theorem, and $K_1(A_i/I_i) = \{0\}$ or $K_0(A_i/I_i)$ is a free group (possibly $K_0(A_i/I_i) = \{0\}$). Suppose there exist isomorphisms $\beta: I_1 \otimes \mathcal{K} \rightarrow I_2 \otimes \mathcal{K}$, $\alpha: (A_1/I_1) \otimes \mathcal{K} \rightarrow (A_2/I_2) \otimes \mathcal{K}$, and $\eta_*: K_*(A_1 \otimes \mathcal{K}) \rightarrow K_*(A_2 \otimes \mathcal{K})$ such that $(K_*(\beta), \eta_*, K_*(\alpha)): K_{\text{six}}(A_1 \otimes \mathcal{K}; I_1 \otimes \mathcal{K}) \rightarrow K_{\text{six}}(A_2 \otimes \mathcal{K}; I_2 \otimes \mathcal{K})$ is an isomorphism. Then $A_1 \otimes \mathcal{K} \cong A_2 \otimes \mathcal{K}$.*

Proof: Let $e_1: 0 \rightarrow I_2 \otimes \mathcal{K} \rightarrow B_1 \rightarrow (A_1/I_1) \otimes \mathcal{K} \rightarrow 0$ be the extension obtained by pushing forward the extension $0 \rightarrow I_1 \otimes \mathcal{K} \rightarrow A_1 \otimes \mathcal{K} \rightarrow (A_1/I_1) \otimes \mathcal{K} \rightarrow 0$ via the isomorphism β and let $e_2: 0 \rightarrow I_2 \otimes \mathcal{K} \rightarrow B_2 \rightarrow (A_1/I_1) \otimes \mathcal{K} \rightarrow 0$ be the extension obtained by pulling back the extension $0 \rightarrow I_2 \otimes \mathcal{K} \rightarrow A_2 \otimes \mathcal{K} \rightarrow (A_2/I_2) \otimes \mathcal{K} \rightarrow 0$ via the isomorphism α . Note that there exist isomorphisms $\phi_1: A_1 \otimes \mathcal{K} \rightarrow B_1$ and $\phi_2: B_2 \rightarrow A_2 \otimes \mathcal{K}$ such that $(K_*(\beta), K_*(\phi_1), K_*(\text{id}_{(A_1/I_1) \otimes \mathcal{K}})): K_{\text{six}}(A_1 \otimes \mathcal{K}; I_1 \otimes \mathcal{K}) \rightarrow K_{\text{six}}(B_1; I_2 \otimes \mathcal{K})$ and $(K_*(\text{id}_{I_2 \otimes \mathcal{K}}), K_*(\phi_2), K_*(\alpha)): K_{\text{six}}(B_2; I_2 \otimes \mathcal{K}) \rightarrow K_{\text{six}}(A_2 \otimes \mathcal{K}; I_2 \otimes \mathcal{K})$ are isomorphisms. Then $(K_*(\text{id}_{I_2 \otimes \mathcal{K}}), K_*(\phi_2^{-1}) \circ \eta_* \circ K_*(\phi_1^{-1}), K_*(\text{id}_{(A_1/I_1) \otimes \mathcal{K}}))$ is an isomorphism from $K_{\text{six}}(B_1; I_2 \otimes \mathcal{K})$ to $K_{\text{six}}(B_2; I_2 \otimes \mathcal{K})$.

We claim that $[\tau_{e_1}] = [\tau_{e_2}]$ in $\text{Ext}((A_1/I_1) \otimes \mathcal{K}, I_2 \otimes \mathcal{K})$. Since A_1/I_1 satisfies the Universal Coefficient Theorem, we may identify $\text{Ext}((A_1/I_1) \otimes \mathcal{K}, I_2 \otimes \mathcal{K})$ with $\text{KK}^1((A_1/I_1) \otimes \mathcal{K}, I_2 \otimes \mathcal{K})$. Note that $\text{Ext}_{\mathbb{Z}}^1(K_1((A_1/I_1) \otimes \mathcal{K}), K_1(I_2 \otimes \mathcal{K})) = \{0\}$ since $K_1(I_2) = \{0\}$. Suppose $K_1(A_1/I_1) = \{0\}$. Then $K_1(\tau_{e_i}) = \{0\}$. Since $K_1(I_2) = \{0\}$, we have that $K_0(\tau_{e_i}) = \{0\}$. Hence, $K_*(\tau_{e_i}) = \{0\}$. By the Universal Coefficient Theorem, $[\tau_{e_i}]$ can be identified with the element in $\text{Ext}_{\mathbb{Z}}^1(K_0((A_1/I_1) \otimes \mathcal{K}), K_0(I_2 \otimes \mathcal{K}))$ given by $K_{\text{six}}(B_i; I_2 \otimes \mathcal{K})$. Since $(K_*(\text{id}_{I_2 \otimes \mathcal{K}}), K_*(\phi_2^{-1}) \circ \eta_* \circ K_*(\phi_1^{-1}), K_*(\text{id}_{(A_1/I_1) \otimes \mathcal{K}}))$ is an isomorphism from $K_{\text{six}}(B_1; I_2 \otimes \mathcal{K})$ to $K_{\text{six}}(B_2; I_2 \otimes \mathcal{K})$ we have that $K_{\text{six}}(B_1; I_2 \otimes \mathcal{K})$ and $K_{\text{six}}(B_2; I_2 \otimes \mathcal{K})$ induce the same element in $\text{Ext}_{\mathbb{Z}}^1(K_0((A_1/I_1) \otimes \mathcal{K}), K_0(I_2 \otimes \mathcal{K}))$. Hence, $[\tau_{e_1}] = [\tau_{e_2}]$ in $\text{Ext}((A_1/I_1) \otimes \mathcal{K}, I_2 \otimes \mathcal{K})$. Suppose $K_0(A_1/I_1)$ is a free group (possibly the zero group). By the Universal Coefficient, $[\tau_{e_i}]$ is completely determined by $K_*(\tau_{e_i})$. Since $(K_*(\text{id}_{I_2 \otimes \mathcal{K}}), K_*(\phi_2^{-1}) \circ \eta_* \circ K_*(\phi_1^{-1}), K_*(\text{id}_{(A_1/I_1) \otimes \mathcal{K}}))$ is an isomorphism from $K_{\text{six}}(B_1; I_2 \otimes \mathcal{K})$ to $K_{\text{six}}(B_2; I_2 \otimes \mathcal{K})$, we have that $K_*(\tau_{e_1}) = K_*(\tau_{e_2})$. Hence, $[\tau_{e_1}] = [\tau_{e_2}]$ in $\text{Ext}((A_1/I_1) \otimes \mathcal{K}, I_2 \otimes \mathcal{K})$.

In both cases, we have shown that $[\tau_{e_1}] = [\tau_{e_2}]$ in $\text{Ext}((A_1/I_1) \otimes \mathcal{K}, I_2 \otimes \mathcal{K})$, proving our claim. By Proposition 6.1, we have that $B_1 \cong B_2$. Therefore, $A_1 \otimes \mathcal{K} \cong A_2 \otimes \mathcal{K}$. \square

LEMMA 6.4 *Let A be a unital, separable, nuclear C^* -algebra with an essential ideal I such that $I \cong \mathcal{K}$ or $I \cong \mathcal{O}_\infty \otimes \mathcal{K}$ and A/I is isomorphic to a purely infinite Cuntz-Krieger algebra. If $K_1(A/I) = \{0\}$ or $K_0(A/I)$ is a free group (possibly $K_0(A/I) = \{0\}$), then A is isomorphic to a graph algebra.*

Proof: By [ABK14, Theorem 4.4] and [Res06, Proposition 8.3], there exists a finite graph F such that each vertex of F is the base point of at least two loop of length one and there exists an isomorphism $\phi: C^*(F) \otimes \mathcal{K} \rightarrow A/I \otimes \mathcal{K}$. Let $\psi: C^*(G) \otimes \mathcal{K} \rightarrow I$ be an isomorphism such that $K_*(\psi) = \text{id}$, where G is the graph $\{v\}$ with one vertex and no edges if $I \cong \mathcal{K}$ and G is the graph with one vertex $\{v\}$ with infinitely many edges when $I \cong \mathcal{O}_\infty \otimes \mathcal{K}$. By [EKTW, Lemma 5.2(r1) and Proposition 5.5], there exists a graph E with the properties that

- (1) $E^0 = G^0 \sqcup F^0$,
- (2) E^1 is the union of G^1 and F^1 together with a finite nonzero number of edges from each $w \in F^0$ to v , and
- (3) there exist an isomorphism $\alpha_*: K_*(C^*(E)) \rightarrow K_*(A)$ with the property that $(K_*(\psi), \alpha_*, K_*(\phi))$ is an isomorphism from $K_{\text{six}}(C^*(E); I_{\{v\}})$ to $K_{\text{six}}(A; I)$.

Note that $I_{\{v\}} \otimes \mathcal{K}$ is an essential ideal of $C^*(E) \otimes \mathcal{K}$ and there exist an isomorphism $\bar{\alpha}_*: K_*(C^*(E) \otimes \mathcal{K}) \rightarrow K_*(A \otimes \mathcal{K})$ such that $(K_*(\psi \otimes \text{id}_{\mathcal{K}}), \bar{\alpha}_*, K_*(\phi \otimes \text{id}_{\mathcal{K}}))$ is an isomorphism from $K_{\text{six}}(C^*(E) \otimes \mathcal{K}; I_{\{v\}} \otimes \mathcal{K})$ to $K_{\text{six}}(A \otimes \mathcal{K}; I \otimes \mathcal{K})$. Also, note that $I \cong I_{\{v\}} = \mathcal{K}$ or $I \cong I_{\{v\}} \cong \mathcal{O}_\infty \otimes \mathcal{K}$. By Lemma 6.3, $A \otimes \mathcal{K} \cong C^*(E) \otimes \mathcal{K}$. Therefore, A is isomorphic to a graph algebra by Proposition 6.2. \square

LEMMA 6.5 *For each $m \in \mathbb{N}$, for each $n \geq 0$, the smallest nonzero ideal I of $E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_2^{r_k}$ is isomorphic to \mathcal{K} and $(E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_2^{r_k})/I$ is isomorphic to a Cuntz-Krieger algebra with vanishing K_1 -group. Consequently, $E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_2^{r_k}$ and $E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_2^{r_k} \otimes \mathcal{O}_\infty$ are isomorphic to graph algebras.*

Proof: Note that for each $m \in \mathbb{N}$, by [EKTW, Theorem 7.2], E_m^{+1} , E_m^{-1} , $E_m^{+1} \otimes \mathcal{O}_\infty$, and $E_m^{-1} \otimes \mathcal{O}_\infty$ are graph algebras with $E_m^{\pm 1}/\mathcal{K}$ and $(E_m^{\pm 1} \otimes \mathcal{O}_\infty)/(\mathcal{K} \otimes \mathcal{O}_\infty) \cong (E_m^{\pm 1}/\mathcal{K}) \otimes \mathcal{O}_\infty$ are isomorphic to purely infinite Cuntz-Krieger algebras. Therefore, we may assume that $n \geq 1$. For notational convenience, set $A = E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_2^{r_k}$. Note that $I = \bigotimes_{k=1}^{n+1} \mathcal{K}$. Let $J = \mathcal{K} \otimes \bigotimes_{k=1}^n E_2^{r_k}$. Then J is a primitive ideal and $A/J \cong \mathcal{O}_m \otimes \bigotimes_{k=1}^n E_2^{r_k}$. We will show that J/I is stably isomorphic to an \mathcal{O}_2 -absorbing Cuntz-Krieger

algebra, A/J is isomorphic to a Cuntz-Krieger algebra with vanishing boundary maps, and the boundary maps in K -theory induced by the extension $0 \rightarrow J/I \rightarrow A/I \rightarrow A/J \rightarrow 0$ are zero.

We will first prove that J/I is \mathcal{O}_2 -absorbing. Note that it is enough to show that $(\bigotimes_{k=1}^n E_2^{r_k}) / (\bigotimes_{k=1}^n \mathcal{K})$ is \mathcal{O}_2 -absorbing since $J/I \cong \mathcal{K} \otimes (\bigotimes_{k=1}^n E_2^{r_k}) / (\bigotimes_{k=1}^n \mathcal{K})$. Since $E_2^{\pm} / \mathcal{K} \cong \mathcal{O}_2$ which is \mathcal{O}_2 -absorbing by [KP00, Theorem 3.8], we have that $(\bigotimes_{k=1}^n E_2^{r_k}) / (\bigotimes_{k=1}^n \mathcal{K})$ is \mathcal{O}_2 -absorbing for $n = 1$. Suppose $(\bigotimes_{k=1}^m E_2^{r_k}) / (\bigotimes_{k=1}^m \mathcal{K})$ is \mathcal{O}_2 -absorbing for $1 \leq m < n$. Consider the extension

$$\begin{aligned} 0 \rightarrow (E_2^{r_1} \otimes \bigotimes_{k=2}^n \mathcal{K}) / (\bigotimes_{k=1}^n \mathcal{K}) \rightarrow \\ \rightarrow (\bigotimes_{k=1}^n E_2^{r_k}) / (\bigotimes_{k=1}^n \mathcal{K}) \rightarrow (\bigotimes_{k=1}^n E_2^{r_k}) / (E_2^{r_1} \otimes \bigotimes_{k=2}^n \mathcal{K}) \rightarrow 0 \end{aligned}$$

Now, $(E_2^{r_1} \otimes \bigotimes_{k=2}^n \mathcal{K}) / (\bigotimes_{k=1}^n \mathcal{K}) \cong (E_2^{r_1} / \mathcal{K}) \otimes \bigotimes_{k=2}^n \mathcal{K} \cong \mathcal{O}_2 \otimes \bigotimes_{k=2}^n \mathcal{K}$ which is \mathcal{O}_2 -absorbing by [KP00, Theorem 3.8]. Since $(\bigotimes_{k=1}^n E_2^{r_k}) / (E_2^{r_1} \otimes \bigotimes_{k=2}^n \mathcal{K}) \cong E_2^{r_1} \otimes ((\bigotimes_{k=2}^n E_2^{r_k}) / (\bigotimes_{k=2}^n \mathcal{K}))$ and because of the inductive hypothesis, we have that $(\bigotimes_{k=1}^n E_2^{r_k}) / (E_2^{r_1} \otimes \bigotimes_{k=2}^n \mathcal{K})$ is \mathcal{O}_2 -absorbing. Hence, by [KP00, Theorem 3.8] and [TW07, Corollary 4.3], $(\bigotimes_{k=1}^n E_2^{r_k}) / (\bigotimes_{k=1}^n \mathcal{K})$ is \mathcal{O}_2 -absorbing. This proves our claim.

Since J/I is \mathcal{O}_2 -absorbing and J/I has finitely many ideals, by [Kir00], J/I is stably isomorphic to a Cuntz-Krieger algebra with vanishing boundary maps. This is because for any finite T_0 -space X , there exists an \mathcal{O}_2 -absorbing Cuntz-Krieger algebra with primitive ideal space X . We also note that the boundary maps in K -theory induced by the extension $0 \rightarrow J/I \rightarrow A/I \rightarrow A/J \rightarrow 0$ are zero since $K_*(J/I) = \{0\}$.

We now show that A/J is isomorphic to a Cuntz-Krieger algebra with vanishing boundary maps. Recall that $A/J \cong \mathcal{O}_m \otimes \bigotimes_{k=1}^n E_2^{r_k}$. Hence, every simple sub-quotient of A/J is isomorphic to $\mathcal{O}_m \otimes (\mathcal{I}_2 / \mathcal{I}_1)$ where $\mathcal{I}_1, \mathcal{I}_2$ are ideals of $\bigotimes_{k=1}^n E_2^{r_k}$ with $\mathcal{I}_1 \subseteq \mathcal{I}_2$ and $\mathcal{I}_2 / \mathcal{I}_1$ simple. Note that if $\mathcal{I}_1, \mathcal{I}_2$ are ideals of $\bigotimes_{k=1}^n E_2^{r_k}$ with $\mathcal{I}_1 \subseteq \mathcal{I}_2$ and $\mathcal{I}_2 / \mathcal{I}_1$ simple, then $\mathcal{I}_2 / \mathcal{I}_1 \cong \bigotimes_{k=1}^n B_k$ where B_k is a simple sub-quotient of $E_2^{r_k}$. Hence, every simple sub-quotient of $\bigotimes_{k=1}^n E_2^{r_k}$ is either isomorphic to $\bigotimes_{k=1}^n \mathcal{K}$ or is \mathcal{O}_2 -absorbing. Hence, every simple sub-quotient of $\mathcal{O}_m \otimes \bigotimes_{k=1}^n E_2^{r_k}$ is either stably isomorphic to \mathcal{O}_m or \mathcal{O}_2 . So every simple sub-quotient of A/J is stably isomorphic to a Cuntz-Krieger algebra. Consider the extension $e : 0 \rightarrow \mathcal{O}_m \otimes \mathcal{I}_1 \rightarrow \mathcal{O}_m \otimes \mathcal{I}_2 \rightarrow \mathcal{O}_m \otimes (\mathcal{I}_2 / \mathcal{I}_1) \rightarrow 0$ with $\mathcal{I}_2 / \mathcal{I}_1$ simple. If $\mathcal{I}_1 = \{0\}$ and $\mathcal{I}_2 = \bigotimes_{k=1}^n \mathcal{K}$, then $\mathcal{O}_m \otimes \mathcal{I}_1 = \{0\}$ which implies that e has vanishing boundary maps. If $\mathcal{I}_2 / \mathcal{I}_1$ is \mathcal{O}_2 -absorbing, then $K_*(\mathcal{O}_m \otimes (\mathcal{I}_2 / \mathcal{I}_1)) = \{0\}$ which also implies that e has vanishing boundary maps. By [Ben14, Corollary 3.6], we have that $A/J \cong \mathcal{O}_m \otimes \bigotimes_{k=1}^n E_2^{r_k}$ has vanishing boundary maps. Therefore, by [Ben14, Corollary 8.2], A/J is isomorphic to a Cuntz-Krieger algebra with vanishing boundary maps. This finishes the proof of the above claim.

The above claim shows that all the assumptions in [Ben14, Proposition 3.7, Proposition 3.10, and Corollary 8.4] are satisfied. Thus, A/I is isomorphic to a purely infinite Cuntz-Krieger algebra.

We now show that $K_1((E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_2^{r_k})/I) \cong \{0\}$. Since $0 \rightarrow J/I \rightarrow A/I \rightarrow A/J \rightarrow 0$ has vanishing boundary maps and the fact that J/I is \mathcal{O}_2 -absorbing, we have that the surjective map $A/I \rightarrow A/J$ induces an injective map $K_1(A/I) \rightarrow K_1(A/J)$. Since every simple sub-quotient of A/J is stably isomorphic to \mathcal{O}_m or \mathcal{O}_2 and since A/J has finitely many ideals, one can show that $K_1(A/J) = \{0\}$. Therefore, $K_1(A/I) = \{0\}$.

Lemma 6.4 implies that $E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_{m_k}^{r_k}$ and $E_m^{\pm 1} \otimes \bigotimes_{k=1}^n E_{m_k}^{r_k} \otimes \mathcal{O}_\infty$ are isomorphic to graph algebras. \square

LEMMA 6.6 *Let $m_1, m_2, \dots, m_n \in \mathbb{N}$. Then*

- (1) $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$ is stably isomorphic to unital graph algebra if and only if whenever there exists an i such that $m_i \in \{1\} \sqcup \mathbb{Z}_{\geq 3}$, we have that $m_j = 2$ for all $j \neq i$.
- (2) $\bigotimes_{k=1}^n E_{m_k}^{\pm 1} \otimes \mathcal{O}_\infty$ is stably isomorphic to unital graph algebra if and only if whenever there exists an i such that $m_i \in \{1\} \sqcup \mathbb{Z}_{\geq 3}$, we have that $m_j = 2$ for all $j \neq i$.

Proof: We prove (1). (2) is proved in a similar way.

Suppose whenever there exists an i such that $m_i \in \{1\} \sqcup \mathbb{Z}_{\geq 3}$, we have that $m_j = 2$ for all $j \neq i$. By Lemma 6.5, $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$ is isomorphic to a graph algebra. So, also stably isomorphic to a unital graph algebra.

Suppose $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$ is stably isomorphic to graph algebra. Note that $E_m^{\pm 1} \otimes \mathcal{K} \cong E_m^{\pm 1} \otimes \mathcal{K}$ for any m . Therefore, it is enough to prove the case $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$. Note that $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$ has finitely many ideals. Since $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$ is stably isomorphic to a unital graph algebra $C^*(E)$, we have that $C^*(E)$ has finitely many ideals. Therefore, every sub-quotient of $C^*(E)$ is stably isomorphic to a unital graph algebra with finitely many ideals. Consequently, every sub-quotient of $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$ is stably isomorphic to a unital graph algebra with finitely many ideals.

Suppose there exists i and j such that $m_i, m_j \in \{1\} \sqcup \mathbb{Z}_{\geq 3}$. Let $I = \bigotimes_{k=1}^n I_k$ be the ideal of $\bigotimes_{k=1}^n E_{m_k}^{\pm 1}$ where $I_k = \mathcal{K}$ if $k \notin \{i, j\}$, $I_i = E_{m_i}^{\pm 1}$, and $I_j = E_{m_j}^{\pm 1}$. From the above observation we must have that every sub-quotient of I is stably isomorphic to a unital graph algebra with finitely many ideals. Note that I is stably isomorphic to $E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1}$ and $E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1}$ has a quotient isomorphic to $\mathcal{O}_{m_i} \otimes \mathcal{O}_{m_j}$. Therefore, $\mathcal{O}_{m_i} \otimes \mathcal{O}_{m_j}$ is stably isomorphic to a graph algebra.

Let $\mathcal{K} \otimes \mathcal{K}$ be the smallest non-zero ideal of $E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1}$. By the Künneth formula, $K_0(E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1}) \cong \mathbb{Z}$ and $K_1(E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1}) = \{0\}$, and hence the extension $0 \rightarrow \mathcal{K} \otimes \mathcal{K} \rightarrow E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1} \rightarrow (E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1})/(\mathcal{K} \otimes \mathcal{K}) \rightarrow 0$ induces a six-term exact sequence in K -theory of the form

$$\begin{array}{ccccccc}
 \mathbb{Z} & \longrightarrow & \mathbb{Z} & \longrightarrow & K_0((E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1})/(\mathcal{K} \otimes \mathcal{K})) & & \\
 & & \uparrow & & \downarrow & & \\
 K_1((E_{m_i}^{\pm 1} \otimes E_{m_j}^{\pm 1})/(\mathcal{K} \otimes \mathcal{K})) & \longleftarrow & 0 & \longleftarrow & 0 & &
 \end{array}$$

In particular, $K_0((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K}))$ and $K_1((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K}))$ are cyclic groups.

Since $(E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K})$ is stably isomorphic to a graph algebra with finitely many ideals, $(E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K})$ has real rank zero. Therefore, the quotient of $(E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K})$ by the ideal $(\mathcal{K} \otimes E_{m_j}^{+1} + E_{m_i}^{+1} \otimes \mathcal{K})/(\mathcal{K} \otimes \mathcal{K})$ induces the following six-term exact sequence

$$\begin{CD}
 K_0(\mathcal{O}_{m_i}) \oplus K_0(\mathcal{O}_{m_j}) @>>> K_0((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K})) @>>> K_0(\mathcal{O}_{m_i} \otimes \mathcal{O}_{m_j}) \\
 @. @. @VVV \\
 @. @. 0 \\
 @. @. @VVV \\
 K_1(\mathcal{O}_{m_i} \otimes \mathcal{O}_{m_j}) @<<< K_1((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K})) @<<< K_1(\mathcal{O}_{m_i}) \oplus K_1(\mathcal{O}_{m_j})
 \end{CD} \tag{8}$$

Using the Künneth formula, we get

$$K_0(\mathcal{O}_{m_i} \otimes \mathcal{O}_{m_j}) = K_1(\mathcal{O}_{m_i} \otimes \mathcal{O}_{m_j}) = \begin{cases} \mathbb{Z}_{\gcd(m_i-1, m_j-1)} & \text{if } m_i, m_j \geq 3 \\ K_1(\mathcal{O}_{m_i}) \oplus K_0(\mathcal{O}_{m_i}) & \text{if } m_j = 1 \\ K_1(\mathcal{O}_{m_j}) \oplus K_0(\mathcal{O}_{m_j}) & \text{if } m_i = 1. \end{cases}$$

Since $\mathcal{O}_{m_i} \otimes \mathcal{O}_{m_j}$ is stably isomorphic to a unital graph algebra, $\gcd(m_i - 1, m_j - 1) = 1$ if $m_i, m_j \geq 3$ and $m_i = 1$ if and only if $m_j = 1$.

Suppose $m_i, m_j \geq 3$. The exactness of Diagram (8) implies that $K_0((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K})) \cong K_0(\mathcal{O}_{m_i}) \oplus K_0(\mathcal{O}_{m_j}) \cong \mathbb{Z}_{m_i-1} \oplus \mathbb{Z}_{m_j-1}$ which contradicts the fact that $K_0((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K}))$ is a cyclic group.

Suppose $m_i = 1$. Then $m_j = 1$. Then by the exactness of Diagram (8), $K_1((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K}))$ has a sub-group isomorphic to $K_1(\mathcal{O}_{m_i}) \oplus K_1(\mathcal{O}_{m_j}) \cong \mathbb{Z} \oplus \mathbb{Z}$. This can not happen since $K_1((E_{m_i}^{+1} \otimes E_{m_j}^{+1})/(\mathcal{K} \otimes \mathcal{K}))$ is a cyclic group. \square

Let the notation be as in Definition 4.1.

THEOREM 6.7 *Let Γ be a countable graph. Then $C^*(A_\Gamma^\pm)$ is isomorphic to a graph algebra if and only if one of the following holds*

1. $t(\Gamma) = 1, o(\Gamma) = 0$ and $N_k(\Gamma) = 0$ for all k
2. $t(\Gamma) = 0, N_{-1}(\Gamma) + N_1(\Gamma) < \infty$ and

$$\sum_{|k| \neq 1} N_k(\Gamma) \leq 1$$

Proof: Suppose there exists an isomorphism $\psi: C^*(A_\Gamma^\pm) \rightarrow C^*(E)$ for some countable directed graph E . Since $C^*(A_\Gamma^\pm)$ is unital, $C^*(E)$ is unital. Let $\Gamma_i = (V_i, E_i)$ be the co-irreducible components of Γ . To prove (1), let I be the ideal of $C^*(A_\Gamma^\pm)$ generated by $\{\otimes_j J_{ij}\}_i$ where $J_{ij} = C^*(A_{\Gamma_j}^\pm)$ if $j \neq i$ and

$$J_{ii} = \begin{cases} \mathcal{K} & \text{if } 1 < |V_i| < \infty \\ 0 & \text{otherwise.} \end{cases}$$

Then $C^*(A_\Gamma^+)/I \cong \bigotimes_i C^*(A_{\Gamma_i}^+)/J_{ii}$ where $C^*(A_{\Gamma_i}^+)/J_{ii}$ is a Kirchberg algebra if $|V_i| \geq 2$ and $C^*(A_{\Gamma_i}^+)/J_{ii} \cong \mathcal{T}$ otherwise. In particular,

$$\text{Prim}(C^*(A_\Gamma^+)/I) \cong \begin{cases} \prod_{k=1}^{t(\Gamma)} \text{Prim}(\mathcal{T}) & \text{if there exists } i \text{ with } |V_i| = 1 \\ \{\bullet\} & \text{otherwise.} \end{cases}$$

Note that I is generated by projections. Therefore, $\psi(I)$ is generated by projections and hence is a gauge-invariant ideal of $C^*(E)$. Hence, by [BPRS00, Corollary 3.5 and Theorem 3.6], $C^*(E)/\psi(I)$ is isomorphic to a graph algebra. Since $C^*(A_\Gamma^+)/I \cong C^*(E)/\psi(I)$, we have that $C^*(A_\Gamma^+)/I$ is isomorphic to a unital graph algebra. Note that $C^*(A_\Gamma^+)/I$ is \mathcal{O}_∞ -absorbing (if there exists i such that $|V_i| \geq 2$) or $C^*(A_\Gamma^+)/I \cong \bigotimes_{k=1}^{t(\Gamma)} \mathcal{T}$.

Suppose $C^*(A_\Gamma^+)/I$ is \mathcal{O}_∞ -absorbing. Since any unital \mathcal{O}_∞ -absorbing graph algebra has a finite primitive ideal space, we must have that $t(\Gamma) = 0$. Suppose $C^*(A_\Gamma^+)/I$ is not \mathcal{O}_∞ -absorbing. Then $C^*(A_\Gamma^+)/I \cong \bigotimes_{k=1}^{t(\Gamma)} \mathcal{T}$. Let J be the ideal generated by $\{\bigotimes_j J_{ij}\}_i$ where $J_{ij} = \mathcal{T}$ if $j \neq i$ and $J_{ii} = \mathcal{K}$, then J is an ideal generated by projections such that $(\bigotimes_{k=1}^{t(\Gamma)} \mathcal{T})/J \cong C(\mathbb{T}^{t(\Gamma)})$. Since $C^*(A_\Gamma^+)/I$ is isomorphic to a graph algebra and every ideal generated by projections in a graph algebra is gauge invariant, by [BPRS00, Corollary 3.5 and Theorem 3.6] every quotient of $C^*(A_\Gamma^+)/I$ by an ideal generated by projections is isomorphic to a graph algebra. Hence, $C(\mathbb{T}^{t(\Gamma)}) \cong (\bigotimes_{k=1}^{t(\Gamma)} \mathcal{T})/J$ is isomorphic to a unital graph algebra. Since the only unital commutative graph algebra is isomorphic to finite direct sums of \mathbb{C} and \mathbb{T} , we must have that $t(\Gamma) = 1$.

In both cases, we have shown that $t(\Gamma) \leq 1$. Suppose $o(\Gamma) \neq 0$ or $N_k(\Gamma) \neq 0$ for some k , then there exists an i such that $C^*(A_{\Gamma_i}^+)/J_{ii}$ is a Kirchberg algebra. Hence, by [KP00, Theorem 3.15] and [TW07, Corollary 3.4] $C^*(A_\Gamma^+)/I \cong \bigotimes_i C^*(A_{\Gamma_i}^+)/J_{ii}$ is an \mathcal{O}_∞ -absorbing C^* -algebra. Since every unital graph algebra that is \mathcal{O}_∞ -absorbing must have finitely many ideals and since

$$\text{Prim}(C^*(A_\Gamma^+)/I) \cong \begin{cases} \prod_{k=1}^{t(\Gamma)} \text{Prim}(\mathcal{T}) & \text{if there exists } i \text{ with } |V_i| = 1 \\ \{\bullet\} & \text{otherwise,} \end{cases}$$

we have that $t(\Gamma) = 0$. Hence, we only get a graph algebra in the case $t(\Gamma) = 1$ when all other data vanish.

Suppose $t(\Gamma) = 0$. Note that $1 < |V_i|$ for all i . Thus, $C^*(A_{\Gamma_i}^+)$ is a unital properly infinite C^* -algebra, and $\text{Prim}(C^*(A_{\Gamma_i}^+)) = \{x_i, y_i\}$ with open sets $\{\emptyset, \{x_i\}, \{x_i, y_i\}\}$ when $|V_i| < \infty$ and $\text{Prim}(C^*(A_{\Gamma_i}^+)) \cong \{\bullet\}$ when $|V_i| = \infty$.

We claim that $|\{k: N_k(\Gamma) \neq 0\}| < \infty$ and $N_k(\Gamma) < \infty$ for all k . Suppose first $|\{k: N_k(\Gamma) \neq 0\}| = \infty$ or $N_k(\Gamma) = \infty$ for some k . Then $C^*(A_\Gamma^+) \cong \bigotimes_{i=1}^\infty C^*(A_{\Gamma_i}^+)$ and $C^*(A_\Gamma^+)$ has infinitely many ideals. By Lemma 4.7, $C^*(A_\Gamma^+)$ is \mathcal{O}_∞ -absorbing. Again, using the fact that a unital graph algebra that is

\mathcal{O}_∞ -absorbing has finitely many ideals, we have a contradiction. Therefore, $|\{k: N_k(\Gamma) \neq 0\}| < \infty$ and $N_k(\Gamma) < \infty$ for all k , proving the claims in (2). Note that

$$C^*(A_\Gamma^+) \cong (E_1^0)^{\otimes N_0(\Gamma)} \otimes \bigotimes_{n=1}^{\infty} (E_{1+n}^{-1})^{\otimes N_{-n}(\Gamma)} \otimes \bigotimes_{n=1}^{\infty} (E_{1+n}^{+1})^{\otimes N_n(\Gamma)} \otimes (\mathcal{O}_\infty)^{\otimes o(\Gamma)}$$

By Lemma 6.6, (1) and (2) hold.

In the other direction, we have in the case (1) that $C^*(A_\Gamma^+) \cong \mathcal{T}$ which is isomorphic to a graph algebra. And in case (2) we have that either

$$C^*(A_\Gamma^+) \cong (E_2^{-1})^{\otimes N_{-1}(\Gamma)} \otimes (E_2^{+1})^{\otimes N_1(\Gamma)} \otimes (\mathcal{O}_\infty)^{\otimes o(\Gamma)},$$

$$C^*(A_\Gamma^+) \cong E_m^{+1} \otimes (E_2^{-1})^{\otimes N_{-1}(\Gamma)} \otimes (E_2^{+1})^{\otimes N_1(\Gamma)} \otimes (\mathcal{O}_\infty)^{\otimes o(\Gamma)}$$

for some $m \neq 2$, or

$$C^*(A_\Gamma^+) \cong E_m^{-1} \otimes (E_2^{-1})^{\otimes N_{-1}(\Gamma)} \otimes (E_2^{+1})^{\otimes N_1(\Gamma)} \otimes (\mathcal{O}_\infty)^{\otimes o(\Gamma)}.$$

for some $m \neq 2$. If $o(\Gamma) \geq 1$, then by [KP00, Theorem 3.15], $(\mathcal{O}_\infty)^{\otimes o(\Gamma)} \cong \mathcal{O}_\infty$. Hence, by Lemma 6.5, $C^*(A_\Gamma^+)$ is isomorphic to a graph algebra. \square

REMARK 6.8 *The relation between a (undirected, loop-free) graph Γ and a directed graph G_Γ with $C^*(A_\Gamma^+) \cong C^*(G_\Gamma)$ is somewhat opaque, although the proof given above in principle is constructive. In Figure 2 we present eight graphs presenting the C^* -algebras given by five-vertex graphs of Figure 1 in the unshaded regions.*

We conclude by establishing semiprojectivity and non-semiprojectivity of $C^*(A_\Gamma^+)$ in a number of cases, covering for instance all graphs with 5 or fewer vertices. We note, however, that this theorem does not contain a full answer to the question of which of the C^* -algebras under study are semiprojective. The most basic open case has $N_{-2} = 2$ and may be represented by a graph with 6 vertices.

THEOREM 6.9

1. When $t(\Gamma) > 1$, $C^*(A_\Gamma^+)$ is not semiprojective.
2. When $t(\Gamma) = 1$, $C^*(A_\Gamma^+)$ is semiprojective if and only if

$$o(\Gamma) = \sum_k N_k(\Gamma) = 0.$$

3. When $t(\Gamma) = 0$, $C^*(A_\Gamma^+)$ is semiprojective when $N_{-1}(\Gamma) + N_1(\Gamma) < \infty$ and

$$\sum_{|k| \neq 1} N_k(\Gamma) \leq 1$$




Figure 2: Graphs representing cases from Figure 1

Proof: We first note that by [End13, Corollary 4.4.16], a C^* -algebra of the form $A \otimes \mathcal{T}$ with A unital, nuclear, infinite-dimensional and in the UCT-class can never be semiprojective. This proves (1) and (2) since \mathcal{T} itself is trivially semiprojective.

For (3), we first apply Theorem 6.7 to see that $C^*(A_\Gamma^+)$ in this case is a unital graph algebra. We have seen that when $o(\Gamma) > 0$, $C^*(A_\Gamma^+)$ is strongly purely infinite, and when $o(\Gamma) = 0$, there is a minimal ideal \mathcal{K} in $C^*(A_\Gamma^+)$ so that $C^*(A_\Gamma^+)/\mathcal{K}$ is strongly purely infinite. In either case, [EK] applies to guarantee that the C^* -algebra is semiprojective. \square

REFERENCES

- [AD] Y. Antolin and D. Dreesen, *The Haagerup property is stable under graph products*, arXiv:1305.6748v2.
- [ABK14] S.E. Arklint, R. Bentmann, T. Katsura, *The K -theoretical range of Cuntz-Krieger algebras*, J. Funct. Anal. 266 (2014), no. 8, 5448–5466.
- [AGR] S.E. Arklint, J. Gabe, and E. Ruiz, *Hereditary C^* -subalgebras of graph C^* -algebras*, arXiv:1604.03085.
- [AR15] S.E. Arklint and E. Ruiz, *Corners of Cuntz-Krieger algebras*, Trans. Amer. Math. Soc. 367 (2015), 7595–7612.
- [Arv77] W. Arveson, *Notes on extensions C^* -algebras*, Duke Math. J. 44 (1977), 329–355.
- [BPRS00] T. Bates, D. Pask, I. Raeburn, and W. Szymanski, *C^* -algebras of row-finite graphs*, New York J. Math. 6 (2000), 307–324.
- [Ben14] R. Bentmann, *Kirchberg X -algebras with real rank zero and intermediate cancellation*, J. Noncommut. Geom. 8 (2014), 1061–1081.
- [BK] R. Bentmann and M. Köhler, *Universal coefficient theorems for C^* -algebras over finite topological spaces*, arXiv:1101.5702v3.
- [Bla77] B. Blackadar, *Infinite tensor products of C^* -algebras*, Pacific J. Math. 72 (1977), 313–334.
- [Bla85] ———, *Shape theory for C^* -algebras*, Math. Scand. 56 (1985), 249–275.
- [Bla04] ———, *Semiprojectivity in simple C^* -algebras*, Adv. Stud. Pure Math. 38 (2004), 1–17.
- [BD96] L.G. Brown and M. Dădărlat, *Extensions of C^* -algebras and quasi-diagonality*, J. London Math. Soc. (2) 53 (1996), no. 3, 582–600.

- [CL02] J. Crisp and M. Laca, *On the Toeplitz algebras of right-angled and finite-type Artin groups*, J. Austral. Math. Soc. 72 (2002), 223–245.
- [CL07] ———, *Boundary quotients and ideals of Toeplitz algebras of Artin groups*, J. Funct. Anal. 242 (2007), 127–156.
- [CEL13] J. Cuntz, S. Echterhoff, and X. Li, *On the K -theory of crossed products by automorphic semigroup actions*, Q. J. Math. 64 (2013), 747–784.
- [EK] S. Eilers and T. Katsura, *Semiprojectivity and properly infinite projections in graph C^* -algebras*, arXiv:1512.07277.
- [EKTW] S. Eilers, T. Katsura, M. Tomforde, and J. West, *The ranges of K -theoretical invariants for nonsimple graph algebras*, Trans. Amer. Math. Soc., 368 (2016), 3811–3847.
- [ELP99] S. Eilers, T. A. Loring, and G. K. Pedersen, *Morphisms of extensions of C^* -algebras: Pushing forward the Busby invariant*, Adv. Math. 147 (1999), 74–109.
- [ERR] S. Eilers, G. Restorff, and E. Ruiz, *Strong classification of extensions of classifiable C^* -algebras*, arXiv:1301.7695.
- [ERR09] ———, *Classification of extensions of classifiable C^* -algebras*, Adv. Math. 222 (2009), 2153–2172.
- [ERR10] ———, *On graph C^* -algebras with a linear ideal lattice*, Bull. Malays. Math. Sci. Soc. 33 (2010), no. 2, 233–241.
- [ERR13] ———, *Classifying C^* -algebras with both finite and infinite subquotients*, J. Funct. Anal. 265 (2013), 449–468.
- [End13] D. Enders, *On the structure of certain classes of semiprojective C^* -algebras*, Ph.D. thesis, Westfälische Wilhelms-Universität Münster, 2013.
- [End15] D. Enders, *Semiprojectivity for Kirchberg algebras*, preprint, arXiv:1507.0609, 2015.
- [EL91] R. Exel and T.A. Loring, *Invariants of almost commuting unitaries*, J. Funct. Anal. 95 (1991), 364–376.
- [FL07] X. Fang and S. Liu, *Extension algebras of Cuntz algebras*, J. Math. Anal. Appl. 329 (2007), 655–663.
- [HL+14] D. Hay, M. Loving, M. Montgomery, E. Ruiz, and K. Todd, *Non-stable K -theory for Leavitt path algebras*, Rocky Mountain J. Math. 44 (2014), 1817–1850.

- [Iva10] N. Ivanov, *The K -theory of Toeplitz C^* -algebras of right-angled Artin groups*, Trans. Amer. Math. Soc. 362 (2010), no. 11, 6003–6027.
- [JT91] K.K. Jensen and K. Thomsen, *Elements of KK -theory*, Birkhäuser, Boston, 1991.
- [Kir] E. Kirchberg, *The classification of purely infinite C^* -algebras using Kasparov's theory* 3rd draft. Preprint.
- [Kir00] E. Kirchberg, *Das nicht-kommutative Michael-Auswahlprinzip und die Klassifikation nicht-einfacher Algebren*, C^* -algebras (Münster, 1999), Springer, Berlin, 2000, pp. 92–141.
- [KP00] E. Kirchberg and N. C. Phillips, *Embedding of exact C^* -algebras in the Cuntz algebra \mathcal{O}_2* , J. Reine Angew. Math. 525 (2000), 17–53.
- [KR02] E. Kirchberg and M. Rørdam, *Infinite non-simple C^* -algebras: absorbing the Cuntz algebras \mathcal{O}_∞* , Adv. Math. 167 (2002), 195–264.
- [Li12] X. Li, *Semigroup C^* -algebras and amenability of semigroups*, J. Funct. Anal. 262 (2012), 4302–4340.
- [Li13] X. Li, *Nuclearity of semigroup C^* -algebras and the connection to amenability*, Adv. Math. 244 (2013), 626–662.
- [Lor97] T.A. Loring, *Lifting solutions to perturbing problems in C^* -algebras*, Fields Institute Monographs, vol. 8, American Mathematical Society, Providence, RI, 1997.
- [NR97] G. Niblo and L. Reeves, *Groups acting on $CAT(0)$ cube complexes*, Geom. & Top. 1 (1997), 1–7.
- [Par02] L. Paris, *Artin monoids inject in their groups*, Comment. Math. Helv. 77 (2002), 609–637.
- [PR07] C. Pasnicu and M. Rørdam, *Purely infinite C^* -algebras of real rank zero*, J. Reine Angew. Math. 613 (2007), 51–73.
- [Res06] G. Restorff, *Classification of Cuntz-Krieger algebras up to stable isomorphism*, J. Reine Angew. Math. 598 (2006), 185–210.
- [RR07] G. Restorff and E. Ruiz, *On Rørdam's classification of certain C^* -algebras with one nontrivial ideal II*, Math. Scand. 101 (2007), 280–292.
- [Ror97] M. Rørdam, *Classification of extensions of certain C^* -algebras by their six term exact sequences in K -theory*, Math. Ann. 308 (1997), 93–117.

- [Sor13] A. Sørensen, *Geometric classification of simple graph algebras*, Ergodic Theory Dynam. Systems 33 (2013), no. 4, 1199–1220.
- [TW07] A. Toms and W. Winter, *Strongly self-absorbing C^* -algebras*, Trans. Amer. Math. Soc. 359 (2007), 3999–4029.
- [Voi83] D. Voiculescu, *Asymptotically commuting finite rank unitary operators without commuting approximants*, Acta Sci. Math. (Szeged) 45 (1983), no. 1-4, 429–431.

Søren Eilers
Department of
Mathematical Sciences
University of Copenhagen
Universitetsparken 5
DK-2100 Copenhagen
Denmark

Xin Li
School of
Mathematical Sciences
Queen Mary University
of London
Mile End Road
London E1 4NS
United Kingdom

Efren Ruiz
Department of Mathematics
University of Hawaii, Hilo
200 W. Kawili St., Hilo
Hawaii, 96720-4091
USA

