

Research Article

Solution to the Linear Fractional Differential Equation Using Adomian Decomposition Method

Jin-Fa Cheng¹ and Yu-Ming Chu²

¹ *Department of Mathematics, Xiamen University, Xiamen 361005, China*

² *Department of Mathematics, Huzhou Teachers College, Huzhou 313000, China*

Correspondence should be addressed to Yu-Ming Chu, chuyuming2005@yahoo.com.cn

Received 25 March 2010; Accepted 1 June 2010

Academic Editor: Geraldo Silva

Copyright © 2011 J.-F. Cheng and Y.-M. Chu. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

We obtain the analytical general solution of the linear fractional differential equations with constant coefficients by Adomian decomposition method under nonhomogeneous initial value condition, which is in the sense of the Caputo fractional derivative.

1. Introduction

Fractional differential equations are hot topics both in mathematics and physics. Recently, the fractional differential equations have been the subject of intensive research. There are several methods to obtain the solution, such as the Laplace transform method, power series method, and Green function method. Many remarkable results for the fractional differential equations can be found in the literature [1–11]. In particular, the Adomian decomposition method has attracted the attention of many mathematicians [12–15].

For a better understanding of the fractional derivatives and for a physical understanding of the fractional equations, the readers can refer to the recent publications in [16, 17]. Ebaid [18] suggested a modification of the Adomian method, and a few iterations lead to exact solution. Das [19] compared the variational iteration method with the Adomian method for fractional equations and found that the variational iteration method is much more effective. For other methods of the fractional differential equations, especially the homotopy perturbation method, variational iteration method and differential transform method were presented in [20, 21].

Consider the following n -term fractional differential equation with constant coefficients:

$$a_n \left[{}^C D^{\beta_n} \right] y(t) + a_{n-1} \left[{}^C D^{\beta_{n-1}} \right] y(t) + \cdots + a_1 \left[{}^C D^{\beta_1} \right] y(t) + a_0 \left[{}^C D^{\beta_0} \right] y(t) = f(t), \quad (1.1)$$

where $n + 1 > \beta \geq n > \beta_{n-1} > \cdots > \beta_1 > \beta_0$ and a_i ($i = 0, 1, \dots, n$) is a real constant. In [12], the authors obtain the particular solution of (1.1) of the homogeneous initial value problem of the form

$$y^i(0) = 0, \quad i = 0, 1, \dots, n. \quad (1.2)$$

However, it seems also more meaningful and more complicated for solving general solution of (1.1) under nonhomogeneous initial value condition. Therefore, in this paper, we will remove the restriction of the homogeneous initial value, consider the nonhomogeneous initial value problems of the form

$$y^{j_i}(0) = c_{ij_i}, \quad i = 0, 1, \dots, n, \quad j_i = 1, 2, \dots, l_i, \quad l_i - 1 \leq \beta_i < l_i, \quad (1.3)$$

and obtain the analytical general solution of (1.1), which generalizes the result in [12].

We organize the paper as follows. In Section 2, we give some basic definitions and properties. In Section 3, we obtain the analytical general solution of the linear fractional differential equations by Adomian decomposition method. Some explicit examples are given in Section 4.

2. Basic Definitions and Notations

Definition 2.1 (see [1]). The Riemann-Liouville integral of order p is defined by

$${}_a D_t^{-p} f(t) = \frac{1}{\Gamma(p)} \int_a^t (t - \tau)^{p-1} f(\tau) d\tau, \quad p > 0. \quad (2.1)$$

From Definition 2.1, we clearly see that

$${}_a D_t^{-\alpha} ((t - a)^\nu) = \frac{\Gamma(1 + \nu)}{\Gamma(1 + \nu + \alpha)} (t - a)^{\nu + \alpha}, \quad (2.2)$$

$${}_a D_t^\alpha ((t - a)^\nu) = \frac{\Gamma(1 + \nu)}{\Gamma(1 + \nu - \alpha)} (t - a)^{\nu - \alpha}, \quad (2.3)$$

where $\alpha > 0$ and ν is a real number.

Definition 2.2 (see [1]). For $f(t) \in C^m (m \in \mathbb{N})$, the Caputo fractional derivative of $f(t)$ is defined by

$${}^C_a D_t^p f(t) = \begin{cases} D^{m-p} f^{(m)}(t), & m-1 < p < m, \\ \frac{d^{(m)}}{dt^m} f(t), & p = m, \\ {}_a D_t^p f(t), & p \leq 0. \end{cases} \quad (2.4)$$

Therefore,

$${}^C_a D_t^{-\alpha} ((t-a)^\nu) = {}_a D_t^{-\alpha} ((t-a)^\nu) = \frac{\Gamma(1+\nu)}{\Gamma(1+\nu+\alpha)} (t-a)^{\nu+\alpha}. \quad (2.5)$$

Lemma 2.3 (see [1]). *If $f(t)$ is continuous, then*

$${}^C_a D_t^{-n} f^{(n)}(t) = {}_a D_t^{-n} f^{(n)}(t) = f(t) - \sum_{j=0}^{n-1} \frac{f^{(j)}(a)(t-a)^j}{\Gamma(j+1)}, \quad n \in \mathbb{Z}. \quad (2.6)$$

Lemma 2.4 (see [1]). *If $f(t)$ is continuous, then*

$${}^C_a D_t^{-p} ({}^C_a D_t^{-q}) f(t) = {}_a D_t^{-p} ({}_a D_t^{-q}) f(t) = {}_a D_t^{-p-q} f(t) = {}^C_a D_t^{-p-q} f(t) = {}^C_a D_t^{-q} ({}^C_a D_t^{-p}) f(t). \quad (2.7)$$

Lemma 2.5. *If $f(t)$ is continuous, then*

$${}^C_a D_t^{-p} ({}^C_a D_t^q) f(t) = {}_a D_t^{q-p} f(t) - \sum_{j=0}^{m-1} f^{(j)}(a) \frac{(t-a)^{j-q+p}}{\Gamma(j-q+p+1)}, \quad (2.8)$$

where $m-1 < q < m$ and $q \leq p$.

Proof. From Definition 2.2 and Lemmas 2.3–2.4, we get

$$\begin{aligned} {}^C_a D_t^{-p} ({}^C_a D_t^q) f(t) &= {}_a D_t^{-p} ({}_a D_t^{q-m}) f^{(m)}(t) = {}_a D_t^{-p+q} \left[{}_a D_t^{-q} ({}_a D_t^{q-m}) f^{(m)}(t) \right] \\ &= {}_a D_t^{-p+q} \left[{}_a D_t^{-m} f^{(m)}(t) \right] = {}_a D_t^{-p+q} \left[f(t) - \sum_{j=0}^{m-1} \frac{f^{(j)}(a)(t-a)^j}{\Gamma(j+1)} \right] \\ &= {}_a D_t^{q-p} f(t) - \sum_{j=0}^{m-1} \frac{f^{(j)}(a)(t-a)^{j-q+p}}{\Gamma(j-q+p+1)}. \end{aligned} \quad (2.9)$$

It is easy to see that

$${}^C D_t^{-p} ({}^C D_t^p) f(t) = f(t) - \sum_{j=0}^{m-1} f^{(j)}(a) \frac{(t-a)^j}{\Gamma(j+1)}. \quad (2.10)$$

□

Proposition 2.6 (see [12]). *One has*

$$\sum_{k_1=0}^{\infty} \cdots \sum_{k_n=0}^{\infty} a_{k_1, k_2, \dots, k_{n-1}, k_n} = \sum_{m=0}^{\infty} \sum_{\substack{k_1, \dots, k_{n-1}, k_n \geq 0 \\ k_1 + \dots + k_{n-1} + k_n = m}} a_{k_1, k_2, \dots, k_{n-1}, k_n}. \quad (2.11)$$

Proposition 2.7 (see [12]). *More over, one has*

$$\sum_{m=0}^{\infty} \sum_{\substack{k_1, \dots, k_{n-1}, k_n \geq 0 \\ k_1 + \dots + k_{n-1} + k_n = m}} a_{k_1, k_2, \dots, k_{n-1}, k_n} = \sum_{s=0}^{\infty} \sum_{\substack{k_1, \dots, k_{n-1} \geq 0 \\ k_1 + \dots + k_{n-1} = s}} \sum_{k_n=0}^{\infty} a_{k_1, k_2, \dots, k_{n-1}, k_n}. \quad (2.12)$$

3. The Analytical Solution of the Linear Constant Coefficient Fractional Differential Equation

For simplicity, if $a = 0$, then we denote ${}^C D_t^p$ or ${}^C D_t^{-p}$ by ${}^C D^p$ or ${}^C D^{-p}$.

In this section, we use Adomian decomposition method to discuss the general form of the linear fractional differential equations with constant coefficients, and apply and some basic transformation and integration to obtain the solution of the equations.

Let us consider the following n -term linear fractional differential equations with constant coefficients:

$$\begin{aligned} a_n [{}^C D^{\beta_n}] y(t) + a_{n-1} [{}^C D^{\beta_{n-1}}] y(t) + \cdots + a_1 [{}^C D^{\beta_1}] y(t) + a_0 [{}^C D^{\beta_0}] y(t) &= f(t), \\ y^{(j_i)}(0) &= c_{ij_i}, \quad i = 0, 1, \dots, n, \quad j_i = 1, 2, \dots, l_i, \quad l_i - 1 \leq \beta_i < l_i, \end{aligned} \quad (3.1)$$

where $n+1 > \beta_n \geq n > \beta_{n-1} \cdots > \beta_1 > \beta_0$, a_i and c_{ij_i} are real constants, ${}^C D^\alpha = {}^C D_t^\alpha$ denotes Caputo fractional derivative of order α .

Applying ${}^C D^{-\beta_n}$ to both sides of (1.1) and utilizing Lemma 2.5, we get

$$\begin{aligned} y(t) + \frac{a_{n-1}}{a_n} {}^C D^{\beta_{n-1}-\beta_n} y(t) + \cdots + \frac{a_0}{a_n} {}^C D^{\beta_0-\beta_n} y(t) \\ = \frac{1}{a_n} {}^C D^{-\beta_n} f(t) + \frac{1}{a_n} \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} y^{(j_i)}(0) \frac{t^{\beta_n-\beta_i+j_i}}{\Gamma(1+\beta_n-\beta_i+j_i)}. \end{aligned} \quad (3.2)$$

By the Adomian decomposition method, we obtain the recursive relationship as follows:

$$\begin{aligned}
 y_0(t) &= \frac{1}{a_n} {}^C D^{-\beta_n} f(t) + \frac{1}{a_n} \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} y^{(j_i)}(0) \frac{t^{\beta_n-\beta_i+j_i}}{\Gamma(1+\beta_n-\beta_i+j_i)}, \\
 y_1(t) &= -\left(\frac{a_{n-1}}{a_n} {}^C D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} {}^C D^{\beta_0-\beta_n}\right) y_0(t), \\
 y_2(t) &= (-1)^2 \left(\frac{a_{n-1}}{a_n} {}^C D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} {}^C D^{\beta_0-\beta_n}\right)^2 y_0(t), \\
 &\vdots \\
 y_s(t) &= (-1)^s \left(\frac{a_{n-1}}{a_n} {}^C D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} {}^C D^{\beta_0-\beta_n}\right)^s y_0(t), \\
 &\vdots
 \end{aligned} \tag{3.3}$$

By Adomian decomposition method, adding all terms of the recursion, we obtain the solution of (1.1) as

$$\begin{aligned}
 y(t) &= \sum_{s=0}^{\infty} y_s(t) \\
 &= \sum_{s=0}^{\infty} (-1)^s \left(\frac{a_{n-1}}{a_n} {}^C D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} {}^C D^{\beta_0-\beta_n}\right)^s y_0(t) \\
 &= \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \left(\frac{a_{n-1}}{a_n} D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} D^{\beta_0-\beta_n}\right)^s D^{-\beta_n} f(t) \\
 &\quad + \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \left(\frac{a_{n-1}}{a_n} D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} D^{\beta_0-\beta_n}\right)^s \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} y^{(j_i)}(0) \frac{t^{\beta_n-\beta_i+j_i}}{\Gamma(1+\beta_n-\beta_i+j_i)}.
 \end{aligned} \tag{3.4}$$

Let

$$\begin{aligned}
 I_1 &= \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \left(\frac{a_{n-1}}{a_n} D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} D^{\beta_0-\beta_n}\right)^s D^{-\beta_n} f(t) \\
 I_2 &= \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \left(\frac{a_{n-1}}{a_n} D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} D^{\beta_0-\beta_n}\right)^s \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} y^{(j_i)}(0) \frac{t^{\beta_n-\beta_i+j_i}}{\Gamma(1+\beta_n-\beta_i+j_i)}.
 \end{aligned} \tag{3.5}$$

Then,

$$y(t) = I_1 + I_2. \tag{3.6}$$

Next, we estimate I_1 and I_2 , respectively.

For I_1 , by [12] we obtain

$$\begin{aligned}
& \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \left(\frac{a_{n-1}}{a_n} D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} D^{\beta_0-\beta_n} \right)^s D^{-\beta_n} f(t), \\
&= \int_0^t \frac{1}{a_n} \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0 + k_1 + \dots + k_{n-2} = m}} (m; k_0, k_1, \dots, k_{n-2}) \\
&\quad \times \prod_{p=0}^{n-2} \left(\frac{a_p}{a_n} \right)^{k_p} (t - \tau)^{(\beta_n - \beta_{n-1})m + \beta_n + \sum_{j=0}^{n-2} (\beta_{n-1} - \beta_j)k_{j-1}} \\
&\quad \times E_{\beta_n - \beta_{n-1}, \beta_n + \sum_{j=0}^{n-2} (\beta_{n-1} - \beta_j)k_j}^{(m)} \left(-\frac{a_{n-1}}{a_n} D^{\beta_n - \beta_{n-1}} \right) f(\tau) d\tau.
\end{aligned} \tag{3.7}$$

For I_2 , by the initial conditions (2.10) we get

$$\begin{aligned}
& \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \left(\frac{a_{n-1}}{a_n} D^{\beta_{n-1}-\beta_n} + \dots + \frac{a_0}{a_n} D^{\beta_0-\beta_n} \right)^s \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} y^{(j_i)}(0) \frac{t^{\beta_n - \beta_i + j_i}}{\Gamma(1 + \beta_n - \beta_i + j_i)} \\
&= \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \sum_{\substack{k_0, k_1, \dots, k_{n-1} \geq 0 \\ k_0 + k_1 + \dots + k_{n-1} = s}} \frac{s!}{k_0! k_1! \dots k_{n-1}!} \left(\frac{a_{n-1}}{a_n} \right)^{k_{n-1}} \left(\frac{a_{n-2}}{a_n} \right)^{k_{n-2}} \dots \left(\frac{a_0}{a_n} \right)^{k_0} \\
&\quad \times D^{k_{n-1}(\beta_{n-1} - \beta_n) + k_{n-2}(\beta_{n-2} - \beta_n) + \dots + k_0(\beta_0 - \beta_n)} \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} \frac{t^{\beta_n - \beta_i + j_i}}{\Gamma(1 + \beta_n - \beta_i + j_i)}.
\end{aligned} \tag{3.8}$$

Using formulas (2.2) and (2.3), the above expression can be written as

$$\begin{aligned}
& \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \sum_{\substack{k_0, k_1, \dots, k_{n-1} \geq 0 \\ k_0 + k_1 + \dots + k_{n-1} = s}} \frac{s!}{k_0! k_1! \dots k_{n-1}!} \left(\frac{a_{n-1}}{a_n} \right)^{k_{n-1}} \left(\frac{a_{n-2}}{a_n} \right)^{k_{n-2}} \dots \left(\frac{a_0}{a_n} \right)^{k_0} \\
&\quad \times \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} \frac{\Gamma(\mathcal{A}) t^{\beta_n - \beta_i + j_i + k_{n-1}(\beta_n - \beta_{n-1}) + k_{n-2}(\beta_n - \beta_{n-2}) + \dots + k_0(\beta_n - \beta_0)}}{\Gamma(\mathcal{A}) \cdot \Gamma(\mathcal{A} + k_{n-1}(\beta_n - \beta_{n-1}) + k_{n-2}(\beta_n - \beta_{n-2}) + \dots + k_0(\beta_n - \beta_0))} \\
&= \frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0 + k_1 + \dots + k_{n-1} = s}} \frac{s!}{k_0! k_1! \dots k_{n-1}!} \left(\frac{a_{n-1}}{a_n} \right)^{k_{n-1}} \left(\frac{a_{n-2}}{a_n} \right)^{k_{n-2}} \dots \left(\frac{a_0}{a_n} \right)^{k_0} \\
&\quad \times \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} \frac{t^{\beta_n - \beta_i + j_i + k_{n-1}(\beta_n - \beta_{n-1}) + k_{n-2}(\beta_n - \beta_{n-2}) + \dots + k_0(\beta_n - \beta_0)}}{\Gamma(\mathcal{A} + k_{n-1}(\beta_n - \beta_{n-1}) + k_{n-2}(\beta_n - \beta_{n-2}) + \dots + k_0(\beta_n - \beta_0))}
\end{aligned}$$

$$\begin{aligned}
 &= \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} D^{(\beta_i-j_i-1)} \left[\frac{1}{a_n} \sum_{s=0}^{\infty} (-1)^s \sum_{\substack{k_0, k_1, \dots, k_{n-1} \geq 0 \\ k_0+k_1+\dots+k_{n-1}=s}} \frac{s!}{k_0!k_1! \dots k_{n-1}!} \right. \\
 &\quad \times \left(\frac{a_{n-1}}{a_n} \right)^{k_{n-1}} \left(\frac{a_{n-2}}{a_n} \right)^{k_{n-2}} \dots \left(\frac{a_0}{a_n} \right)^{k_0} \\
 &\quad \times \left. \frac{t^{\beta_{n-1}+k_{n-1}(\beta_n-\beta_{n-1})+k_{n-2}(\beta_n-\beta_{n-2})+\dots+k_0(\beta_n-\beta_0)}}{\Gamma(\beta_n+k_{n-1}(\beta_n-\beta_{n-1})+k_{n-2}(\beta_n-\beta_{n-2})+\dots+k_0(\beta_n-\beta_0))} \right], \tag{3.9}
 \end{aligned}$$

where \mathcal{A} denotes $1 + \beta_n - \beta_i + j_i$.

Using Propositions 2.6 and 2.7, the above solution is equivalent to the following form:

$$\begin{aligned}
 &\sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} D^{(\beta_i-j_i-1)} \left[\frac{1}{a_n} \sum_{k_{n-1}=0}^{\infty} \sum_{m=0}^{\infty} \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0+k_1+\dots+k_{n-2}=m}} (-1)^{k_{n-1}+m} \frac{(k_{n-1}+m)!}{k_0!k_1! \dots k_{n-1}!} \right. \\
 &\quad \times \left(\frac{a_{n-1}}{a_n} \right)^{k_{n-1}} \left(\frac{a_{n-2}}{a_n} \right)^{k_{n-2}} \dots \left(\frac{a_0}{a_n} \right)^{k_0} \\
 &\quad \times \left. \frac{t^{\beta_{n-1}+k_{n-1}(\beta_n-\beta_{n-1})+k_{n-2}(\beta_n-\beta_{n-2})+\dots+k_0(\beta_n-\beta_0)}}{\Gamma(\beta_n+k_{n-1}(\beta_n-\beta_{n-1})+k_{n-2}(\beta_n-\beta_{n-2})+\dots+k_0(\beta_n-\beta_0))} \right] \\
 &= \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} D^{(\beta_i-j_i-1)} \left[\frac{1}{a_n} \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0+k_1+\dots+k_{n-2}=m}} \frac{m!}{k_0!k_1! \dots k_{n-2}!} \right. \\
 &\quad \times \prod_{r=0}^{n-2} \left(\frac{a_r}{a_n} \right)^{k_r} t^{(\beta_n-\beta_{n-1})m + \sum_{\xi=0}^{n-2} (\beta_{n-1}-\beta_{\xi})k_{\xi} + \beta_n - 1} \\
 &\quad \times \sum_{k_{n-1}=0}^{\infty} (-1)^{k_{n-1}} \left(\frac{a_{n-1}}{a_n} \right)^{k_{n-1}} \frac{(k_{n-1}+m)!}{k_{n-1}!} \\
 &\quad \times \left. \frac{t^{k_{n-1}(\beta_n-\beta_{n-1})}}{\Gamma(k_{n-1}(\beta_n-\beta_{n-1}) + (\beta_n-\beta_{n-1})m + \sum_{\delta=0}^{n-2} (\beta_{n-1}-\beta_{\delta})k_{\delta} + \beta_n)} \right]
 \end{aligned}$$

$$\begin{aligned}
&= \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} D^{(\beta_i-j_i-1)} \left[\frac{1}{a_n} \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0 + k_1 + \dots + k_{n-2} = m}} (m; k_0, k_1, \dots, k_{n-2}) \right. \\
&\quad \times \prod_{r=0}^{n-2} \left(\frac{a_r}{a_n} \right)^{k_r} t^{(\beta_n - \beta_{n-1})m + \sum_{\xi=0}^{n-2} (\beta_{n-1} - \beta_{\xi}) k_{\xi} + \beta_{n-1}} \\
&\quad \left. \times E_{\beta_n - \beta_{n-1}, \sum_{\delta=0}^{n-2} (\beta_{n-1} - \beta_{\delta}) k_{\delta} + \beta_n}^{(m)} \left(-\frac{a_{n-1}}{a_n} t^{\beta_n - \beta_{n-1}} \right) \right]. \tag{3.10}
\end{aligned}$$

Therefore,

$$\begin{aligned}
y(t) &= I_1 + I_2 \\
&= \int_0^t \frac{1}{a_n} \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0 + k_1 + \dots + k_{n-2} = m}} (m; k_0, k_1, \dots, k_{n-2}) \\
&\quad \times \prod_{p=0}^{n-2} \left(\frac{a_p}{a_n} \right)^{k_p} (t - \tau)^{(\beta_n - \beta_{n-1})m + \beta_n + \sum_{j=0}^{n-2} (\beta_{n-1} - \beta_j) k_j - 1} \\
&\quad \times E_{\beta_n - \beta_{n-1}, \beta_n + \sum_{j=0}^{n-2} (\beta_{n-1} - \beta_j) k_j}^{(m)} \left(-\frac{a_{n-1}}{a_n} D^{\beta_n - \beta_{n-1}} \right) f(\tau) d\tau \\
&\quad + \sum_{i=0}^n a_i \sum_{j_i=0}^{l_i-1} c_{ij_i} D^{(\beta_i-j_i-1)} \left[\frac{1}{a_n} \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0 + k_1 + \dots + k_{n-2} = m}} (m; k_0, k_1, \dots, k_{n-2}) \right. \\
&\quad \times \prod_{r=0}^{n-2} \left(\frac{a_r}{a_n} \right)^{k_r} t^{(\beta_n - \beta_{n-1})m + \sum_{\xi=0}^{n-2} (\beta_{n-1} - \beta_{\xi}) k_{\xi} + \beta_{n-1}} \\
&\quad \left. \times E_{\beta_n - \beta_{n-1}, \sum_{\delta=0}^{n-2} (\beta_{n-1} - \beta_{\delta}) k_{\delta} + \beta_n}^{(m)} \left(-\frac{a_{n-1}}{a_n} t^{\beta_n - \beta_{n-1}} \right) \right], \tag{3.11}
\end{aligned}$$

where

$$(m; k_0, k_1, \dots, k_{n-2}) = \frac{m!}{k_0! k_1! \dots k_{n-2}!}, \tag{3.12}$$

and $E_{\lambda, \mu}^{(i)}(y)$ is the Mittag-Leffler function

$$E_{\lambda, \mu}^{(i)}(y) = \frac{d^i}{dy^i} E_{\lambda, \mu}(y) = \sum_{j=0}^{\infty} \frac{(i+j)! y^j}{j! \Gamma(\lambda j + \lambda i + \mu)}. \tag{3.13}$$

Substituting the Green function

$$\begin{aligned}
 G_n(t) &= \frac{1}{a_n} \sum_{m=0}^{\infty} \frac{(-1)^m}{m!} \sum_{\substack{k_0, k_1, \dots, k_{n-2} \geq 0 \\ k_0 + k_1 + \dots + k_{n-2} = m}} (m; k_0, k_1, \dots, k_{n-2}) \\
 &\times \prod_{p=0}^{n-2} \left(\frac{a_p}{a_n} \right)^{k_p} t^{(\beta_n - \beta_{n-1})m + \beta_n + \sum_{j=0}^{n-2} (\beta_{n-1} - \beta_j)k_j - 1} \\
 &\times E_{\beta_n - \beta_{n-1}, \beta_n + \sum_{j=0}^{n-2} (\beta_{n-1} - \beta_j)k_j}^{(m)} \left(-\frac{a_{n-1}}{a_n} D^{\beta_n - \beta_{n-1}} \right)
 \end{aligned} \tag{3.14}$$

into the above expression, we know that

$$y(t) = \int_0^t G_n(t - \tau) f(\tau) d\tau + \sum_{i=0}^n a_i \sum_{j=0}^{i-1} c_{ij} G_n^{(\beta_i - j - 1)}(t) \tag{3.15}$$

is the analytical general solution of (1.1).

4. Illustrative Examples

In order to verify our conclusions, we give some examples.

(1). Consider an initial value problem for the relaxation-oscillation equation (see [1])

$$\begin{aligned}
 {}^C D^\alpha y(t) + Ay(t) &= f(t), \quad t > 0, \\
 y^{(j)}(0) &= b_j, \quad j = 1, 2, \dots, m-1, \quad m-1 < \alpha < m,
 \end{aligned} \tag{4.1}$$

where b_j are real constants.

Utilizing Lemma 2.5 and applying ${}^C D^{-\alpha}$ to both sides of (4.1), we obtain

$$y(t) + AD^{-\alpha} y(t) = D^{-\alpha} f(t) + \sum_{j=0}^{m-1} y^{(j)}(0) \frac{t^j}{\Gamma(j+1)}. \tag{4.2}$$

According to the above procedure of solving the linear fractional differential equations with constant coefficients and using the Adomian decomposition method, let

$$\begin{aligned}
 y_0(t) &= D^{-\alpha} f(t) + \sum_{j=0}^{m-1} y^{(j)}(0) \frac{t^j}{\Gamma(j+1)}, \\
 y_1(t) &= -AD^{-\alpha} y_0(t), \\
 y_2(t) &= -AD^{-\alpha} y_1(t) = (-A)^2 D^{-2\alpha} y_0(t), \\
 &\vdots \\
 y_s(t) &= -AD^{-\alpha} y_{s-1}(t) = (-A)^s D^{-s\alpha} y_0(t), \\
 &\vdots
 \end{aligned} \tag{4.3}$$

Adding all of the above terms, we obtain the solution of the equation by Adomian decomposition method as follows:

$$\begin{aligned}
 y(t) &= \sum_{s=0}^{\infty} y_s(t) \\
 &= \sum_{s=0}^{\infty} (-A)^s D^{-s\alpha} y_0(t) \\
 &= \sum_{s=0}^{\infty} (-A)^s D^{-(s-1)\alpha} f(t) + \sum_{s=0}^{\infty} (-A)^s D^{-s\alpha} \sum_{j=0}^{m-1} y^{(j)}(0) \frac{t^j}{\Gamma(j+1)} \\
 &= \sum_{s=0}^{\infty} (-A)^s \frac{1}{\Gamma((s+1)\alpha)} \int_0^t (t-\tau)^{(s+1)\alpha-1} f(\tau) d\tau + \sum_{j=0}^{m-1} b_j \sum_{s=0}^{\infty} (-A)^s \frac{t^{j+s\alpha}}{\Gamma(j+s\alpha+1)} \\
 &= \int_0^t \sum_{s=0}^{\infty} (t-\tau)^{\alpha-1} \cdot \frac{[-A(t-\tau)^\alpha]^s}{\Gamma(s\alpha+\alpha)} f(\tau) d\tau + \sum_{j=0}^{m-1} b_j D^{(\alpha-j-1)} \left[\sum_{s=0}^{\infty} t^{\alpha-1} \frac{(-At^\alpha)^s}{\Gamma(s\alpha+\alpha)} \right] \\
 &= \int_0^t (t-\tau)^{\alpha-1} E_{\alpha,\alpha}(-A(t-\tau)^\alpha) f(\tau) d\tau + \sum_{j=0}^{m-1} b_j D^{(\alpha-j-1)} \left[t^{\alpha-1} E_{\alpha,\alpha}(-At^\alpha) \right] \\
 &= \int_0^t G_2(t-\tau) f(\tau) d\tau + \sum_{j=0}^{m-1} b_j D^{(\alpha-j-1)} G_2(t),
 \end{aligned} \tag{4.4}$$

where $G_2(t) = t^{\alpha-1} E_{\alpha,\alpha}(-At^\alpha)$.

It is easy to see that

$$y(t) = \int_0^t G_2(t-\tau) f(\tau) d\tau + b_0 D^{(\alpha-1)} G_2(t), \quad (0 < \alpha < 1). \tag{4.5}$$

(2). Consider an initial value problem for the nonhomogeneous Bagley-Torvik equation (see [5])

$$Ay''(t) + B^C D^{3/2}y(t) + Cy(t) = f(t), \quad t > 0, \quad (4.6)$$

$$y^{(i)}(0) = a_i, \quad i = 0, 1,$$

where a_i are real constants.

Utilizing Lemma 2.5 and applying ${}^C D^{-2}$ to both sides of (4.6), we obtain

$$y(t) + \frac{1}{A} [BD^{-1/2}y(t) + CD^{-2}y(t)]$$

$$= \frac{1}{A} D^{-2}f(t) + \sum_{i=0}^1 y^{(i)}(0) \frac{t^i}{\Gamma(i+1)} + \frac{B}{A} \sum_{i=0}^1 y^{(i)}(0) \frac{t^{1/2+i}}{\Gamma(3/2+i)} \quad (4.7)$$

$$= \frac{1}{A} D^{-2}f(t) + \sum_{i=0}^1 y^{(i)}(0) \left[\frac{t^i}{\Gamma(i+1)} + \frac{B}{A} \cdot \frac{t^{1/2+i}}{\Gamma(3/2+i)} \right]$$

According to the above procedure of solving the linear fractional differential equation with constant coefficients and using the Adomian decomposition method, let

$$y_0(t) = \frac{1}{A} D^{-2}f(t) + \sum_{i=0}^1 y^{(i)}(0) \left[\frac{t^i}{\Gamma(i+1)} + \frac{B}{A} \cdot \frac{t^{1/2+i}}{\Gamma(3/2+i)} \right],$$

$$y_1(t) = -\frac{1}{A} [BD^{-1/2}y_0(t) + CD^{-2}y_0(t)] = -\frac{C}{A} \left(\frac{B}{C} I + D^{-3/2} \right) D^{-1/2}y_0(t),$$

$$y_2(t) = -\frac{C}{A} \left(\frac{B}{C} I + D^{-3/2} \right) D^{-1/2}y_1(t) = \left(-\frac{C}{A} \right)^2 \left(\frac{B}{C} I + D^{-3/2} \right)^2 D^{-2/2}y_0(t), \quad (4.8)$$

$$\vdots$$

$$y_n(t) = -\frac{C}{A} \left(\frac{B}{C} I + D^{-3/2} \right) D^{-1/2}y_{n-1}(t) = \left(-\frac{C}{A} \right)^n \left(\frac{B}{C} I + D^{-3/2} \right)^n D^{-n/2}y_0(t),$$

$$\vdots$$

Adding all of the above terms, we obtain the solution of the equation by Adomian decomposition method as follows:

$$\begin{aligned}
y(t) &= \sum_{n=0}^{\infty} y_n(t) \\
&= \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \left(\frac{B}{C}I + D^{-3/2}\right)^n D^{-n/2} y_0(t) \\
&= \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \left(\frac{B}{C}I + D^{-3/2}\right)^n D^{-n/2} \left[\frac{1}{A} D^{-2} f(t) \right] \\
&\quad + \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \left(\frac{B}{C}I + D^{-3/2}\right)^n D^{-n/2} \sum_{i=0}^1 y^{(i)}(0) \left[\frac{t^i}{\Gamma(i+1)} + \frac{B}{A} \cdot \frac{t^{1/2+i}}{\Gamma(3/2+i)} \right] \\
&= \frac{1}{A} \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \sum_{k=0}^n C_n^{n-k} \left(\frac{B}{C}I\right)^{n-k} D^{-3k/2} D^{-n/2} D^{-2} f(t) \\
&\quad + \sum_{i=0}^1 a_i \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \sum_{k=0}^n C_n^{n-k} \left(\frac{B}{C}I\right)^{n-k} D^{-3k/2} D^{-n/2} \left[\frac{t^i}{\Gamma(i+1)} + \frac{B}{A} \cdot \frac{t^{1/2+i}}{\Gamma(3/2+i)} \right] \\
&= \frac{1}{A} \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \sum_{k=0}^n \frac{n!}{(n-k)!k!} \left(\frac{B}{C}I\right)^{n-k} D^{-(3k/2+n/2+2)} f(t) \\
&\quad + \sum_{i=0}^1 a_i \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \sum_{k=0}^n \frac{n!}{(n-k)!k!} \left(\frac{B}{C}I\right)^{n-k} D^{-(3k/2+n/2)} \left[\frac{t^i}{\Gamma(i+1)} + \frac{B}{A} \cdot \frac{t^{1/2+i}}{\Gamma(3/2+i)} \right] \\
&= \frac{1}{A} \int_0^t \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \sum_{k=0}^n \frac{n!}{(n-k)!k!} \left(\frac{B}{C}I\right)^{n-k} \frac{1}{\Gamma(3k/2+n/2+2)} (t-\tau)^{3k/2+n/2+2-1} f(\tau) d\tau \\
&\quad + \sum_{i=0}^1 a_i \sum_{n=0}^{\infty} \left(-\frac{C}{A}\right)^n \sum_{k=0}^n \frac{n!}{(n-k)!k!} \left(\frac{B}{C}I\right)^{n-k} \left[\frac{t^{i+3k/2+n/2}}{\Gamma(i+3k/2+n/2+1)} \right. \\
&\quad \quad \quad \left. + \frac{B}{A} \cdot \frac{t^{1/2+i+(3k/2)+(n/2)}}{\Gamma(i+3k/2+n/2+3/2)} \right] \\
&= \frac{1}{A} \int_0^t \sum_{k=0}^{\infty} \sum_{m=0}^{\infty} \left(-\frac{C}{A}\right)^{k+m} \frac{(k+m)!}{m!k!} \left(\frac{B}{C}I\right)^m \frac{1}{\Gamma(3/2k+(k+m)/2+2)} \\
&\quad \quad \quad \times (t-\tau)^{3k/2+(k+m)/2+1} f(\tau) d\tau \\
&\quad + \sum_{i=0}^1 a_i D^{(1-i)} \left[\sum_{k=0}^{\infty} \sum_{m=0}^{\infty} \left(-\frac{C}{A}\right)^{k+m} \frac{(k+m)!}{m!k!} \left(\frac{B}{C}I\right)^m \frac{t^{3k/2+(k+m)/2+1}}{\Gamma(3/2k+(k+m)/2+2)} \right] \\
&\quad + \sum_{i=0}^1 a_i D^{(1/2-i)} \left[\sum_{k=0}^{\infty} \sum_{m=0}^{\infty} \left(-\frac{C}{A}\right)^{k+m} \frac{(k+m)!}{m!k!} \left(\frac{B}{C}I\right)^m \frac{t^{3k/2+(k+m)/2+1}}{\Gamma(3/2k+(k+m)/2+2)} \right]
\end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{A} \int_0^t \sum_{k=0}^{\infty} \frac{(-C/A)^k}{k!} (t-\tau)^{2k+1} \sum_{m=0}^{\infty} \frac{(k+m)!}{m!} \cdot \frac{(-B/A)(t-\tau)^{1/2})^m}{\Gamma(k/2 + m/2 + 3k/2 + 2)} f(\tau) d\tau \\
 &+ A \sum_{i=0}^1 a_i D^{(1-i)} \left[\frac{1}{A} \sum_{k=0}^{\infty} \frac{(-C/A)^k}{k!} (t-\tau)^{2k+1} \sum_{m=0}^{\infty} \frac{(k+m)!}{m!} \cdot \frac{(-B/A)(t-\tau)^{1/2})^m}{\Gamma(k/2 + m/2 + 3k/2 + 2)} \right] \\
 &+ A \sum_{i=0}^1 a_i D^{(1/2-i)} \left[\frac{1}{A} \sum_{k=0}^{\infty} \frac{(-C/A)^k}{k!} (t-\tau)^{2k+1} \right. \\
 &\quad \left. \times \sum_{m=0}^{\infty} \frac{(k+m)!}{m!} \cdot \frac{(-B/A)(t-\tau)^{1/2})^m}{\Gamma((k/2) + (m/2) + (3k/2) + 2)} \right] \\
 &= \frac{1}{A} \int_0^t \sum_{k=0}^{\infty} \frac{(-C/A)^k}{k!} (t-\tau)^{2k+1} E_{1/2,3k/2+2}^{(k)} \left(-\frac{B}{A} (t-\tau)^{1/2} \right) f(\tau) d\tau \\
 &+ A \sum_{i=0}^1 a_i D^{(1-i)} \left[\frac{1}{A} \sum_{k=0}^{\infty} \frac{(-C/A)^k}{k!} t^{2k+1} E_{1/2,3k/2+2}^{(k)} \left(-\frac{B}{A} t^{1/2} \right) \right] \\
 &+ A \sum_{i=0}^1 a_i D^{(1/2-i)} \left[\frac{1}{A} \sum_{k=0}^{\infty} \frac{(-C/A)^k}{k!} t^{2k+1} E_{1/2,3k/2+2}^{(k)} \left(-\frac{B}{A} t^{1/2} \right) \right] \\
 &= \int_0^t G_3(t-\tau) f(\tau) d\tau + A \sum_{i=0}^1 a_i \left[D^{(1-i)} G_3(t) + D^{(1/2-i)} G_3(t) \right],
 \end{aligned}$$

(4.9)

where $G_3(t) = (1/A) \sum_{k=0}^{\infty} ((-C/A)^k / k!) t^{2k+1} E_{1/2,3k/2+2}^{(k)} (-B/A) t^{1/2}$.

Acknowledgment

The authors wish to thank the anonymous referees for their very careful reading of the manuscript and fruitful comments and suggestions. This work was supported by the National Natural Science Foundation of China (No. 60850005), the Natural Science Foundation of Zhejiang Province (Nos. D7080080 and Y607128) and the Innovation Team Foundation of the Department of Education of Zhejiang Province (No. T200924).

References

- [1] I. Podlubny, *Fractional Differential Equations: An Introduction to Fractional Derivatives, Fractional Differential Equations, to Methods of Their Solution and Some of Their Application*, vol. 198 of *Mathematics in Science and Engineering*, Academic Press, San Diego, Calif, USA, 1999.
- [2] K. Diethelm and N. J. Ford, "Numerical solution of the Bagley-Torvik equation," *BIT*, vol. 42, no. 3, pp. 490–507, 2002.
- [3] K. S. Miller and B. Ross, *An Introduction to the Fractional Calculus and Fractional Differential Equations*, John Wiley & Sons, New York, NY, USA, 1993.
- [4] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi, *Higher Transcendental Functions. Vol. III*, McGraw-Hill, London, UK, 1955.
- [5] S. S. Ray and R. K. Bera, "Analytical solution of the Bagley Torvik equation by Adomian decomposition method," *Applied Mathematics and Computation*, vol. 168, no. 1, pp. 398–410, 2005.

- [6] K. Diethelm and N. J. Ford, "Analysis of fractional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 265, no. 2, pp. 229–248, 2002.
- [7] K. Diethelm, "An algorithm for the numerical solution of differential equations of fractional order," *Electronic Transactions on Numerical Analysis*, vol. 5, pp. 1–6, 1997.
- [8] J. Biazar, E. Babolian, and R. Islam, "Solution of the system of ordinary differential equations by Adomian decomposition method," *Applied Mathematics and Computation*, vol. 147, no. 3, pp. 713–719, 2004.
- [9] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives: Theory and Applications*, Gordon and Breach, Yverdon, Switzerland, 1993.
- [10] B. J. West, M. Bologna, and P. Grigolini, *Physics of Fractal Operators*, Springer, New York, NY, USA, 2003.
- [11] T. M. Atanackovic and B. Stankovic, "On a system of differential equations with fractional derivatives arising in rod theory," *Journal of Physics A*, vol. 37, no. 4, pp. 1241–1250, 2004.
- [12] Y. Hu, Y. Luo, and Z. Lu, "Analytical solution of the linear fractional differential equation by Adomian decomposition method," *Journal of Computational and Applied Mathematics*, vol. 215, no. 1, pp. 220–229, 2008.
- [13] G. Adomian, "A review of the decomposition method in applied mathematics," *Journal of Mathematical Analysis and Applications*, vol. 135, no. 2, pp. 501–544, 1988.
- [14] G. Adomian, *Solving Frontier Problems of Physics: The Decomposition Method*, vol. 60 of *Fundamental Theories of Physics*, Kluwer Academic Publishers, Dordrecht, The Netherlands, 1994.
- [15] V. Daftardar-Gejji and H. Jafari, "Adomian decomposition: a tool for solving a system of fractional differential equations," *Journal of Mathematical Analysis and Applications*, vol. 301, no. 2, pp. 508–518, 2005.
- [16] J.-H. He, G.-Ch. Wu, and F. Austin, "The variational iteration method which should be followed," *Nonlinear Science Letters A*, vol. 1, no. 1, pp. 1–30, 2010.
- [17] S. Barat, S. Das, and P. K. Gupta, "A note on fractional schrodinger equation," *Nonlinear Science Letters A*, vol. 1, no. 1, pp. 91–94, 2010.
- [18] A. E. Ebaid, "Exact orbits of planetary motion using the Adomian decomposition method," *Nonlinear Science Letters A*, vol. 1, no. 3, pp. 249–252, 2010.
- [19] S. Das, "Solution of fractional vibration equation by the variational iteration method and modified decomposition method," *International Journal of Nonlinear Sciences and Numerical Simulation*, vol. 9, no. 4, pp. 361–366, 2008.
- [20] A. Golbabai and K. Sayevand, "The homotopy perturbation method for multi-order time fractional differential equations," *Nonlinear Science Letters A*, vol. 1, no. 2, pp. 147–154, 2010.
- [21] Y. Keskin and G. Oturanc, "The reduced differential transform method: a new approach to factional partial differential equations," *Nonlinear Science Letters A*, vol. 1, no. 2, pp. 207–217, 2010.



Hindawi

Submit your manuscripts at
<http://www.hindawi.com>

