



# Enumeration of Basic Ideals in Type $B$ Lie Algebras

Jonathan Nilsson

Department of Mathematics

Uppsala University

P. O. Box 480

SE-751 06, Uppsala

Sweden

[jonathan.nilsson@math.uu.se](mailto:jonathan.nilsson@math.uu.se)

## Abstract

The number of ad-nilpotent ideals of the Borel subalgebra of the classical Lie algebra of type  $B_n$  is determined using combinatorial arguments involving a generalization of Dyck-paths. We also solve a similar problem for the untwisted affine Lie algebra of type  $\tilde{B}_n$ , where we instead enumerate a certain class of ideals called *basic ideals*. This leads to an explicit formula for the number of basic ideals in  $\tilde{B}_n$ , which gives rise to a new integer sequence.

## 1 Introduction

Catalan numbers enumerate many objects appearing in various parts of mathematics, see Stanley [11]. One algebraic example comes from Lie theory, where Catalan numbers enumerate ad-nilpotent ideals contained in the Borel subalgebra of  $\mathfrak{sl}_n(\mathbb{C})$ , see for example Cellini and Papi [5]. In the present paper we solve a similar enumeration problem and arrive at a new integer sequence which gives the number of so called basic ideals in a certain family of simple Lie algebras. More specifically, let  $\mathfrak{g}$  be a finite dimensional simple complex Lie algebra with a fixed triangular decomposition

$$\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$$

and let  $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}_+$  be the corresponding Borel subalgebra. An ideal  $\mathfrak{i}$  of  $\mathfrak{b}$  is called *ad-nilpotent* provided that the adjoint action of each element of  $\mathfrak{i}$  on  $\mathfrak{b}$  is nilpotent. When  $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$ ,

the set of strictly upper triangular matrices is a typical example of an ad-nilpotent ideal, but there are many other such ideals contained in this one. This leads to the problem of enumerating ad-nilpotent ideals, which has been done in various contexts. See for example the work of Cellini and Papi [5, 6], Shi [9], Athanasiadis [1, 2] and Krattenthaler, Orsina, and Papi [8]. For example, as mentioned above, when  $\mathfrak{g} = \mathfrak{sl}_n(\mathbb{C})$ , the number of ad-nilpotent ideals in  $\mathfrak{b}$  is given by the  $n$ 'th Catalan number  $\frac{1}{n+1} \binom{2n}{n}$ .

In the case when  $\mathfrak{g}$  is an untwisted affine Lie algebra (as defined in Section 2), any ad-nilpotent ideal of  $\mathfrak{b}$  is contained in the center of  $\mathfrak{g}$ , and the latter is 1-dimensional, so the corresponding enumeration problem becomes trivial. Baur and Mazorchuk [3] formulated another problem, where instead of ad-nilpotent ideals, one considers the so called *basic ideals* (see the definition in Section 4). The enumeration problem for basic ideals was solved for the affine algebra  $\tilde{\mathfrak{sl}}_n$  which led to a new integer sequence.

In the present paper we solve the enumeration problem for basic ideals for untwisted affine Lie algebras of affine type  $\tilde{B}_n$ . We prove the following statement.

**Theorem 1.** *The number  $\tilde{b}_n$  of basic ideals in the Lie algebra of type  $\tilde{B}_n$  is given by the formula*

$$\tilde{b}_n = (3n + 5)2^{2n-2} - 2(3n - 1) \binom{2n-2}{n-1}.$$

To prove Theorem 1 we establish a combinatorial scheme in which basic ideals for type  $B$  are encoded using certain pairs of type  $B$ -analogues for Dyck-paths. This gives an explicit but complicated formula for the number of basic ideals. The simplification of this expression to the simple form in Theorem 1 is a nontrivial combinatorial computation which was performed by Christian Krattenthaler. The above simple formula, in particular, implies that the sequence  $\tilde{b}_n$  satisfies the following non-homogeneous linear recurrence relation.

$$\tilde{b}_n - 8\tilde{b}_{n-1} + 16\tilde{b}_{n-2} = \frac{24}{n-1} \binom{2n-6}{n-2}, \quad n \geq 4.$$

We do not have any direct combinatorial explanation for this recursion.

The paper is organized as follows. Section 2 contains some of the established Lie theory regarding ideals and the structure of affine untwisted Lie algebras. The concepts of basic and combinatorial ideals are also introduced here following the definition of Bauer and Mazorchuk [3]. In Section 3 we focus on finite dimensional Lie algebras of type  $B$ . We introduce *B-paths*, an analogue for Dyck paths in type  $B$ , and show that the set of such paths bijectively corresponds to the set of ad-nilpotent ideals in type  $B$ . We use this to show that the number of ad-nilpotent ideals in the Lie algebra of type  $B_n$  equals  $\binom{2n}{n}$ , the type  $B$  Catalan number, which agrees with a previous result [9, 5, 2]. After this we want to generalize these ideas to the affine case. In Section 4 we study the root system structure of the affine Lie algebras of type  $B$  and in Section 5 we show that there is a bijective correspondence between basic ideals and a certain set of pairs of  $B$ -paths called *admissible* pairs. This reduces our problem to an enumeration of the admissible pairs. Finally, in Section 6 we write down an explicit formula for the number of admissible pairs by considering a number of cases. We then proceed to simplify this expression using a number of combinatorial lemmas, and after a lot of simplification we arrive at the short expression of Theorem 1.

## 2 Preliminaries

Throughout this section, let  $\mathfrak{g}$  be a finite dimensional simple Lie algebra with Cartan subalgebra  $\mathfrak{h}$ . Let  $\Phi$  denote the corresponding (finite) root system and let  $\Delta$  be a fixed base of  $\Phi$ . Then the Lie algebra is the direct sum of its root spaces  $\mathfrak{g} = \mathfrak{h} \oplus \left( \bigoplus_{\alpha \in \Phi} \mathfrak{g}_\alpha \right)$ , moreover these root spaces are all one-dimensional.

### 2.1 Presentation of ideals

The set  $\Phi$  is ordered by defining  $\alpha \prec \beta$  whenever  $\beta - \alpha \in \text{span}_{\mathbb{Z}^+}(\Delta)$ , in other words, whenever  $\beta - \alpha$  is a sum of simple roots. The set of roots splits into two parts with respect to this order. By defining  $\Phi^+ := \{\alpha \in \Phi | \alpha \succ 0\}$  and  $\Phi^- := \{\alpha \in \Phi | \alpha \prec 0\}$  and letting  $\mathfrak{n}_-$  and  $\mathfrak{n}_+$  be the sum of the corresponding root spaces, we have  $\mathfrak{g} = \mathfrak{n}_- \oplus \mathfrak{h} \oplus \mathfrak{n}_+$  which is called a triangular decomposition of  $\mathfrak{g}$ . The subalgebra  $\mathfrak{b} = \mathfrak{h} \oplus \mathfrak{n}_+$  is called the Borel subalgebra. If  $\mathfrak{i}$  is a nonzero ideal of  $\mathfrak{b}$  it is  $\mathfrak{h}$ -stable, hence it is the direct sum of entire root spaces. Now, if  $x \in \mathfrak{g}_\alpha \subset \mathfrak{i}$  and  $y \in \mathfrak{g}_\beta$ , then  $\mathfrak{i}$  also contains  $[x, y]$  which lies in  $\mathfrak{g}_{\alpha+\beta}$  so  $\mathfrak{i} \supset \mathfrak{g}_{\alpha+\beta}$ . Continuing this inductively letting  $\beta$  vary over simple roots we see that in fact  $x \in \mathfrak{g}_\alpha \subset \mathfrak{i}$  and  $\beta \succ \alpha$  implies  $\mathfrak{i} \supset \mathfrak{g}_\beta$ . This shows that ad-nilpotent ideals of the Borel-subalgebra corresponds to coideals of the poset  $\Phi^+$ . This fact is used in Section 3 to enumerate the number of ad-nilpotent ideals when the Lie algebra is of type  $B$ .

### 2.2 Affine Lie algebras

Given the Lie algebra  $\mathfrak{g}$  there is a natural way of constructing a corresponding infinite-dimensional Lie algebra  $\tilde{\mathfrak{g}}$ . The algebra  $\tilde{\mathfrak{g}}$  is obtained as follows. First consider the loop algebra  $\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]$  endowed with the bracket  $[x \otimes t^m, y \otimes t^n] = [x, y] \otimes t^{m+n}$ . The universal central extension of the Loop algebra is explicitly,  $(\mathfrak{g} \otimes \mathbb{C}[t, t^{-1}]) \oplus \mathbb{C}c$  where the bracket is modified as follows:

$$[x \otimes t^m + \lambda c, y \otimes t^n + \mu c] = [x, y] \otimes t^{m+n} + m(x|y)\delta_{m+n,0}c$$

where  $(\cdot|\cdot)$  is the Killing form of  $\mathfrak{g}_n$ . The algebra  $\tilde{\mathfrak{g}}$  is now obtained by extending the latter by a derivation  $d$  which acts as follows  $[d, (x \otimes t^m)] = m(x \otimes t^m)$  and  $[d, c] = 0$ . The resulting Lie algebra  $\tilde{\mathfrak{g}}$  is called the *untwisted affine Kac-Moody algebra* corresponding to  $\mathfrak{g}$ . The algebra  $\tilde{\mathfrak{g}}$  has a triangular decomposition  $\tilde{\mathfrak{g}} = \tilde{\mathfrak{n}}_- \oplus \tilde{\mathfrak{h}} \oplus \tilde{\mathfrak{n}}_+$ , where

$$\tilde{\mathfrak{n}}_+ = \text{Span}(\{x \otimes 1 | x \in \mathfrak{n}_+\} \cup \{x \otimes t^k | x \in \mathfrak{g}, k > 0\})$$

$$\tilde{\mathfrak{n}}_- = \text{Span}(\{y \otimes 1 | y \in \mathfrak{n}_-\} \cup \{y \otimes t^k | y \in \mathfrak{g}, k < 0\})$$

$$\tilde{\mathfrak{h}} = \text{Span}(\{c, d\} \cup \{h \otimes 1 | h \in \mathfrak{h}\}).$$

Define  $\delta \in \tilde{\mathfrak{h}}^*$  by  $\delta(d) = 1$ ,  $\delta(c) = 0$  and  $\delta(h \otimes t^k) = 0$  for any  $h \in \mathfrak{h}$ ,  $t \in \mathbb{Z}$ . Then  $\delta$  is a root of  $\tilde{\mathfrak{g}}$ , called the indivisible positive imaginary root. For  $x \in \mathfrak{g}_\alpha$  we see that  $x \otimes t^k$  lies in the root space  $\tilde{\mathfrak{g}}_{\alpha+k\delta}$  (where we identify  $\alpha \in \mathfrak{h}^*$  with its trivial extension to  $\tilde{\mathfrak{h}}^*$ ). Thus the root system of  $\tilde{\mathfrak{g}}$  is  $(\Delta + \mathbb{Z}\delta) \cup \mathbb{N}\delta \cup -\mathbb{N}\delta$ , in particular the root system is infinite. If  $\alpha_{max}$  is the maximal root of  $\Phi$ , then  $\{\delta - \alpha_{max}\} \cup \Delta$  is a basis for the root system of  $\tilde{\mathfrak{g}}$ .

We shall need the following lemma regarding the ordering of the roots.

**Lemma 2.** *If  $\alpha, \beta$  are positive roots of  $\mathfrak{g}$  and let  $\alpha_{max}$  be the maximal root of  $\mathfrak{g}$ , then  $\delta - \alpha \succ \beta$  if and only if  $\alpha + \beta \prec \alpha_{max}$ , and equality holds in the latter relation if and only if  $\delta - \alpha$  covers  $\beta$ .*

*Proof.* We have  $\delta - \alpha \succ \beta$  if and only if  $\alpha + \beta \prec \alpha_0 + \alpha_{max}$ . Writing each root of this inequality in the base given above we see that the  $\alpha_0$  coordinate is 0 on the left and 1 on the right, hence this is equivalent to  $\alpha + \beta \prec \alpha_{max}$ . Moreover  $\alpha + \beta = \alpha_{max}$  is equivalent to  $\delta - \alpha = \beta + \alpha_0$ , in other words,  $\delta - \alpha$  covers  $\beta$ .  $\square$

The corresponding Borel subalgebra  $\tilde{\mathfrak{b}} = \tilde{\mathfrak{h}} \oplus \tilde{\mathfrak{n}}_+$  does not contain any nontrivial ad-nilpotent ideals (outside the center  $Z(\mathfrak{b})$  of  $\mathfrak{b}$ ). For, if such an ideal  $\mathfrak{i}$  existed, containing an element  $x \in \mathfrak{b} \setminus Z(\mathfrak{b})$  we could commute it with positive root elements to obtain  $y \in \mathfrak{i}$  lying in a root space corresponding to a positive root:  $y \in \mathfrak{g}_\alpha + \mathfrak{k}\delta$  with  $k \geq 0$ . If  $\alpha \succ 0$  we can commute further with a positive root vector from  $\mathfrak{g}_{\delta-\alpha}$  to see that  $\mathfrak{i}$  includes  $\mathfrak{g}_{(k+1)\delta}$ . If  $\alpha \prec 0$ ,  $k$  is positive and by commuting with a positive root vector from  $\mathfrak{g}_{-\alpha}$  we see that  $\mathfrak{i}$  includes  $\mathfrak{g}_{k\delta}$ . If  $\alpha = 0$  we have  $k > 0$  and it is obvious that  $\mathfrak{i}$  includes  $\mathfrak{g}_{k\delta}$ . In any case,  $\mathfrak{i}$  contains an element  $z$  belonging to an imaginary root space. But then  $ad z$  is clearly not nilpotent, contradicting the existence of nontrivial ad-nilpotent ideals of  $\tilde{\mathfrak{b}}_{\mathfrak{n}}$ .

## 2.3 Combinatorial and basic ideals

Lacking ad-nilpotent ideals we can instead consider a combinatorial analogue, called *combinatorial ideals* which were introduced by Bauer and Mazorchuk [3]. A combinatorial ideal of  $\mathfrak{b}$  is an ideal  $\mathfrak{i}$  of  $\tilde{\mathfrak{b}}$  contained in  $\tilde{\mathfrak{n}}_+$  which has finite codimension and is the union of entire root spaces:  $\mathfrak{i} \cap \tilde{\mathfrak{g}}_\alpha \in \{\{0\}, \tilde{\mathfrak{g}}_\alpha\}$  for each positive root  $\alpha$ . For any combinatorial ideal  $\mathfrak{i}$  we define  $\text{supp}(\mathfrak{i})$ , called the *support* of  $\mathfrak{i}$ , to be the set of positive roots whose root spaces are contained in  $\mathfrak{i}$ , that is,  $\text{supp}(\mathfrak{i}) = \{\alpha \in \tilde{\mathfrak{h}}^* | \tilde{\mathfrak{g}}_\alpha \subset \mathfrak{i}\}$ .

The set of combinatorial ideal is infinite. This leads us to consider an equivalence relation on the set of combinatorial ideals by defining  $\mathfrak{i} = \bigoplus_{\alpha \in \text{supp}(\mathfrak{i})} \mathfrak{g}_\alpha$  to be equivalent to all combinatorial ideals of form  $\bigoplus_{\alpha \in \text{supp}(\mathfrak{i})} \mathfrak{g}_{\alpha+k\delta}$  for  $k \in \mathbb{Z}$ . We define the finite subset  $D$  of the root system by  $D := \{\delta\} \cup \Phi^+ \cup (\Phi^- + \delta)$ . Note that  $D$  is precisely the principal order ideal generated by  $\delta$ . The combinatorial ideals whose support intersects  $D$  non trivially are called *basic ideals*. One can easily show [3] that the set of basic ideals contains precisely one element of each equivalence class of combinatorial ideals defined above. The set of basic ideals is clearly finite as its elements can be identified with a subsets of  $D$ , and in this paper we determine its size when  $\mathfrak{g}$  is a simple Lie algebra of type  $B$  and rank  $n$ , see Section 4 and onward.

## 3 The simple Lie algebra of type $B$

### 3.1 Root system structure

Following the notation of Bourbaki [4], the root system of type  $B_n$  can be constructed in  $\mathbb{R}^n$  as follows. Let  $\{e_i\}_{i=1}^n$  be the standard orthogonal basis for  $\mathbb{R}^n$ . For  $1 \leq i \leq n-1$  define

$\alpha_i = e_i - e_{i+1}$ , and let  $\alpha_n = e_n$ . Then the (simple roots)  $\alpha_1, \dots, \alpha_n$  constitutes a base for a root system of type  $B_n$  where the positive roots are given by

$$\sum_{j \leq i \leq k} \alpha_i, \quad 1 \leq j \leq k \leq n; \quad \sum_{j \leq i \leq k} \alpha_i + \sum_{k+1 \leq i \leq n} 2\alpha_i, \quad 1 \leq j < k < n.$$

We identify each such root with the integer pair  $(r, s)$ , such that  $r$  is the minimal value of  $i$  for which the  $\alpha_i$ -coefficient of the root in the simple basis is non-zero, and  $s = r + h - 1$  where  $h$  is the height of the root (the sum of its coefficients in the simple basis). For example, in  $B_7$  we write  $(3, 10)$  for the root  $(0, 0, 1, 1, 2, 2, 2)$ , and  $(2, 5)$  for  $(0, 1, 1, 1, 1, 0, 0)$ .

Positive roots are partially ordered via

$$\alpha \prec \beta \iff \beta - \alpha \text{ is a } \mathbb{Z}_{\geq 0}\text{-linear combination of simple roots.}$$

This order can of course be extended to all of  $\mathfrak{h}^*$ . It is easy to check that  $(r, s) \prec (\hat{r}, \hat{s})$  if and only if  $\hat{r} \leq r$  and  $\hat{s} \geq s$ . The  $(\prec)$ -poset structure on positive roots can now be visualized in a modified version of the Hasse-diagram. Consider a diagram consisting of rows and columns. If  $(i, j)$  is a root then we write it in the  $i$ 'th row (from top to bottom) and  $j$ 'th column (from left to right), in other words, we write  $(i, j)$  in the  $(i, j)$ 'th position. Then  $\prec$ -greater roots are always in the north and east directions.

Not all positions in the diagram will contain roots. We obviously have  $i \geq 1$  and  $j \geq 1$ . We must also have  $i \leq j$ , since positive roots have positive height, and we must have  $i \leq 2n - j$  (compare the original expression of a general positive root). Thus, all roots lie in a triangular scheme. For example, in  $B_4$  the positive roots are ordered as in Figure 1.




Figure 1: Positive roots in  $B_4$

Here  $\alpha \rightarrow \beta$  means that  $\beta$  covers  $\alpha$  with respect to  $\prec$ , that is  $\beta$  is a minimal element of the set  $\{\gamma | \alpha \prec \gamma\}$ . Thus we have  $\alpha \prec \beta$  if and only if there is a directed path from  $\alpha$  to  $\beta$ .

Any ad-nilpotent ideal  $\mathfrak{i}$  in  $\mathfrak{n}_+$  can be decomposed as a direct sum of some root spaces corresponding to positive roots. If  $\mathfrak{i}$  contains some nonzero  $x$  from a root space  $L_\gamma$ , it contains



*Proof.* Let  $\mathfrak{i}$  be an ad-nilpotent ideal in  $\mathfrak{b}_n$ . Consider the  $(2n - 1) \times n$  rectangular picture described above in which boxes are identified with positive roots of  $\mathfrak{g}_n$ . If  $\mathfrak{i}$  contains a root element for some root  $\alpha$ , the fact that  $\mathfrak{i}$  is an ideal implies that  $\mathfrak{i}$  contains root elements for all roots  $\beta$  such that  $\beta \succ \alpha$ . From this it follows that there is a unique path  $p = p(\mathfrak{i})$  which has the following properties.

- $p$  starts at  $(-1, 0)$ ;
- each step of  $p$  goes to the right (along  $(1, 0)$ ) or down (along  $(0, -1)$ );
- $p$  terminates at a point of the form  $(2n - 1 - x, -x)$ ,  $x \in \{0, 1, \dots, n\}$ ;
- $p$  separates all boxes corresponding to elements of  $\mathfrak{i}$  from all other boxes.

Substituting each  $(1, 0)$ -step in  $p$  with  $r$  and each  $(0, -1)$ -step in  $p$  with  $f$ , we obtain a  $B_n$ -path.

Conversely, given a  $B_n$ -path we make the reverse substitution and obtain a path satisfying the first three conditions above. It is easy to see that the linear span of root elements corresponding to the roots lying to the northeast of this path is an ad-nilpotent ideal of  $\mathfrak{b}_n$ . The claim follows.  $\square$

Here is an explicit example in type  $B_4$ . The  $B_4$ -path  $rrrfrfrr$  is drawn below; it corresponds to the ideal spanned by the root spaces of the roots  $(1, 3)$ ,  $(1, 4)$ ,  $(1, 5)$ ,  $(1, 6)$ ,  $(1, 7)$ ,  $(2, 4)$ ,  $(2, 5)$  and  $(2, 6)$ . Also note that any pair of (non-zero) elements from the root spaces of the roots  $(1, 3)$  and  $(2, 4)$  generates this ideal.




Figure 3: An example of the  $B_4$ -path  $rrrfrfrr$

The path  $rfrfrrfrf$  gives the improper coideal containing all the roots, i.e. the one corresponding to the Borel subalgebra. The path  $rrrrrrrr$  corresponds to the empty coideal containing no roots.

We can now state a corollary regarding the number of ad-nilpotent ideals in  $B_n$ . This result was proved in a similar way (but less detailed) by Krattenthaler, Orsina, and Papi [8].

**Corollary 6.** *The number of  $B_n$ -paths, hence also the number of ad-nilpotent ideals in  $\mathfrak{b}_n$ , equals  $\binom{2n}{n}$ .*





roots of form  $d + k\delta$  for  $d \in D$ ,  $k \geq 0$ , and the negative roots correspond to  $k < 0$  (except 0 which is not a root at all).

We now describe the poset structure of  $D$  in terms of its partition above. We have  $D_1 = \{\delta\}$  and  $\delta$  is the maximum element of the poset  $D$ .  $D_2$  is the set of positive roots of  $B_n$  which is fully embedded into  $D$  as a poset. Thus the poset structure on  $D_2$  is precisely as in Section 3.  $D_3$  consists of roots of form  $\delta - \alpha$  where  $\alpha$  is a positive root of  $B_n$ . These are ordered in the reversed way, since  $\alpha \prec \beta \Leftrightarrow \delta - \alpha \succ \delta - \beta$ . The roots of  $D_3$  can be organized into a digram similarly to how we organized positive roots of  $B_n$  but reflected with respect to the horizontal line. More explicitly, in the position  $(i, j)$  we write  $\delta - (n - i + 1, 2n - j)$  whenever  $(n - i + 1, 2n - j)$  is a positive root. Then for  $\hat{i} \leq i$  and  $\hat{j} \geq j$ ,  $\delta - (n - \hat{i} + 1, 2n - \hat{j}) \succ \delta - (n - i + 1, 2n - j)$ , so greater elements are again in the north and east directions. Putting the diagrams  $D_2$  and  $D_3$  together we obtain a diagram which is illustrated below in the case  $n = 4$ .




Figure 5:  $D_2$  and  $D_3$

However, we have yet to consider  $\prec$ -relations between  $D_2$  and  $D_3$  in the diagram. Because of the  $\alpha_0$ -coordinate, a root in the upper part can not be  $\prec$ -smaller than a root in the lower part. Applying Lemma 2 we obtain the following result which specifies precisely when a root in the lower part is  $\prec$ -smaller than a root in the upper part.

**Corollary 7.** *For  $i > 1$  we have  $(i, j) \prec \delta - (\hat{i}, \hat{j})$  if and only if  $\hat{j} \leq 2n - j$ , while  $(1, j) \prec \delta - (\hat{i}, \hat{j})$  if and only if  $\hat{j} \leq 2n - j$  and  $\hat{i} > 1$ .*

Let us translate this result to our diagram above. Since the  $(i, j)$ 'th position in the upper part contains the root  $\delta - (n - i + 1, 2n - j)$ , any element in the  $j$ 'th column (except the one

in the first row) at the lower part is  $\prec$ -smaller than all elements in the  $j$ 'th column in the upper part. The root  $(1, j)$  in the first row and  $j$ 'th column is  $\prec$ -smaller than all elements in the  $j$ 'th column above, except for the one in the bottom row. For example, our  $\tilde{B}_4$  diagram is now completed as follows.




Figure 6: Completed  $\tilde{B}_4$  diagram

Here again, a root is  $\prec$ -smaller than another root if and only if there is a path of arrows from the first one to the second one. Note that the top row of the bottom part is linked to the second to bottom row in the upper part and vice versa.

Our goal is now to express the number of coideals in such a poset (for an arbitrary  $n$ ) in terms of pairs of paths. To this end, construct again a rectangle in  $\mathbb{R}^2$ , this time with the northwest corner in  $(0, n + 1)$  and the southeast corner in  $(2n - 1, -n)$ , where  $1 \times 1$ -blocks correspond to roots of  $D \setminus \{\delta\}$ . The rectangle contains the two triangles in a natural way as illustrated below in the case  $n = 4$ .

We have drawn two paths in the picture, a red one and a blue one. These paths together specify a coideal of the poset in the following way: we take all roots corresponding to boxes to the northeast of the blue path together with all roots corresponding to boxes from the lower triangle to the northeast of the red path. Every coideal can be obtained this way, but the converse is not true, some choices of paths do not correspond to coideals. For example, in the above diagram, since the red path specifies that the root in box marked by (1) belongs to the coideal, the root in the box marked by (2) in the top triangle should be in the coideal as well. Thus the blue path can not be taken “too high” given the red path. The choice in the picture above indeed does correspond to a coideal. A pair of paths which do correspond to a coideal will be called an *admissible* pair of paths.



- each step of  $q$  goes to the left (along  $(-1, 0)$ ) or up (along  $(0, 1)$ );
- $q$  terminates at a point of the form  $(y, y + 1)$ ,  $y \in \{0, 1, \dots, n\}$ ;
- $q$  separates all boxes corresponding to elements of  $D_3 \cap \text{supp}(\mathbf{i})$  from boxes corresponding to elements of  $D_3 \setminus \text{supp}(\mathbf{i})$ .

By substituting each  $(1, 0)$ -step in  $p$  with  $r$  and each  $(0, -1)$ -step in  $p$  with  $f$  we obtain a  $B_n$ -path  $p(\mathbf{i})$ , and by substituting each  $(-1, 0)$ -step in  $q$  with  $r$  and each  $(0, 1)$ -step in  $q$  with  $f$  we obtain another  $B_n$ -path  $q(\mathbf{i})$ . Thus, with the basic ideal  $\mathbf{i}$  we have associated a pair of  $B_n$ -paths  $(p(\mathbf{i}), q(\mathbf{i}))$ . It is also clear that different basic ideals give rise to different pairs of paths.

However, as mentioned before, not every pair  $(p, q)$  corresponds to a basic ideal. All pairs of the form  $(p(\mathbf{i}), q(\mathbf{i}))$  for some basic ideal  $\mathbf{i}$  are called *admissible*. For example, the pair  $(rrrfrfrr, rrrrrrff)$  is admissible; it is precisely the pair from the previous picture.

We now specify the conditions for a pair  $(p, q)$  of  $B_n$ -paths to be admissible.

For  $1 \leq i < j \leq 2n$  define  $\mathfrak{B}_n(i, j)$  as the set of all  $B_n$ -paths in which the first and the second occurrences of  $f$  (reading the word from left to right) are at the positions  $i$  and  $j$ . Also, for  $1 \leq i \leq 2n$  let  $\mathfrak{B}_n(i)$  be the set of paths where the first occurrence of  $f$  is at position  $i$ . For example  $rrrfrfff \in \mathfrak{B}_4(4, 6) \subset \mathfrak{B}_4(4)$  and  $rrrfff \in \mathfrak{B}_3(4, 5) \subset \mathfrak{B}_3(4)$ . Note that  $B_n$  paths also are defined for  $n = 1$ , and we have  $|\mathfrak{B}_1(2)| = |\mathfrak{B}_1(3)| = 1$ . We shall also abuse notation by defining  $\{rr \cdots rr\} = \mathfrak{B}_n(2n + 1)$ , that is, if a path contains no  $f$ , its first  $f$  can be thought of to be one step after the last letter of the path. Similarly, write  $\{rr \cdots rr\} = \mathfrak{B}_n(2n + 1, 2n + 2)$  and  $rr \cdots rfr \cdots rr \in \mathfrak{B}_n(k, 2n + 1)$  (here the only occurrence of  $f$  is at place  $k$ ).

Using the notation above we can state the following criterion for a pair of  $B_n$ -paths  $(p, q)$  to be admissible.

**Proposition 8.** *Let  $p \in \mathfrak{B}_n(a, b)$  where  $3 \leq a < b \leq 2n$ . Then  $(p, q)$  is an admissible pair of paths if and only if  $q \in \mathfrak{B}_n(i, j)$  where  $i \geq 2n + 4 - b$  and  $j \geq 2n + 4 - a$ .*

*Proof.* Let  $3 \leq a < b \leq 2n$  and  $p \in \mathfrak{B}_n(a, b)$ . Realize  $(p, q)$  as a pair of paths in the diagram described above. By Proposition 5, the path  $p$  specifies a coideal of the poset  $D_2$  and  $q$  specifies a coideal of the poset  $D_3$ , so we need only consider  $\prec$ -arrows between the two triangles. Such arrows originate from the two upper rows of the lower diagram. The root corresponding to the box whose northwest corner is in  $(a - 2, 0)$  is in the coideal, and it  $(\prec)$ -covers the root corresponding to the box with northwest corner in  $(a - 2, 3)$ , so this latter root must also belong to the coideal. Similarly, since the root corresponding to the box with northwest corner  $(b - 3, -1)$  is in the coideal, the root corresponding to the box with northwest corner in  $(b - 3, 2)$  must also be in the coideal. Thus, for  $p$  as above,  $(p, q)$  is admissible if and only if the two boxes with northwest corners  $(a - 2, 3)$  and  $(b - 3, 2)$  respectively lie to the northwest of the path  $q$ . This is equivalent to  $q \in \mathfrak{B}_n(i, j)$  with  $i \geq 2n + 4 - b$  and  $j \geq 2n + 4 - a$ .  $\square$

We illustrate the proposition above with an example.

**Example 9.** Let  $n = 6$  and let  $p = rrrrfrrfrfrr \in \mathfrak{B}_6(5, 8)$ . The path  $p$  is drawn in red in the bottom triangle of the following picture.




Figure 8: A pair of paths

The box marked by (1) is linked to the box marked (1') linked via the  $\prec$ -relation, and the box marked by (2) is linked to the box marked (2') linked via the  $\prec$ -relation. Hence  $q$  must be chosen such that the boxes marked (1') and (2') are both to the northeast of the path  $q$ . In other words,  $q \in \mathfrak{B}_6(i, j)$  where  $i \geq 2n + 4 - b = 8$  and  $j \geq 2n + 4 - a = 11$ . The smallest coideal allowed for this choice of the path  $p$  is of course given by  $q = (rrrrrrrrfrrff)$ , and the largest ideal corresponds to when  $q = rrrrrrrrrrrr$ . All paths “between” these also give admissible pairs  $(p, q)$ . One such choice,  $q = rrrrrrrrfrrr$ , is drawn in blue color in the picture.

**Corollary 10.** For  $p \in \mathfrak{B}_n(a, b)$  where  $3 \leq a < b \leq 2n$ , the number of  $q$ 's such that  $(p, q)$  is admissible is given by

$$\sum_{i \geq 2n+4-b} \sum_{j \geq 2n+4-a} |\mathfrak{B}_n(i, j)|.$$

Comparing with the picture of the poset though, we note that the argument above does not make sense when  $a = 2$  or when  $b = 2n + 1$ , so unfortunately we need to consider a few special cases as well.

## 6 Counting paths

In this section we prove Theorem 1. We start with the following lemma.

**Lemma 11.** *For  $n \geq 2$  and  $2 \leq i < j \leq 2n + 1$  we have*

$$|\mathfrak{B}_n(i, j)| = |\mathfrak{B}_{n-1}(j - 2)|.$$

*Proof.* For such  $i, j$  and  $n$ , define a map  $\phi : \mathfrak{B}_n(i, j) \rightarrow \mathfrak{B}_{n-1}(j - 2)$  which just deletes the first (leftmost)  $r$  and the first  $f$  from a given path. It is easy to check that  $\phi(p) \in \mathfrak{B}_{n-1}(j - 2)$  and that  $\phi$  is a bijection (its inverse just appends an  $r$  to the left and inserts an  $f$  at position  $i$ ). Hence the sets have the same cardinality.  $\square$

The numbers  $|\mathfrak{B}_n(i)|$  are given explicitly by the following lemma.

**Lemma 12.** *For  $n \geq 2$  and  $1 \leq i \leq 2n$  we have*

$$|\mathfrak{B}_n(i)| = \begin{cases} \sum_{k=1}^{n-1} 2^k \left( \binom{2n-i-k-1}{n-i} - \binom{2n-i-k-1}{n-i-k} \right), & \text{if } i \leq n; \\ 2^{2n-i}, & \text{if } i > n. \end{cases}$$

*Proof.* Each path of  $\mathfrak{B}_n(i)$  can be specified by first choosing a Dyck-path starting with  $r^{i-1}f$ , and then replacing its tail consisting of some number of  $f$ 's by any sequence of  $r$  and  $f$  of the same length. Baur and Mazorchuk [3, Proposition 13] showed that the number of Dyck-paths of semilength  $n$  with the first peak at height  $i$  and the last peak at height  $k$  is given by the expression

$$\binom{2n-i-k-2}{n-i-1} - \binom{2n-i-k-2}{n-i-k-1}.$$

Since a Dyck-path having the last peak at height  $k$  ends with  $f^k$ , it gives rise to  $2^k$  possible  $B$ -paths. Summing over  $k$  we obtain the number of all such  $B$ -paths. When  $i > n$ , this argument degenerates, but a path starting with  $r^{i-1}f$  with  $i > n$  can clearly be completed by choosing the remaining symbols arbitrarily, which gives  $2^{2n-i}$  possibilities. Here is a geometric illustration.




Figure 9: Modication of the Dyck path  $rfrrrfrfff$

The red path in Figure 9 corresponds to the Dyck-path  $rfrrrfrfff$ . It can be modified to a  $B$ -path by replacing the three  $f$ 's at the end by any three letter word on  $\{r, f\}$ . There are  $2^3$  such choices, the choice  $rfr$  gives the  $B$ -path  $rfrrrfrfr$  which is displayed as a dashed path in the figure.  $\square$

To proceed we need a number of lemmas.

## 6.1 Lemmata

**Lemma 13.** For integers  $n \geq 0$  we have

$$\sum_{k=0}^n k2^{n-k} = 2^{n+1} - n - 2.$$

*Proof.* Induction on  $n$ .  $\square$

**Lemma 14.** For integers  $n \geq 0$  we have

$$\sum_{k=0}^n k^2 2^{n-k} = 6 \cdot 2^n - n^2 - 4n - 6.$$

*Proof.* Induction on  $n$ .  $\square$

**Lemma 15.** For integers  $0 \leq m \leq n$  we have

$$\sum_{s=m}^n \binom{s}{m} = \binom{n+1}{m+1}.$$

*Proof.* This follows from Pascal's rule.  $\square$

**Lemma 16.** For all positive integers  $A$  and  $B$  with  $A \geq B$ , we have

$$\sum_{\ell=0}^{A-B} 2^\ell \binom{A-\ell}{B} = \sum_{s=0}^{A-B} \binom{A+1}{s}.$$

*Proof.* We first convert the sum on the left into standard hypergeometric notation, as given by

$${}_pF_q \left[ \begin{matrix} a_1, \dots, a_p \\ b_1, \dots, b_q \end{matrix}; z \right] = \sum_{\ell=0}^{\infty} \frac{(a_1)_\ell, \dots, (a_p)_\ell}{\ell! (b_1)_\ell, \dots, (b_q)_\ell} z^\ell,$$

where the Pochhammer symbol  $(\alpha)_\ell$  is defined by

$$(\alpha)_\ell := \alpha(\alpha+1)\cdots(\alpha+\ell-1)$$

for  $\ell \geq 1$ , and  $(\alpha)_0 := 1$ .

In this notation we have

$$\sum_{\ell=0}^{A-B} 2^\ell \binom{A-\ell}{B} = \binom{A}{B} {}_2F_1 \left[ \begin{matrix} 1, -A+B \\ -A \end{matrix}; 2 \right].$$

We now apply the following transformation formula for hypergeometric functions, see Slater [10, (1.8.10)].

$${}_2F_1 \left[ \begin{matrix} a, -N \\ c \end{matrix}; z \right] = (1-z)^N \frac{(a)_N}{(c)_N} {}_2F_1 \left[ \begin{matrix} -N, c-a \\ 1-a-N \end{matrix}; \frac{1}{1-z} \right],$$

where  $N$  is a nonnegative integer.

Choosing  $a = 1 + \varepsilon$ ,  $N = A - B$ ,  $c = -A + \varepsilon$  and  $z = 2$  gives the formula

$${}_2F_1 \left[ \begin{matrix} 1, -A+B \\ -A \end{matrix}; 2 \right] = (-1)^{A-B} \lim_{\varepsilon \rightarrow 0} \frac{(1+\varepsilon)_{A-B}}{(-A+\varepsilon)_{A-B}} {}_2F_1 \left[ \begin{matrix} -A+B, -A-1 \\ -A+B-\varepsilon \end{matrix}; -1 \right].$$

Applying this formula to our previous expression we obtain

$$\begin{aligned} \sum_{\ell=0}^{A-B} 2^\ell \binom{A-\ell}{B} &= \binom{A}{B} \lim_{\varepsilon \rightarrow 0} \frac{(1+\varepsilon)_{A-B}}{(B+1-\varepsilon)_{A-B}} {}_2F_1 \left[ \begin{matrix} -A+B, -A-1 \\ -A+B-\varepsilon \end{matrix}; -1 \right] \\ &= \sum_{s=0}^{A-B} \frac{(-A-1)_s}{s!} (-1)^s \\ &= \sum_{s=0}^{A-B} \binom{A+1}{s}. \end{aligned}$$

□

We are now ready to prove Theorem 1. The proof will occupy the remainder of the paper, so we shall divide it into a number of parts.

*Layout of the proof:* The number  $\tilde{b}_n$  equals the number of admissible pairs of  $B_n$ -paths. We shall count these by counting the numbers of admissible pairs  $(p, q)$  where  $p \in \mathfrak{B}(a, b)$  and  $q \in \mathfrak{B}(i, j)$  and then taking the sum over all relevant such integer quadruples  $(a, b, i, j)$ . We partition these quadruples into four disjoint classes and we proceed by separately computing the number of admissible pairs belonging to each class. The rest of the proof is a simplification of the obtained expression.



## 6.2 Case I

Consider those  $(a, b, i, j)$  where  $a \geq 3$ ,  $4 \leq b \leq 2n$  and  $i \leq 2n + 1$ . This is the most general case. For each such  $a$  and  $b$ , we have  $|\mathfrak{B}_n(a, b)| = |\mathfrak{B}_{n-1}(b-2)|$  choices for  $p$ , and for each such choice we choose  $j$  between  $(2n+4-a)$  and  $2n+1$ , and for such  $a, b, j$  we have  $(b+j-2n-4)$  ways to choose  $i$ , and then  $|\mathfrak{B}_n(i, j)| = |\mathfrak{B}_{n-1}(j-2)|$  ways to complete the path. So for this case, the number of admissible paths can be written

$$s_1 = \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)| \sum_{a=3}^{b-1} \sum_{j=2n+4-a}^{2n+1} (b+j-2n-4) |\mathfrak{B}_{n-1}(j-2)|.$$

Here is an attempt of illustration: when  $n = 4$ , the  $p$ 's for which the number of  $(p, q)$  are counted in this case are all drawn on top of each other in the picture below.




Figure 10: Case I

## 6.3 Case II

Next consider the case when  $a \geq 3$ ,  $b = 2n + 1$  and  $i \leq 2n + 1$ , that is, the case when  $p$  only contains one  $f$ . The path  $p$  is then determined by the choice of  $a$ , and the path  $q$  can be chosen as in Case I, except that there are now less restriction on  $i$ : any number  $2 \leq i \leq j-1$  will do, so there are  $(j-2)$  choices.

$$s_2 = \sum_{a=3}^{2n} \sum_{j=2n+4-a}^{2n+1} (j-2) |\mathfrak{B}_{n-1}(j-2)|.$$

Again, an illustration for  $n = 4$ :




Figure 11: Case II

### 6.4 Case III

Next consider the case when  $a = 2$ ,  $b \leq 2n$  and  $i \leq 2n + 1$  (which corresponds to the paths  $p$  starting with  $rf$ ). When  $b$  is chosen, and the number of  $p$  is counted, the possibilities for  $q$  are very restricted. We have one choice for  $j$ , and  $(b - 3)$  choices for  $i$ , that is

$$s_3 = \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)|(b-3).$$

This is illustrated by the following picture.




Figure 12: Case III

### 6.5 Case IV

This case considers all remaining admissible pairs of paths. When  $a = 2$ ,  $b = 2n + 1$  and  $i \leq 2n + 1$ , the path  $p$  is fixed and  $q$  is determined by its parameter  $i$  for which there are  $(2n - 1)$  choices.

Everything up to now has been restricted by the condition  $i \leq 2n + 1$ . It remains to consider the case when  $q = rr \cdots rr$ . In this case we have  $\binom{2n}{n}$  possibilities for  $p$ .

Similarly, we have to consider the case  $p = rr \cdots rr$ , where we also have  $\binom{2n}{n}$  possibilities for  $q$ . Here the single case  $(p, q) = (rr \cdots rr, rr \cdots rr)$  has to be subtracted though, in order

not to count it twice.

$$s_4 = (2n - 1) + \binom{2n}{n} + \binom{2n}{n} - 1.$$

## 6.6 Simplification of the total sum

All in all, in summing up we have  $\tilde{b}_n = s_1 + s_2 + s_3 + s_4$ , or explicitly

$$\begin{aligned} \tilde{b}_n &= \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)| \sum_{a=3}^{b-1} \sum_{j=2n+4-a}^{2n+1} (b+j-2n-4) |\mathfrak{B}_{n-1}(j-2)| \\ &+ \sum_{a=3}^{2n} \sum_{j=2n+4-a}^{2n+1} (j-2) |\mathfrak{B}_{n-1}(j-2)| \\ &+ \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)|(b-3) \\ &+ 2 \binom{2n}{n} + (2n-2). \end{aligned}$$

In the first and second term (row) the variable  $a$  only determines the index of a sum, so we can eliminate this variable.

$$\begin{aligned} \tilde{b}_n &= \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)| \sum_{j=2n-b+5}^{2n+1} (b+j-2n-4)^2 |\mathfrak{B}_{n-1}(j-2)| \\ &+ \sum_{j=4}^{2n+1} (j-3)(j-2) |\mathfrak{B}_{n-1}(j-2)| \\ &+ \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)|(b-3) \\ &+ 2 \binom{2n}{n} + (2n-2). \end{aligned}$$

We next replace  $j$  by  $j-2n+b-3$  as index variable in the first expression, and the third and first terms are factored together. This gives

$$\begin{aligned} \tilde{b}_n &= \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)| \left( (b-3) + \sum_{j=2}^{b-2} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right) \\ &+ \left( \sum_{j=4}^{2n+1} (j-3)(j-2) |\mathfrak{B}_{n-1}(j-2)| \right) + 2 \binom{2n}{n} + (2n-2). \end{aligned}$$

In the sum going from  $j = 2$  to  $b - 2$  we can sum from 1 instead since the new first term will be 0. The last term is  $(b - 3)^2$ , and  $(b - 3)^2 + (b - 3) = (b - 3)(b - 2)$  so

$$\begin{aligned}\tilde{b}_n &= \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)| \left( (b-3)(b-2) + \sum_{j=1}^{b-3} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right) \\ &\quad + \left( \sum_{j=4}^{2n+1} (j-3)(j-2) |\mathfrak{B}_{n-1}(j-2)| \right) + 2 \binom{2n}{n} + (2n-2).\end{aligned}$$

Similarly we take out the last term of the last sum and write

$$\begin{aligned}\tilde{b}_n &= \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)| \left( (b-3)(b-2) + \sum_{j=1}^{b-3} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right) \\ &\quad + \left( \sum_{j=4}^{2n} (j-3)(j-2) |\mathfrak{B}_{n-1}(j-2)| \right) + (2n-2)(2n-1) + 2 \binom{2n}{n} + (2n-2).\end{aligned}$$

Now the last sum over  $j$  can be absorbed into the first term, and we can factor two terms at the end to obtain

$$\begin{aligned}\tilde{b}_n &= \sum_{b=4}^{2n} |\mathfrak{B}_{n-1}(b-2)| \left( 2(b-3)(b-2) + \sum_{j=1}^{b-3} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right) \\ &\quad + 4n(n-1) + 2 \binom{2n}{n}.\end{aligned}$$

Next we want to apply Lemma 12 to rewrite the functions  $\mathfrak{B}$  in a more explicit form. Since  $\mathfrak{B}$  is a piecewise defined function, we first split up our sums in the corresponding way.

$$\begin{aligned}\tilde{b}_n &= \sum_{b=4}^{n+1} |\mathfrak{B}_{n-1}(b-2)| \left( 2(b-3)(b-2) + \sum_{j=1}^{b-3} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right) \\ &\quad + \sum_{b=n+2}^{2n} |\mathfrak{B}_{n-1}(b-2)| \left( 2(b-3)(b-2) + \sum_{j=1}^{b-3} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right) \\ &\quad + 4n(n-1) + 2 \binom{2n}{n} \\ &= \sum_{b=4}^{n+1} |\mathfrak{B}_{n-1}(b-2)| \left( 2(b-3)(b-2) + \sum_{j=1}^{b-3} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right)\end{aligned}$$

$$\begin{aligned}
& + \sum_{b=n+2}^{2n} |\mathfrak{B}_{n-1}(b-2)| \left( 2(b-3)(b-2) + \left( \sum_{j=1}^{b-n-2} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right. \right. \\
& \quad \left. \left. + \sum_{j=b-n-1}^{b-3} (j-1)^2 |\mathfrak{B}_{n-1}(j+2n-b+1)| \right) \right) \\
& + 4n(n-1) + 2 \binom{2n}{n}.
\end{aligned}$$

Now the sums are partitioned properly and we can apply Lemma 12. To avoid using a lot of binomial coefficients we temporarily introduce the notation

$$D(n, i, j) := \binom{2n-i-j-2}{n-i-1} - \binom{2n-i-j-2}{n-i-j-1}.$$

Lemma 12 now gives

$$\begin{aligned}
\tilde{b}_n &= \sum_{b=4}^{n+1} \sum_{k=1}^{n-2} 2^k D(n-1, b-3, k) \left( 2(b-3)(b-2) + \sum_{j=1}^{b-3} (j-1)^2 2^{b-j-3} \right) \\
& + \sum_{b=n+2}^{2n} 2^{2n-b} \left( 2(b-3)(b-2) + \left( \sum_{j=1}^{b-n-2} (j-1)^2 \sum_{k=1}^{n-2} 2^k D(n-1, 2n+j-b, k) \right. \right. \\
& \quad \left. \left. + \sum_{j=b-n-1}^{b-3} (j-1)^2 2^{b-j-3} \right) \right) \\
& + 4n(n-1) + 2 \binom{2n}{n}.
\end{aligned}$$

Changing summation variables such that both sums over  $b$  goes up to  $n-2$  we obtain

$$\begin{aligned}
\tilde{b}_n &= \sum_{b=1}^{n-2} \sum_{k=1}^{n-2} 2^k D(n-1, b, k) \left( 2b(b+1) + \sum_{j=1}^b (j-1)^2 2^{b-j} \right) \\
& + \sum_{b=0}^{n-2} 2^{n-b-2} \left( 2(b+n-1)(b+n) + \left( \sum_{j=1}^b (j-1)^2 \sum_{k=1}^{n-2} 2^k D(n-1, n+j-b-2, k) \right. \right. \\
& \quad \left. \left. + \sum_{j=b+1}^{b+n-1} (j-1)^2 2^{b+n-j-1} \right) \right) \\
& + 4n(n-1) + 2 \binom{2n}{n}.
\end{aligned}$$

Using Lemma 14 (and changing variables) we simplify the following two expressions:

$$\sum_{j=1}^b (j-1)^2 2^{b-j} = 3 \cdot 2^b - b^2 - 2b - 3$$

$$\sum_{j=b+1}^{b+n-1} (j-1)^2 2^{b+n-j-1} = 2^{n-1}(b^2 + 2b + 3) - n^2 - b^2 - 2bn - 2.$$

Inserting this into our expression for  $\tilde{b}_n$  we have

$$\begin{aligned} \tilde{b}_n &= \sum_{b=1}^{n-2} \sum_{k=1}^{n-2} 2^k D(n-1, b, k) (3 \cdot 2^b + b^2 - 3) \\ &+ \sum_{b=0}^{n-2} 2^{n-b-2} \left( 2(b+n-1)(b+n) + \left( \sum_{j=1}^b (j-1)^2 \sum_{k=1}^{n-2} 2^k D(n-1, n+j-b-2, k) \right. \right. \\ &\quad \left. \left. + 2^{n-1}(b^2 + 2b + 3) - n^2 - b^2 - 2bn - 2 \right) \right) \\ &+ 4n(n-1) + 2 \binom{2n}{n}. \end{aligned}$$

This can be rewritten as

$$\begin{aligned} \tilde{b}_n &= \sum_{b=1}^{n-2} \sum_{k=1}^{n-2} 2^k D(n-1, b, k) (3 \cdot 2^b + b^2 - 3) \\ &+ \sum_{b=0}^{n-2} 2^{n-b-2} \left( \sum_{j=1}^b (j-1)^2 \sum_{k=1}^{n-2} 2^k D(n-1, n+j-b-2, k) \right) \\ &+ \sum_{b=0}^{n-2} (2^{2n-b-3}(b^2 + 2b + 3) + 2^{n-b-2}(n^2 + b^2 + 2bn - 2b - 2n - 2)) \\ &+ 4n(n-1) + 2 \binom{2n}{n}. \end{aligned}$$

Here the last sum over  $b$  equals  $2^{2n+1} - 2^n(n+3) - 4n(n-1)$  which is easily proved by application of Lemma 13 and Lemma 14. Canceling the term  $4n(n-1)$  we have

$$\begin{aligned}
\tilde{b}_n &= \sum_{b=1}^{n-2} \sum_{k=1}^{n-2} 2^k D(n-1, b, k) (3 \cdot 2^b + b^2 - 3) \\
&+ \sum_{b=0}^{n-2} 2^{n-b-2} \left( \sum_{j=1}^b (j-1)^2 \sum_{k=1}^{n-2} 2^k D(n-1, n+j-b-2, k) \right) \\
&+ 2^{2n+1} - 2^n(n+3) + 2 \binom{2n}{n}.
\end{aligned}$$

Expanding our function  $D$  we more explicitly have

$$\begin{aligned}
\tilde{b}_n &= \sum_{b=1}^{n-2} \sum_{k=1}^{n-2} 2^k \left( \binom{2n-b-k-4}{n-b-2} - \binom{2n-b-k-4}{n-b-k-2} \right) (3 \cdot 2^b + b^2 - 3) \\
&+ \sum_{b=0}^{n-2} 2^{n-b-2} \left( \sum_{j=1}^b (j-1)^2 \sum_{k=1}^{n-2} 2^k \left( \binom{n+b-j-k-2}{b-j} - \binom{n+b-j-k-2}{b-j-k} \right) \right) \\
&+ 2^{2n+1} - 2^n(n+3) + 2 \binom{2n}{n}.
\end{aligned}$$

Changing order of summation to merge the two sums over  $k$ , and rewriting some binomial coefficients we obtain

$$\begin{aligned}
\tilde{b}_n &= \sum_{k=1}^{n-2} 2^k \sum_{b=1}^{n-2} \left( (3 \cdot 2^b + b^2 - 3) \left( \binom{2n-b-k-4}{n-b-2} - \binom{2n-b-k-4}{n-2} \right) \right. \\
&+ 2^{n-b-2} \sum_{j=1}^b (j-1)^2 \left( \binom{n+b-j-k-2}{n-k-2} - \binom{n+b-j-k-2}{n-2} \right) \left. \right) \\
&- 2^n(n+3) + 2^{2n+1} + 2 \binom{2n}{n}. \tag{1}
\end{aligned}$$

## 6.7 Further simplification

The remaining simplification of the above expression, including the proof of Lemma 16 above, is due to Christian Krattenthaler.

We continue by simplifying the sum over  $j$  in (1). For this purpose we write

$$(j-1)^2 = (n+b-j-k)(n+b-j-k-1) - (n+b-j-k-1)(2n+2b-2k-3) + (n+b-k-2)^2$$

and use this to simplify the binomial coefficients. Thus the sum over  $j$  becomes

$$\begin{aligned}
& (n-k)(n-k-1) \sum_{j=1}^b \binom{n+b-j-k}{n-k} - n(n-1) \sum_{j=1}^b \binom{n+b-j-k}{n} \\
& - (2n+2b-2k-3) \left( (n-k-1) \sum_{j=1}^b \binom{n+b-j-k-1}{n-k-1} \right. \\
& \quad \left. - (n-1) \sum_{j=1}^b \binom{n+b-j-k-1}{n-1} \right) \\
& + (n+b-k-2)^2 \left( \sum_{j=1}^b \binom{n+b-j-k-2}{n-k-2} \right. \\
& \quad \left. - \sum_{j=1}^b \binom{n+b-j-k-2}{n-2} \right).
\end{aligned}$$

Changing variables in each sum and then removing zero-terms we obtain

$$\begin{aligned}
& (n-k)(n-k-1) \sum_{j=n-k}^{n+b-k-1} \binom{j}{n-k} - n(n-1) \sum_{j=n-k}^{n+b-k-1} \binom{j}{n} \\
& - (2n+2b-2k-3) \left( (n-k-1) \sum_{j=n-k-1}^{n+b-k-2} \binom{j}{n-k-1} - (n-1) \sum_{j=n-k-1}^{n+b-k-2} \binom{j}{n-1} \right) \\
& + (n+b-k-2)^2 \left( \sum_{j=n-k-2}^{n+b-k-3} \binom{j}{n-k-2} - \sum_{j=n-k-2}^{n+b-k-3} \binom{j}{n-2} \right) \\
& = (n-k)(n-k-1) \sum_{j=n-k}^{n+b-k-1} \binom{j}{n-k} - n(n-1) \sum_{j=n}^{n+b-k-1} \binom{j}{n} \\
& - (2n+2b-2k-3) \left( (n-k-1) \sum_{j=n-k-1}^{n+b-k-2} \binom{j}{n-k-1} - (n-1) \sum_{j=n-1}^{n+b-k-2} \binom{j}{n-1} \right) \\
& + (n+b-k-2)^2 \left( \sum_{j=n-k-2}^{n+b-k-3} \binom{j}{n-k-2} - \sum_{j=n-2}^{n+b-k-3} \binom{j}{n-2} \right).
\end{aligned}$$

Here we apply Lemma 15 six times. This gives



$$\begin{aligned}
& (n-k)(n-k-1) \binom{n+b-k}{n-k+1} - n(n-1) \binom{n+b-k}{n+1} \\
& - (2n+2b-2k-3) \left( (n-k-1) \binom{n+b-k-1}{n-k} - (n-1) \binom{n+b-k-1}{n} \right) \\
& + (n+b-k-2)^2 \left( \binom{n+b-k-2}{n-k-1} - \binom{n+b-k-2}{n-1} \right).
\end{aligned}$$

This simplifies to

$$\binom{n+b-k-1}{n-k+1} + \binom{n+b-k-2}{n-k+1} - \binom{n+b-k-1}{n+1} - \binom{n+b-k-2}{n+1}. \quad (2)$$

So inserting (2) back into (1) we now have

$$\begin{aligned}
\tilde{b}_n &= \sum_{k=1}^{n-2} 2^k \sum_{b=1}^{n-2} \left( (3 \cdot 2^b + b^2 - 3) \left( \binom{2n-b-k-4}{n-b-2} - \binom{2n-b-k-4}{n-2} \right) \right. \\
& \quad \left. + 2^{n-b-2} \left( \binom{n+b-k-1}{n-k+1} + \binom{n+b-k-2}{n-k+1} \right) \right. \\
& \quad \left. - \binom{n+b-k-1}{n+1} - \binom{n+b-k-2}{n+1} \right) \\
& - 2^n(n+3) + 2^{2n+1} + 2 \binom{2n}{n}. \quad (3)
\end{aligned}$$

On the second line we reverse the order of the  $b$ -summation, that is, we replace  $b$  by  $n-1-b$  which yields

$$\begin{aligned}
\tilde{b}_n &= \sum_{k=1}^{n-2} 2^k \sum_{b=1}^{n-2} \left( (3 \cdot 2^b + b^2 - 3) \left( \binom{2n-b-k-4}{n-b-2} - \binom{2n-b-k-4}{n-2} \right) \right. \\
& \quad \left. + 2^{b-1} \left( \binom{2n-b-k-2}{n-k+1} + \binom{2n-b-k-3}{n-k+1} \right) \right. \\
& \quad \left. - \binom{2n-b-k-2}{n+1} - \binom{2n-b-k-3}{n+1} \right) \\
& - 2^n(n+3) + 2^{2n+1} + 2 \binom{2n}{n}.
\end{aligned}$$

Introducing a new summation variable  $\ell = b + k$  we can write this as

$$\begin{aligned}
\tilde{b}_n &= \sum_{k=1}^{n-2} 2^k \sum_{b=1}^{n-2} (b^2 - 3) \left( \binom{2n-b-k-4}{n-k-2} - \binom{2n-b-k-4}{n-2} \right) \\
&+ \sum_{k=1}^{n-2} \sum_{\ell=k+1}^{2n-2} 2^\ell \left( 3 \binom{2n-\ell-4}{n-k-2} + 3 \binom{2n-\ell-4}{n-2} \right) \\
&+ \frac{1}{2} \left( \binom{2n-\ell-2}{n-k+1} + \binom{2n-\ell-3}{n-k+1} - \binom{2n-\ell-2}{n+1} - \binom{2n-\ell-3}{n+1} \right) \\
&- 2^n(n+3) + 2^{2n+1} + 2 \binom{2n}{n}.
\end{aligned}$$

Next we apply Lemma 16 to all the sums over  $\ell$ . This gives

$$\begin{aligned}
\tilde{b}_n &= \sum_{k=1}^{n-2} 2^k \sum_{b=1}^{n-2} (b^2 - 3) \left( \binom{2n-b-k-4}{n-k-2} - \binom{2n-b-k-4}{n-2} \right) \\
&+ 3 \sum_{k=1}^{n-2} 2^{k+1} \sum_{s=0}^{n-3} \binom{2n-k-4}{s} - 3 \sum_{k=1}^{n-2} 2^{k+1} \sum_{s=0}^{n-k-3} \binom{2n-k-4}{s} \\
&+ \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n-4} \binom{2n-k-2}{s} + \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n-5} \binom{2n-k-3}{s} \\
&- \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n-k-4} \binom{2n-k-2}{s} - \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n-k-5} \binom{2n-k-3}{s} \\
&- 2^n(n+3) + 2^{2n+1} + 2 \binom{2n}{n}. \tag{4}
\end{aligned}$$

To simplify (4) we first note that

$$\begin{aligned}
(1+1)^{2n-k-2} &= \sum_{s=0}^{2n-k-2} \binom{2n-k-2}{s} = \sum_{s=0}^{n+1} \binom{2n-k-2}{s} + \sum_{s=n+2}^{2n-k-2} \binom{2n-k-2}{s} \\
&= \sum_{s=0}^{n+1} \binom{2n-k-2}{s} + \sum_{s=0}^{n-k-4} \binom{2n-k-2}{n-k-4-s} \\
&= \sum_{s=0}^{n+1} \binom{2n-k-2}{s} + \sum_{s=0}^{n-k-4} \binom{2n-k-2}{s}.
\end{aligned}$$

Hence we have

$$\sum_{s=0}^{n-k-4} \binom{2n-k-2}{s} = 2^{2n-k-2} - \sum_{s=0}^{n+1} \binom{2n-k-2}{s}.$$

Using this, and similar identities, we transform the expression (4) to

$$\begin{aligned} \tilde{b}_n &= \sum_{k=1}^{n-2} 2^k \sum_{b=1}^{n-2} (b^2 - 3) \left( \binom{2n-b-k-4}{n-k-2} - \binom{2n-b-k-4}{n-2} \right) \\ &+ 3 \sum_{k=1}^{n-2} 2^{k+1} \sum_{s=0}^{n-3} \binom{2n-k-4}{s} - 3(n-2)2^{2n-3} + 3 \sum_{k=1}^{n-2} 2^{k+1} \sum_{s=0}^{n-2} \binom{2n-k-4}{s} \\ &+ \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n-4} \binom{2n-k-2}{s} + \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n-5} \binom{2n-k-3}{s} \\ &- (n-2)2^{2n-2} + \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n+1} \binom{2n-k-2}{s} - (n-2)2^{2n-3} + \sum_{k=1}^{n-2} 2^k \sum_{s=0}^{n+1} \binom{2n-k-3}{s} \\ &- 2^n(n+3) + 2^{2n+1} + 2 \binom{2n}{n}. \end{aligned} \tag{5}$$

In the second, third and fourth line we can interchange summations over  $k$  and  $s$  and apply Lemma 16 again. However, one has to be a bit careful since the sums over  $k$  are in fact shorter than required by the lemma. For example, the first of these sums can be simplified as follows.

$$\begin{aligned} 3 \sum_{k=1}^{n-2} 2^{k+1} \sum_{s=0}^{n-3} \binom{2n-k-4}{s} &= 3 \sum_{s=0}^{n-3} \sum_{k=1}^{n-2} 2^{k+1} \binom{2n-k-4}{s} \\ &= 3 \sum_{s=0}^{n-3} \left( \sum_{k=1}^{2n-s-4} 2^{k+1} \binom{2n-k-4}{s} - \sum_{k=n-1}^{2n-s-4} 2^{k+1} \binom{2n-k-4}{s} \right) \\ &= 3 \sum_{s=0}^{n-3} \left( 4 \sum_{k=0}^{2n-s-5} \binom{2n-4}{k} - 2^n \sum_{k=0}^{n-s-3} \binom{n-2}{k} \right) \\ &= 12 \sum_{k=0}^{n-3} (n-2) \binom{2n-4}{k} + 12 \sum_{k=0}^{n-3} (n-k-2) \binom{2n-4}{n+k-2} \\ &\quad - 3 \cdot 2^n \sum_{k=0}^{n-3} (n-k-2) \binom{n-2}{k} \\ &= 6(n-2) \left( 2^{2n-4} - \binom{2n-4}{n-2} \right) + 6(n-2)2^{2n-4} - 3 \cdot 2^n (n-2)2^{n-3} \\ &= 3(n-2)2^{2n-3} - 6(n-2) \binom{2n-4}{n-2}. \end{aligned}$$

We treat the other double sums over  $k$  and  $s$  in (5) similarly, and after many, many calculations we arrive at the expression

$$\begin{aligned} \tilde{b}_n &= \sum_{k=1}^{n-2} 2^k \sum_{b=1}^{n-2} (b^2 - 3) \left( \binom{2n-b-k-4}{n-k-2} - \binom{2n-b-k-4}{n-2} \right) \\ &\quad + (3n+5)2^{2n-2} + (n^2-1)2^{n-1} - 4(1-7n+7n^2+3n^3) \frac{(2n-3)!}{(n-2)!(n+1)!}. \end{aligned} \quad (6)$$

Now we turn to the sum over  $b$ . By expressing

$$b^2 = (2n-b-k-2)(2n-b-k-3) - (4n-2k-5)(2n-b-k-3) + (2n-k-3)^2$$

and proceeding by application of Lemma 15 as in the beginning of this section, and then rewriting the binomial coefficients we obtain

$$\begin{aligned} &\sum_{b=1}^{n-2} (b^2 - 3) \left( \binom{2n-b-k-4}{n-k-2} - \binom{2n-b-k-4}{n-2} \right) \\ &= (n-k)(n-k-1) \binom{2n-k-2}{n-k+1} - n(n-1) \binom{2n-k-2}{n+1} \\ &\quad - (4n-2k-5)(n-k-1) \binom{2n-k-3}{n-k} + (4n-2k-5)(n-1) \binom{2n-k-3}{n} \\ &\quad + ((2n-k-3)^2 - 3) \binom{2n-k-4}{n-k-1} - ((2n-k-3)^2 - 3) \binom{2n-k-4}{n-1} \\ &= -2n^2 \binom{2n-k-2}{n+1} + (2n^2-1) \binom{2n-k-3}{n} - 3 \binom{2n-k-4}{n-3} + 3 \binom{2n-k-4}{n-1} \\ &\quad + (2n-5) \binom{2n-k-3}{n-3} - 2(n-2)^2 \binom{2n-k-3}{n-2} \\ &\quad - 2(n-1)(n-2) \binom{2n-k-3}{n-1} + 2(n+1)(n-1) \binom{2n-k-3}{n+1} \\ &\quad + 2 \binom{2n-k-2}{n-3} - 2(n-2) \binom{2n-k-2}{n-2} + 2(n-2)(n-1) \binom{2n-k-2}{n-1}. \end{aligned}$$

Inserting this into (6) we can apply Lemma 16 again to all the sums over  $k$ . After considerable simplification we arrive at

$$\tilde{b}_n = (3n+5)2^{2n-2} - 2(3n-1) \binom{2n-2}{n-1}.$$

This concludes the proof.

## 7 The resulting sequence

The above formula yields the integer sequence  $\{\tilde{b}_n\}_{n \geq 2}$ , which could not be found in OEIS [12]. Its first entries are

$n$	2	3	4	5	6	7	8	9	10
$\tilde{b}_n$	24	128	648	3160	14984	69536	317264	1427912	6355080.

## 8 Acknowledgements

I am particularly grateful to Professor Christian Krattenthaler, for his help with simplification of the summation formula which led to the short expression for the sequence in Theorem 1. I also appreciate the advice of Professor Volodymyr Mazorchuk and the helpful comments from the referee.

## References

- [1] C. A. Athanasiadis, Characteristic polynomials of subspace arrangements and finite fields, *Adv. Math.* **122** (1996), 193–233.
- [2] C. A. Athanasiadis, Deformations of Coxeter hyperplane arrangements and their characteristic polynomials, *Adv. Stud. Pure Math.* **27** (2000), 1–26.
- [3] K. Baur and V. Mazorchuk, Combinatorial analogues of ad-nilpotent ideals for untwisted affine Lie algebras, preprint <http://arxiv.org/abs/1108.3659>.
- [4] N. Bourbaki, *Lie Groups and Lie Algebras Chapters 4–6*, Springer, 2002.
- [5] P. Cellini and P. Papi, ad-Nilpotent ideals of a Borel subalgebra. *J. Algebra* **225** (2000), 130–141.
- [6] P. Cellini and P. Papi, ad-Nilpotent ideals of a Borel subalgebra II, *J. Algebra* **258** (2002), 112–121.
- [7] R. Grimaldi, *Discrete and Combinatorial Mathematics: an Applied Introduction*, Addison-Wesley, 2003.
- [8] C. Krattenthaler, L. Orsina, and P. Papi, Enumeration of ad-nilpotent  $\mathfrak{b}$ -ideals for simple Lie algebras, *Adv. in Appl. Math.* **28** (2002), 478–522.
- [9] J. Shi, The number of  $\oplus$ -sign types, *Quart. J. Math. Oxford* **48** (1997), 93–105.
- [10] L. J. Slater, *Generalized Hypergeometric Functions*, Cambridge University Press, 1966.
- [11] R. P. Stanley, *Enumerative Combinatorics: Volume 2*, Cambridge University Press, 1999.
- [12] N. Sloane, The on-line encyclopedia of integer sequences, <http://oeis.org>.

---

2010 *Mathematics Subject Classification*: Primary 17B05; Secondary 05E10.

*Keywords*: Borel subalgebra, ideal, Dyck path, Catalan number, poset.

---

Received April 17 2012; revised versions received August 15 2012; November 17 2012. Published in *Journal of Integer Sequences*, November 20 2012.

---

Return to [Journal of Integer Sequences home page](#).