

Solitons of the spacelike mean curvature flow in generalized Robertson-Walker spacetimes

Márcio Batista, Giovanni Molica Bisci,
Henrique F. de Lima and Wallace F. Gomes

ABSTRACT. Our purpose in this paper is to study solitons of the spacelike mean curvature flow in a generalized Robertson-Walker (GRW) spacetime $-I \times_f M^n$. Under suitable constraints on the warping function f and on the curvatures of the Riemannian fiber M^n , we apply suitable maximum principles in order to obtain nonexistence and uniqueness results concerning these solitons. Applications to standard models of GRW spacetimes, namely, the Einstein-de Sitter spacetime, steady state type spacetimes, de Sitter and anti-de Sitter spaces, are given. Furthermore, we establish new Calabi-Bernstein type results related to entire spacelike mean curvature flow graphs constructed over the Riemannian fiber of the ambient spacetime.

CONTENTS

| | |
|---|-----|
| 1. Introduction | 555 |
| 2. Spacelike hypersurfaces | 556 |
| 3. Spacelike mean curvature flow solitons | 558 |
| 4. Nonexistence of spacelike mean curvature flow solitons | 560 |
| 5. Uniqueness and further nonexistence results | 566 |
| 6. Entire spacelike mean curvature flow graphs | 571 |
| Acknowledgements | 576 |
| References | 576 |

Received November 12, 2021.

2010 *Mathematics Subject Classification.* 53C42, 53E10.

Key words and phrases. Generalized Robertson-Walker spacetime; complete spacelike mean curvature flow solitons; Einstein-de Sitter spacetime; steady state type spacetimes; de Sitter and anti-de Sitter spaces; Calabi-Bernstein type results.

The first author is partially supported by CNPq [Grant: 308440/2021-8 and 405468/2021-0], and FAPEAL [Grant: E:60030.0000001758/2022], and the third author is partially supported by CNPq [Grant: 301970/2019-0]. The second author is partially supported by INdAM-GNAMPA Research Project 2020 titled *Equazioni alle derivate parziali: problemi e modelli*. The fourth author is partially supported by CAPES [Finance Code 001], Brazil.

1. Introduction

Let \mathbb{R}_1^{n+1} be the $(n+1)$ -dimensional Minkowski space $(\mathbb{R}_1^{n+1}, \bar{g})$ with its standard Lorentzian metric

$$\bar{g} = -dx_1^2 + \sum_{i=2}^{n+1} dx_i^2.$$

Let $\psi : \Sigma^n \rightarrow \mathbb{R}_1^{n+1}$ be a spacelike immersion (which means that it has a Riemannian induced metric) in the Minkowski space. The *spacelike mean curvature flow* associated to ψ is a family of smooth spacelike immersions $\Psi_t = \Psi(t, \cdot) : \Sigma^n \rightarrow \mathbb{R}_1^{n+1}$ with corresponding images $\Sigma_t^n = \Psi_t(\Sigma^n)$ satisfying the following evolution equation

$$\begin{cases} \frac{\partial \Psi}{\partial t} = \vec{H} \\ \Psi(0, x) = \psi(x) \end{cases}$$

on some time interval, where \vec{H} stands for the (non-normalized) mean curvature vector of the spacelike submanifold Σ_t^n in \mathbb{R}_1^{n+1} .

Mean curvature flow in the Minkowski space and, more generally, in a Lorentzian manifold has been extensively studied by several authors (see, for instance, [1, 18, 19, 20, 21, 22, 23, 28, 29, 31, 32, 34, 40]) and, according to [19], an important justification for this interest is the fact that spacelike translating solitons can be regarded as a natural way of foliating spacetimes by almost null like hypersurfaces. Particular examples may give insight into the structure of certain spacetimes at null infinity and have possible applications in General Relativity (for more details, see [19]).

More recently, Lambert and Lotay [33] proved long-time existence and convergence results for spacelike solutions to mean curvature flow in the n -dimensional pseudo-Euclidean space \mathbb{R}_m^n of index m , which are entire or defined on bounded domains and satisfying Neumann or Dirichlet boundary conditions. In [24], Guilfoyle and Klingenberg proved the longtime existence for mean curvature flow of a smooth n -dimensional spacelike submanifold of an $(n + m)$ -dimensional manifold whose metric satisfies the so-called timelike curvature condition. Meanwhile, Alías, de Lira and Rigoli [10] introduced the general definition of self-similar mean curvature flow in a Riemannian manifold \bar{M}^{n+1} endowed with a vector field K and establishing the corresponding notion of mean curvature flow soliton. In particular, when \bar{M}^{n+1} is a Riemannian warped product of the type $I \times_f M^n$ and $K = f(t)\partial_t$, they applied weak maximum principles to guarantee that a complete n -dimensional mean curvature flow soliton is a slice of \bar{M}^{n+1} . In [17], Colombo, Mari and Rigoli also studied some properties of mean curvature flow solitons in general Riemannian manifolds and in warped products, focusing on splitting and rigidity results under various geometric conditions, ranging from the stability of the soliton to the fact that the image of its

Gauss map be contained in suitable regions of the sphere. Moreover, they also investigated the case of entire mean curvature flow graphs. When the ambient space is a Lorentzian product space, the first and third authors [14] established nonexistence results for complete spacelike translating solitons under suitable curvature constraints on the curvatures of the Riemannian base of the ambient space. In particular, they obtained Calabi-Bernstein type results for entire translating graphs constructed over this Riemannian base. For this, they proved a version of the Omori-Yau's maximum principle for complete spacelike translating solitons. Besides, they also constructed new examples of rotationally symmetric spacelike translating solitons embedded in such an ambient space.

Proceeding with this picture, here we extend the techniques developed in [6, 10, 13, 14, 17] to study complete spacelike mean curvature flow solitons immersed in a generalized Robertson-Walker (GRW) spacetime, that is, a Lorentzian warped product $-I \times_f M^n$ with 1-dimensional negative definite base I and n -dimensional Riemannian fiber M^n . Under suitable constraints on the warping function f and on the curvatures of M^n , we apply suitable maximum principles in order to obtain nonexistence and uniqueness results concerning these solitons. Applications to standard GRW spacetimes as, for instance, the Einstein-de Sitter and steady state type spacetimes, are given. Furthermore, we establish new Calabi-Bernstein type results related to entire spacelike mean curvature flow graphs constructed over the Riemannian fiber of the ambient spacetime.

2. Spacelike hypersurfaces

Let (M^n, g_M) be a connected, n -dimensional, oriented Riemannian manifold, $I \subset \mathbb{R}$ an open interval and $f : I \rightarrow \mathbb{R}$ a positive smooth function. Also, in the product manifold $\overline{M}^{n+1} = I \times M^n$ let π_I and π_M denote the canonical projections onto the factors I and M^n , respectively.

The class of Lorentzian manifolds which will be of our concern here is the one obtained by furnishing \overline{M}^{n+1} with the Lorentzian metric \overline{g} given by

$$\overline{g}(u, v)_p = -g_I((\pi_I)_*u, (\pi_I)_*v)_{\pi_I(p)} + (f \circ \pi_I)^2(p)g_M((\pi_M)_*u, (\pi_M)_*v)_{\pi_M(p)},$$

for all $p \in \overline{M}^{n+1}$ and $u, v \in T_p\overline{M}$, where g_I stands for the standard metric of $I \subset \mathbb{R}$. Along this work, we will simply write

$$\overline{M}^{n+1} = -I \times_f M^n. \quad (1)$$

According to the nomenclature established in [12], we say that \overline{M}^{n+1} is a *generalized Robertson Walker* (GRW) spacetime with warping function f and Riemannian fiber M^n . When M^n has constant sectional curvature, (1) has been known in the mathematical literature as a Robertson-Walker (RW) spacetime, an allusion to the fact that, for $n = 3$, it is an exact solution of Einstein's field equations (see, for instance, [37, Chapter 12]).

In this setting, we will consider the timelike conformal closed vector field $\mathcal{K} = f(\pi_I)\partial_t$ globally defined on \bar{M} , where $\partial_t = \frac{\partial}{\partial t}$ stands for the coordinate timelike vector field tangent to I . From the relationship between the Levi-Civita connections of \bar{M} and those ones of I and M^n (see [37, Proposition 7.35]), it follows that

$$\bar{\nabla}_V \mathcal{K} = f'(\pi_I)V, \tag{2}$$

for all $V \in \mathfrak{X}(\bar{M})$, where $\bar{\nabla}$ is the Levi-Civita connection of \bar{g} .

Let Σ^n be an n -dimensional connected manifold. A smooth immersion $\psi : \Sigma^n \rightarrow \bar{M}^{\overline{n+1}}$ is said to be a *spacelike hypersurface* if Σ^n , furnished with the metric g induced from \bar{g} via ψ , is a Riemannian manifold. We will denote by ∇ the Levi-Civita connection of g . Since \bar{M} is time-orientable, it follows from the connectedness of Σ^n that one can uniquely choose a globally defined timelike unit vector field $N \in \mathfrak{X}^\perp(\Sigma)$, having the same time-orientation of ∂_t , that is, such that $\bar{g}(N, \partial_t) < 0$. In this case, one says that N is the *future-pointing Gauss map* of Σ^n and will always assume such a timelike orientation for Σ^n . From the inverse Cauchy-Schwarz inequality (see [37, Proposition 5.30]), we have that $\bar{g}(N, \partial_t) \leq -1$, with the equality holding at a point $p \in \Sigma^n$ if, and only if, $N = \partial_t$ at p .

We will denote by A and $H = -\text{trace}(A)$ the shape operator and the (non-normalized) mean curvature function of the spacelike hypersurface $\psi : \Sigma^n \rightarrow \bar{M}^{\overline{n+1}}$ with respect to its future-pointing Gauss map N . Throughout this paper, the mean curvature H taken with respect to such a choice of orientation N will be called the *future mean curvature* of Σ^n . In particular, for a fixed $t_* \in I$, from [7, Example 5.6] we have that the *slice* $\{t_*\} \times M^n$ has constant future mean curvature $H = n \frac{f'(t_*)}{f(t_*)}$ with respect to $N = \partial_t$.

Now, we consider two particular functions naturally attached to a spacelike hypersurface Σ^n immersed into a GRW spacetime $\bar{M}^{\overline{n+1}} = -I \times_f M^n$, namely, the height function $h = (\pi_I)|_\Sigma$ and the hyperbolic angle function $\Theta = \bar{g}(N, \partial_t)$, where we recall that N denotes the future-pointing Gauss map of Σ^n and, consequently, $\Theta \leq -1$. A simple computation shows that

$$\bar{\nabla} \pi_I = -\bar{g}(\bar{\nabla} \pi_I, \partial_t) \partial_t = -\partial_t. \tag{3}$$

So, from (3) we have

$$\nabla h = (\bar{\nabla} \pi_I)^\top = -\partial_t^\top = -\partial_t - \Theta N. \tag{4}$$

Thus, (4) gives the following relation

$$|\nabla h|^2 = \Theta^2 - 1, \tag{5}$$

where $|\cdot|$ stands for the norm of a tangent vector field on Σ^n in the metric g .

On the other hand, from (2) we have that

$$\bar{\nabla}_V \partial_t = \frac{f'(\pi_I)}{f(\pi_I)} \{V + \bar{g}(V, \partial_t) \partial_t\}, \tag{6}$$

for any tangent vector V on \overline{M}^{n+1} .

Hence, from (4) and (6) we deduce that, for any $X \in \mathfrak{X}(\Sigma)$, the Hessian of h in the metric g is given by

$$\begin{aligned} \nabla^2 h(X, X) &= g(\nabla_X \nabla h, X) \\ &= -\frac{f'(h)}{f(h)}\{|X|^2 + g(X, \nabla h)^2\} + g(AX, X)\Theta. \end{aligned} \quad (7)$$

Hence, from (7) we obtain that the Laplacian of h in the metric g is

$$\Delta h = -\frac{f'(h)}{f(h)}\{n + |\nabla h|^2\} - H\Theta. \quad (8)$$

3. Spacelike mean curvature flow solitons

We recall that the spacelike mean curvature flow $\Psi : [0, T) \times \Sigma^n \rightarrow \overline{M}^{n+1}$ of a spacelike hypersurface $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ in a $(n + 1)$ -dimensional Lorentzian manifold \overline{M}^{n+1} , satisfying $\Psi(0, \cdot) = \psi(\cdot)$, looks for solutions of the equation

$$\frac{\partial \Psi}{\partial t} = \vec{H},$$

where $\vec{H}(t, \cdot)$ is the (non-normalized) mean curvature vector of $\Sigma_t^n = \Psi(t, \Sigma^n)$ (see, for instance, [33]). In our context, according to [10, Definition 1.1] and [17, Definition 1.1], a spacelike hypersurface $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ immersed in a GRW spacetime $\overline{M}^{n+1} = -I \times_f M^n$ is said a *spacelike mean curvature flow soliton* with respect to $\mathcal{K} = f(t)\partial_t$ and with *soliton constant* $c \in \mathbb{R}$ if its (non-normalized) future mean curvature function satisfies

$$H = cf(h)\Theta. \quad (9)$$

In fact, considering that Ψ is a self-similar mean curvature flow with respect to some vector field X , we can reason as in [10, Proposition 2.1] to deduce that the corresponding mean curvature vector satisfies

$$\vec{H} = cX^\perp,$$

for some constant $c \in \mathbb{R}$. In our setting, X is equal to $\mathcal{K} = f(t)\partial_t$ and, hence, assuming that $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ satisfies equation (9) means that it is a solution of the mean curvature flow evolution equation.

Adopting the terminology introduced in [10] and [17], we will also consider the *soliton function*

$$\zeta_c(t) = nf'(t) + cf^2(t). \quad (10)$$

So, each slice $M_{t_*} = \{t_*\} \times M^n$ is a spacelike mean curvature flow soliton with respect to $\mathcal{K} = f(t)\partial_t$ and with soliton constant c given by

$$c = -n \frac{f'(t_*)}{f(t_*)^2}. \quad (11)$$

Moreover, t_* is implicitly given by the condition $\zeta_c(t_*) = 0$.

We close this section quoting important examples which will be addressed along the next sections.

Example 3.1. The 4-dimensional Einstein-de Sitter spacetime is modeled by the GRW spacetime $-\mathbb{R}^+ \times_{t^{\frac{2}{3}}} \mathbb{R}^3$, where \mathbb{R}^3 stands for the 3-dimensional Euclidean space endowed with its canonical metric. This spacetime is a classical exact solution to the Einstein field equation without cosmological constant. It is an open Friedmann-Robertson-Walker model, which incorporates homogeneity and isotropy (the cosmological principle) and permitted expansion (for more details, see [37, Chapter 12]). Here, we consider the $(n + 1)$ -dimensional Einstein-de Sitter spacetime $-\mathbb{R}^+ \times_{t^{\frac{2}{3}}} \mathbb{R}^n$. From (11) we conclude that the slice

$$\left\{ \left(-\frac{2n}{3c}\right)^{\frac{3}{5}} \right\} \times \mathbb{R}^n$$

is the only one that is a spacelike mean curvature flow soliton with respect to $\mathcal{K} = t^{\frac{2}{3}} \partial_t$ and with soliton constant $c < 0$.

Example 3.2. According to the terminology introduced by Albujeer and Alías [4], a GRW spacetime $-\mathbb{R} \times_{e^t} M^n$ is called a *steady state type* spacetime. This terminology is due to the fact that the steady state model of the universe \mathcal{H}^4 , proposed by Bondi-Gold [16] and Hoyle [26] when looking for a model of the universe which looks the same not only at all points and in all directions (that is, spatially isotropic and homogeneous) but also at all times, is isometric to the RW spacetime $-\mathbb{R} \times_{e^t} \mathbb{R}^3$ (for more details, see [25]). From (11) we conclude that the slice

$$\left\{ \log\left(-\frac{n}{c}\right) \right\} \times M^n$$

is the only one that is a spacelike mean curvature flow soliton with respect to $\mathcal{K} = e^t \partial_t$ and with soliton constant $c < 0$.

Example 3.3. From [35, Example 4.2], the $(n + 1)$ -dimensional de Sitter space \mathbb{S}_1^{n+1} is isometric to the RW spacetime $-\mathbb{R} \times_{\cosh t} \mathbb{S}^n$, where \mathbb{S}^n denotes the n -dimensional unit Euclidean sphere endowed with its standard metric. Taking into account the terminology introduced in [5], the open half-space $\mathbb{R}^+ \times \mathbb{S}^n \subset \mathbb{S}_1^{n+1}$ (respect. $\mathbb{R}^- \times \mathbb{S}^n \subset \mathbb{S}_1^{n+1}$) is called the *chronological future* (respect. *past*) of \mathbb{S}_1^{n+1} with respect to the totally geodesic *equator* $\{0\} \times \mathbb{S}^n$. From (11) we see that the equator is a spacelike mean curvature flow soliton with respect to $\mathcal{K} = \cosh t \partial_t$ and constant soliton $c = 0$ and the slices

$$\left\{ \sinh^{-1}\left(\frac{-n \pm \sqrt{n^2 - 4c^2}}{2c}\right) \right\} \times \mathbb{S}^n$$

are spacelike mean curvature flow soliton with respect to $\mathcal{K} = \cosh t \partial_t$ and with soliton constant $0 < |c| \leq \frac{n}{2}$.

Example 3.4. Taking into account once more [35, Example 4.2], we consider the open region of \mathbb{S}_1^{n+1} which is isometric to the RW spacetime $-\mathbb{R}^+ \times_{\sinh t}$

\mathbb{H}^n , where \mathbb{H}^n denotes the n -dimensional hyperbolic space endowed with its standard metric. From (11) we have that the slices

$$\left\{ \cosh^{-1} \left(\frac{-n - \sqrt{n^2 + 4c^2}}{2c} \right) \right\} \times \mathbb{H}^n$$

are spacelike mean curvature flow soliton with respect to $\mathcal{K} = \sinh t \partial_t$ and with soliton constant $c < 0$.

Example 3.5. Motivated by [35, Example 4.3], we will consider the open subset of the $(n+1)$ -dimensional anti-de Sitter space \mathbb{H}_1^{n+1} which is isometric to the RW spacetime $-\left(-\frac{\pi}{2}, \frac{\pi}{2}\right) \times_{\cos t} \mathbb{H}^n$. In analogy with the nomenclature of the de Sitter space, the open half-space $(0, \frac{\pi}{2}) \times \mathbb{H}^n \subset \mathbb{H}_1^{n+1}$ (respect. $(-\frac{\pi}{2}, 0) \times \mathbb{H}^n \subset \mathbb{H}_1^{n+1}$) will be called the *chronological future* (respect. *past*) of \mathbb{H}_1^{n+1} with respect to the totally geodesic *equator* $\{0\} \times \mathbb{H}^n$. From (11) we see that the equator is a spacelike mean curvature flow soliton with respect to $\mathcal{K} = \cos t \partial_t$ and constant soliton $c = 0$ and the slices

$$\left\{ \sin^{-1} \left(\frac{-n \pm \sqrt{n^2 + 4c^2}}{2c} \right) \right\} \times \mathbb{H}^n$$

are spacelike mean curvature flow soliton with respect to $\mathcal{K} = \cos t \partial_t$ and with soliton constant $c \neq 0$.

4. Nonexistence of spacelike mean curvature flow solitons

We initiate quoting the generalized maximum principle of Omori [36] and Yau [41] (see also [11] for a modern and accessible reference to the generalized maximum principle of Omori-Yau).

Lemma 4.1. *Let Σ^n be an n -dimensional complete Riemannian manifold whose Ricci curvature is bounded from below and let $u \in C^\infty(\Sigma)$ be a smooth function which is bounded from above on Σ^n . Then there exists a sequence of points $\{p_k\}_{k \geq 1}$ in Σ^n such that*

$$\lim_k u(p_k) = \sup_\Sigma u, \quad \lim_k |\nabla u(p_k)| = 0 \quad \text{and} \quad \limsup_k \Delta u(p_k) \leq 0.$$

It is not difficult to see that Lemma 4.1 is equivalent to the following one.

Lemma 4.2. *Let Σ^n be an n -dimensional complete Riemannian manifold whose Ricci curvature is bounded from below and let $u \in C^\infty(\Sigma)$ be a smooth function which is bounded from below on Σ^n . Then there exists a sequence of points $\{p_k\}_{k \geq 1}$ in Σ^n such that*

$$\lim_k u(p_k) = \inf_\Sigma u, \quad \lim_k |\nabla u(p_k)| = 0 \quad \text{and} \quad \liminf_k \Delta u(p_k) \geq 0.$$

In the results of this subsection, we will suppose that the GRW spacetime $\overline{M}^{n+1} = -I \times_f M^n$ obeys the *strong null convergence condition* (SNCC) which

was introduced by Alías and Colares [8] and corresponds to the following suitable constraints on the sectional curvature K_M of the Riemannian fiber M^n of the GRW spacetime $-I \times_f M^n$

$$K_M \geq \sup_I (f f'' - f'^2). \tag{12}$$

We observe that the SNCC is a suitable change on the so-called *null convergence condition* (NCC), which means that the Ricci curvature of \overline{M}^{n+1} being non-negative on null or lightlike directions (for more details concerning the NCC, see [35]).

Considering an immersed spacelike hypersurface in a GRW spacetime obeying (12), the next lemma gives sufficient conditions to its Ricci curvature with respect to a suitable conformal metric be bounded from below. For this, we will suppose that such a spacelike hypersurface lies in a *timelike bounded region*

$$\mathcal{B}_{t_1, t_2} := \{(t, p) \in -I \times_f M^n : t_1 \leq t \leq t_2 \text{ and } p \in M^n\}.$$

Lemma 4.3. *Let $\overline{M}^{n+1} = -I \times_f M^n$ be a GRW spacetime which obeys (12) and let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a spacelike hypersurface contained in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$. If the second fundamental form and the hyperbolic angle function Θ are bounded, then the Ricci curvature $\widehat{\text{Ric}}$ of Σ^n with respect to the conformal metric $\hat{g} := \frac{1}{f^2(h)}g$ is bounded from below.*

PROOF. We recall that the curvature tensor R of Σ^n can be described in terms of its Weingarten operator A and the curvature tensor \overline{R} of the ambient $-I \times_f M^n$ by the so-called Gauss' equation given by

$$g(R(X, Y)Z, W) = \bar{g}(\overline{R}(X, Y)Z, W) - g(AX, Z)g(AY, W) + g(AX, W)g(AY, Z), \tag{13}$$

for every tangent vector fields $X, Y, Z \in \mathfrak{X}(\Sigma)$. Here, as in [37], the curvature tensor R is given by

$$R(X, Y)Z = \nabla_{[X, Y]}Z - [\nabla_X, \nabla_Y]Z,$$

where $[,]$ denotes the Lie bracket and $X, Y, Z \in \mathfrak{X}(\Sigma)$.

Let us consider a smooth vector field $X \in \mathfrak{X}(\Sigma)$ and take a (local) orthonormal frame $\{E_1, \dots, E_n\}$. It follows from Gauss equation (13) that the Ricci curvature Ric of Σ^n with respect to the induced metric g satisfies

$$\text{Ric}(X, X) \geq \sum_i \bar{g}(\overline{R}(X, E_i)X, E_i) - \frac{H^2}{4}|X|^2. \tag{14}$$

To estimate the first summand on the right-hand side of (14), let us consider $X^* = (\pi_M)_*(X)$ and $E_i^* = (\pi_M)_*(E_i)$. So, from [37, Proposition 7.42] and (4) we

have

$$\begin{aligned} \sum_i \bar{g}(\bar{R}(X, E_i)X, E_i) &= \sum_i g(R_M(X^*, E_i^*)X^*, E_i^*)_M \\ &\quad + (n-1)((\log f)'(h))^2 |X|^2 \\ &\quad - (n-2)(\log f)''(h)g(X, \nabla h)^2 \\ &\quad - (\log f)''(h)|\nabla h|^2 |X|^2, \end{aligned} \quad (15)$$

where R_M denotes the curvature tensor of M^n . But, writing

$$X^* = X + \bar{g}(X, \partial_t)\partial_t,$$

we can estimate the first summand on the right-hand side of (15) to get

$$\begin{aligned} \sum_i g(R_M(X^*, E_i^*)X^*, E_i^*)_M &= f^2(h)(|X^*_M|^2 |E^*_M|^2 \\ &\quad - g(X^*, E^*)_M^2)K_M(X^*, E^*) \\ &\geq \frac{1}{f^2(h)}((n-1)|X|^2 + |\nabla h|^2 |X|^2 \\ &\quad + (n-2)g(X, \nabla h)^2) \min_i K_M(X^*, E_i^*). \end{aligned} \quad (16)$$

Consequently, since our ambient space obeys (12), from (16) we have that

$$\begin{aligned} \sum_i g(R_M(X^*, E_i^*)X^*, E_i^*) &\geq ((n-1)|X|^2 + |\nabla h|^2 \\ &\quad + (n-2)g(X, \nabla h)^2)(\log f)''(h). \end{aligned} \quad (17)$$

Substituting (17) into (15), we get

$$\begin{aligned} \sum_i \bar{g}(\bar{R}(X, E_i)X, E_i) &\geq ((n-1)|X|^2 + |\nabla h|^2 \\ &\quad + (n-2)g(X, \nabla h)^2)(\log f)''(h) \\ &\quad + (n-1)((\log f)'(h))^2 |X|^2 \\ &\quad - (n-2)(\log f)''(h)g(X, \nabla h)^2 \\ &\quad - (\log f)''(h)|\nabla h|^2 |X|^2 \\ &= (n-1)\frac{f''(h)}{f(h)} |X|^2. \end{aligned} \quad (18)$$

Then, taking into account that $|A|^2 \geq \frac{H^2}{n}$, from (14) and (18) we arrive at the following lower estimate

$$\text{Ric}(X, X) \geq -\left((n-1)\frac{|f''(h)|}{f(h)} + \frac{n|A|^2}{4} \right) |X|^2. \quad (19)$$

On the other hand, we have the following equation (see, for instance, [15, Section 1J], [30, Section A] or [39, page 168])

$$\widehat{\text{Ric}}(X, X) = \text{Ric}(X, X) + \frac{1}{f^2(h)} \left\{ (n-2)f(h)\nabla^2 f(h)(X, X) + (f(h)\Delta f(h) - (n-1)|\nabla f(h)|^2)|X|^2 \right\}. \tag{20}$$

Consequently, from equation (20) we get

$$\begin{aligned} \widehat{\text{Ric}}(X, X) = \text{Ric}(X, X) + \frac{1}{f^2(h)} \left\{ (n-2)f(h)(f''(h)g(\nabla h, X)^2 \right. \\ \left. + f'(h)\nabla^2 h(X, X)) + (f(h)(f''(h)|\nabla h|^2 + f'(h)\Delta h) \right. \\ \left. - (n-1)(f'(h))^2|\nabla h|^2)|X|^2 \right\}. \end{aligned} \tag{21}$$

Hence, considering (5), (7), (8) and (19) into (21), we obtain after a straightforward computation the following lower estimate

$$\begin{aligned} \widehat{\text{Ric}}(X, X) \geq \left\{ (n-1)\frac{(f'(h))^2}{f(h)} - (n-1)\left(\frac{|f''(h)|}{f(h)} \right. \right. \\ \left. \left. + (n+1)\frac{(f'(h))^2}{f^2(h)}\right)\Theta^2 - (n-\sqrt{n}-2)\frac{|f'(h)|}{f(h)}|A||\Theta| \right. \\ \left. - \frac{n|A|^2}{4} \right\}|X|^2. \end{aligned} \tag{22}$$

Therefore, taking into account that $|A|$ and $|\Theta|$ are bounded and that Σ^n lies in a timelike bounded region of the ambient space, from (22) we conclude that $\widehat{\text{Ric}}$ is bounded from below. □

Aiming to simplify the notation, along our main results we will consider the *modified soliton function* as being the function

$$\bar{\zeta}_c(t) := f'(t)\zeta_c(t), \tag{23}$$

where ζ_c is the soliton function defined in (10). So, we are in a position to state and prove our first nonexistence result concerning spacelike mean curvature flow solitons immersed in a GRW spacetime.

Theorem 4.4. *Let $\overline{M}^{n+1} = -I \times_f M^n$ be a GRW spacetime satisfying (12). There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = f(t)\partial_t$ with soliton constant c , whose second fundamental form and hyperbolic angle function are bounded, and lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$ such that $\bar{\zeta}_c(t)$ has strict sign for all $t \in [t_1, t_2]$.*

PROOF. By contradiction, let us assume the existence of a complete spacelike mean curvature flow soliton ψ satisfying the assumptions of Theorem 4.4. As before, consider on Σ^n the metric $\hat{g} = \frac{1}{f^2(h)}g$, which is conformal to its induced

metric g . Denoting by $\hat{\Delta}$ the Laplacian with respect to the metric \hat{g} , from (5) and (8) we have

$$\hat{\Delta}h = -f(h)f'(h)\{n + (n-1)|\nabla h|^2\} - Hf^2(h)\Theta. \quad (24)$$

Thus, from (24) we get

$$\begin{aligned} \hat{\Delta}f(h) &= -nf(h)(f'(h))^2 - Hf'(h)f^2(h)\Theta \\ &\quad + f^3(h)\{(\log f)''(h) - (n-2)((\log f)'(h))^2\}|\nabla h|^2. \end{aligned} \quad (25)$$

For any positive real number α , with a straightforward computation from (25) we get

$$\begin{aligned} \hat{\Delta}f^{-\alpha}(h) &= -\alpha f^{-\alpha-1}(h)\{-nf(h)(f'(h))^2 - Hf'(h)f^2(h)\Theta \\ &\quad + f^3(h)\{(\log f)''(h) - (n+\alpha-3)((\log f)'(h))^2\}|\nabla h|^2\} \\ &= -\alpha f^{-\alpha-1}(h)\{-nf(h)(f'(h))^2\Theta^2 - Hf'(h)f^2(h)\Theta \\ &\quad + f^3(h)\{(\log f)''(h) - (\alpha-3)((\log f)'(h))^2\}|\nabla h|^2\}. \end{aligned} \quad (26)$$

Hence, from (9), (23) and (26) we obtain

$$\begin{aligned} \hat{\Delta}f^{-\alpha}(h) &= \alpha f^{-\alpha}(h)\Theta^2\bar{\zeta}_c(h) - \alpha f^{2-\alpha}(h)\{(\log f)''(h) \\ &\quad - (\alpha-3)((\log f)'(h))^2\}|\nabla h|^2. \end{aligned} \quad (27)$$

At this point, let us assume that $\bar{\zeta}_c(t) > 0$ for all $t_1 \leq t \leq t_2$. Since we are assuming that $|A|$ and Θ are bounded, we can apply Lemmas 4.1 and 4.3 to guarantee the existence of a sequence of points $\{p_k\}_{k \geq 1}$ in Σ^n such that

$$\begin{aligned} \lim_k f^{-\alpha}(h)(p_k) &= \sup_{\Sigma} f^{-\alpha}(h), \quad \lim_k |\hat{\nabla} f^{-\alpha}(h)(p_k)|_{\hat{g}} = 0, \\ \text{and } \limsup_k \hat{\Delta}f^{-\alpha}(h)(p_k) &\leq 0, \end{aligned} \quad (28)$$

where $|\cdot|_{\hat{g}}$ and $\hat{\nabla}$ denote, respectively, the norm and gradient with respect to the metric \hat{g} .

But, it is not difficult to verify that

$$|\hat{\nabla} f^{-\alpha}(h)|_{\hat{g}} = \alpha f^{-\alpha}(h)|f'(h)||\nabla h|. \quad (29)$$

So, since $\Sigma^n \subset \mathcal{B}_{t_1, t_2}$, with $|f'(t)| > 0$ for all $t_1 \leq t \leq t_2$, from (28) and (29) we get that

$$\lim_k |\nabla h(p_k)| = 0. \quad (30)$$

Consequently, from (5) and (30) we have that

$$\lim_k \Theta^2(p_k) = 1. \quad (31)$$

Moreover, from (27), (30) and (31) we obtain

$$\begin{aligned}
 0 \geq \limsup_k \hat{\Delta} f^{-\alpha}(h)(p_k) &\geq \alpha \limsup_k \left\{ f^{-\alpha}(h) \Theta^2 \bar{\zeta}_c(h) \right\}(p_k) \\
 &\quad - \alpha \limsup_k \left\{ f^{2-\alpha}(h) \right\} (\log f)''(h) \\
 &\quad - (\alpha - 3) ((\log f)'(h))^2 \|\nabla h\|^2(p_k) \quad (32) \\
 &= \alpha \sup_{\Sigma} f^{-\alpha}(h) \limsup_k \bar{\zeta}_c(p_k) \geq 0.
 \end{aligned}$$

Hence, since $\sup_{\Sigma} f^{-\alpha}(h) > 0$ and $\bar{\zeta}_c(t) > 0$ for all $t_1 \leq t \leq t_2$, (32) gives us a contradiction.

Finally, in the case $\bar{\zeta}_c(t) < 0$ for all $t_1 \leq t \leq t_2$, using Lemma 4.2 instead of Lemma 4.1 we get

$$\begin{aligned}
 0 \leq \liminf_k \hat{\Delta} f^{-\alpha}(h)(p_k) &\leq \alpha \liminf_k \left\{ f^{-\alpha}(h) \Theta^2 \bar{\zeta}_c(h) \right\}(p_k) \\
 &\quad + \alpha \liminf_k \left\{ f^{2-\alpha}(h) \right\} (\log f)''(h) \\
 &\quad - (\alpha - 3) ((\log f)'(h))^2 \|\nabla h\|^2(p_k) \quad (33) \\
 &= \alpha \inf_{\Sigma} f^{-\alpha}(h) \liminf_k \bar{\zeta}_c(p_k) \leq 0.
 \end{aligned}$$

Therefore, since $\inf_{\Sigma} f^{-\alpha}(h) > 0$ and $\bar{\zeta}_c(t) < 0$ for all $t_1 \leq t \leq t_2$, (33) also leads us to a contradiction. \square

Taking into account that the $(n+1)$ -dimensional Einstein-de Sitter spacetime $-\mathbb{R}^+ \times_{\frac{2}{t^3}} \mathbb{R}^n$ (see Example 3.1) satisfies (12), from Theorem 4.4 we obtain the following consequence.

Corollary 4.5. *Let $\overline{M}^{n+1} = -\mathbb{R}^+ \times_{\frac{2}{t^3}} \mathbb{R}^n$ be the $(n + 1)$ -dimensional Einstein-de Sitter spacetime. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = t^{\frac{2}{3}} \partial_t$ with soliton constant $c \geq 0$, whose second fundamental form and hyperbolic angle function are bounded, and lying in a timelike bounded region of \overline{M}^{n+1} .*

Since a steady state type spacetime (see Example 3.2) whose Riemannian fiber has nonnegative sectional curvature satisfies (12), from Theorem 4.4 we obtain the following application.

Corollary 4.6. *Let $\overline{M}^{n+1} = -\mathbb{R} \times_{e^t} M^n$ be a steady state type spacetime whose Riemannian fiber M^n has nonnegative sectional curvature. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = e^t \partial_t$ with soliton constant $c \geq 0$, whose second fundamental form and hyperbolic angle function are bounded, and lying in a timelike bounded region of \overline{M}^{n+1} .*

Considering the context of Example 3.3, from Theorem 4.4 we also get.

Corollary 4.7. *There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \mathbb{S}_1^{n+1}$ with respect to $\mathcal{K} = \cosh t \partial_t$ having soliton constant $c \geq 0$ (respect. $c \leq 0$), whose second fundamental form and hyperbolic angle function are bounded, and lying in a timelike bounded region contained in the chronological future (respect. past) of \mathbb{S}_1^{n+1} with respect to the equator $\{0\} \times \mathbb{S}^n$.*

From Example 3.4 and Theorem 4.4 we obtain.

Corollary 4.8. *There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow -\mathbb{R}^+ \times_{\sinh t} \mathbb{H}^n \subset \mathbb{S}_1^{n+1}$ with respect to $\mathcal{K} = \sinh t \partial_t$ having soliton constant $c \geq 0$, whose second fundamental form and hyperbolic angle function are bounded, and lying in a timelike bounded region of $-\mathbb{R}^+ \times_{\sinh t} \mathbb{H}^n \subset \mathbb{S}_1^{n+1}$.*

Finally, in the setting of Example 3.5, Theorem 4.4 reads as follows.

Corollary 4.9. *There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow -(\frac{\pi}{2}, \frac{\pi}{2}) \times_{\cos t} \mathbb{H}^n \subset \mathbb{H}_1^{n+1}$ with respect to $\mathcal{K} = \cos t \partial_t$ having soliton constant $c \leq 0$ (respect. $c \geq 0$), whose second fundamental form and hyperbolic angle function are bounded, and lying in a timelike bounded region contained in the chronological future (respect. past) of \mathbb{H}_1^{n+1} with respect to the equator $\{0\} \times \mathbb{H}^n$.*

5. Uniqueness and further nonexistence results

We will also adopt the following notation

$$\mathcal{L}_g^p(\Sigma) := \{u : \Sigma^n \rightarrow \mathbb{R} : \int_{\Sigma} |u|^p d\Sigma < +\infty\},$$

where $d\Sigma$ stands for the measure related to the metric g . With this, we quote an extension of Hopf's theorem on a complete Riemannian manifold (Σ^n, g) due to Yau in [42].

Lemma 5.1. *Let u be a smooth function defined on a complete Riemannian manifold (Σ^n, g) , such that Δu does not change sign on Σ^n . If $|\nabla u| \in \mathcal{L}_g^1(\Sigma)$, then Δu vanishes identically on Σ^n .*

In what follows, we will assume that the warping function f of the ambient GRW spacetime $\overline{M}^{n+1} = -I \times_f M^n$ satisfies the following inequality

$$(\log f)'' \leq \gamma((\log f)')^2, \quad (34)$$

for some nonnegative constant γ . As it was observed in [6], the inequality (34) is a mild hypothesis due to the fact that, for instance, when \overline{M}^{n+1} obeys the SNCC (respect. NCC) and its Riemannian fiber M^n is flat (respect. Ricci-flat), we have that (34) is automatically satisfied.

Returning to the study of spacelike mean curvature flow solitons, we get the following result.

Theorem 5.2. *Let $\overline{M}^{n+1} = -I \times_f M^n$ be a GRW spacetime satisfying (34), occurring the equality only at isolated points of I , and whose Riemannian fiber M^n is complete. Let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a complete spacelike mean curvature flow soliton with respect to $\mathcal{K} = f(t)\partial_t$ and with soliton constant c , lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$ such that $\bar{\zeta}_c(t) \geq 0$ for all $t_1 \leq t \leq t_2$. If the height function h is such that $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$, then Σ^n is a slice M_{t_*} for some $t_* \in [t_1, t_2]$ which is implicitly given by the condition $\zeta_c(t_*) = 0$.*

PROOF. We will consider again the conformal metric $\hat{g} := \frac{1}{f^2(h)}g$ and we will take $\alpha = \gamma + 3$. Since we are assuming that Σ^n lies in \mathcal{B}_{t_1, t_2} and that $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$, from (29) we get that $|\hat{\nabla} f^{-\alpha}(h)|_{\hat{g}} \in \mathcal{L}_{\hat{g}}^1(\Sigma)$.

Moreover, since $\bar{\zeta}_c(t) \geq 0$ for all $t_1 \leq t \leq t_2$, from (27) and (34) we obtain that $\hat{\Delta} f^{-\alpha}(h) \geq 0$. Consequently, we can apply Lemma 5.1 to infer that $\hat{\Delta} f^{-\alpha}(h) = 0$ on Σ^n .

Therefore, since we are assuming that the equality occurs in (34) just only at isolated points of I , returning to (27) we conclude that $|\nabla h|$ must vanishes identically on Σ^n . Therefore, Σ^n must be a slice M_{t_*} for some $t_* \in [t_1, t_2]$ which is implicitly given by the condition $\zeta_c(t_*) = 0$. □

Applying Theorem 5.2 to the Einstein-de Sitter spacetime, we obtain the following result.

Corollary 5.3. *Let $\overline{M}^{n+1} = -\mathbb{R}^+ \times_{t^{\frac{2}{3}}} \mathbb{R}^n$ be the $(n + 1)$ -dimensional Einstein-de Sitter spacetime. The only complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = t^{\frac{2}{3}}\partial_t$ with soliton constant $c < 0$, lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$ with $t_2 = (-\frac{2n}{3c})^{\frac{3}{5}}$, and such that its height function h satisfies $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$, is the slice $\{(-\frac{2n}{3c})^{\frac{3}{5}}\} \times \mathbb{R}^n$.*

When the ambient space is a steady state type spacetime, Theorem 5.2 gives us the following consequence.

Corollary 5.4. *Let $\overline{M}^{n+1} = -\mathbb{R} \times_{e^t} M^n$ be a steady state type spacetime whose Riemannian fiber M^n is complete. The only complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = e^t\partial_t$ with soliton constant $c < 0$, lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$ with $t_2 = \log(-\frac{n}{c})$, and such that its height function h satisfies $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$, is the slice $\{\log(-\frac{n}{c})\} \times M^n$.*

From Theorem 5.2 we also get the following nonexistence result.

Corollary 5.5. *Let $\overline{M}^{n+1} = -I \times_f M^n$ be a GRW spacetime satisfying (34), occurring the equality only at isolated points of I , and whose Riemannian fiber M^n*

is complete. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = f(t)\partial_t$ and with soliton constant c , lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$ with $\bar{\zeta}_c(t) > 0$ for all $t_1 \leq t \leq t_2$, and such that its height function h satisfies $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$.

When the ambient spacetime is the Einstein-de Sitter spacetime and a steady state type spacetime, respectively, Corollary 5.5 reads as follows.

Corollary 5.6. Let $\overline{M}^{n+1} = -\mathbb{R}^+ \times_{t^{\frac{2}{3}}} \mathbb{R}^n$ be the $(n + 1)$ -dimensional Einstein-de Sitter spacetime. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = t^{\frac{2}{3}}\partial_t$ with soliton constant $c \geq 0$, lying in a timelike bounded region of \overline{M}^{n+1} and such that its height function h satisfies $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$.

Corollary 5.7. Let $\overline{M}^{n+1} = -\mathbb{R} \times_{e^t} M^n$ be a steady state type spacetime whose Riemannian fiber M^n is complete. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = e^t\partial_t$ with soliton constant $c \geq 0$, lying in a timelike bounded region of \overline{M}^{n+1} and such that its height function h satisfies $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$.

We are also able to present a slight different version of Theorem 5.2.

Theorem 5.8. Let $\overline{M}^{n+1} = -I \times_f M^n$ be a GRW spacetime satisfying (34) whose Riemannian fiber M^n is complete. Let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a complete spacelike mean curvature flow soliton with respect to $\mathcal{K} = f(t)\partial_t$ with soliton constant c , lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$ with $\bar{\zeta}_c(t) \geq 0$ and $f'(t)$ vanishing only in isolated points of $[t_1, t_2]$. If $|\nabla h| \in \mathcal{L}_g^1(\Sigma)$, then Σ^n is a slice M_{t_*} for some $t_* \in [t_1, t_2]$ which is implicitly given by the condition $\zeta_c(t_*) = 0$.

PROOF. As in the proof of Theorem 5.2, we get that $\hat{\Delta}f^{-\alpha}(h) = 0$ on Σ^n , for $\alpha = \gamma + 3$. Moreover, since Σ^n lies in \mathcal{B}_{t_1, t_2} , we can also verify that $|\widehat{\nabla}f^{-2\alpha}(h)|_{\hat{g}} \in \mathcal{L}_g^1(\Sigma)$. But, we note that

$$\hat{\Delta}f^{-2\alpha}(h) = 2f^{-\alpha}(h)\hat{\Delta}f^{-\alpha}(h) + 2|\widehat{\nabla}f^{-\alpha}(h)|_{\hat{g}}^2 = 2|\widehat{\nabla}f^{-\alpha}(h)|_{\hat{g}}^2 \geq 0. \tag{35}$$

Thus, we can apply again Lemma 5.1 to obtain that $\hat{\Delta}f^{-2\alpha}(h) = 0$ on Σ^n . Hence, since we are assuming that $f'(t) > 0$ for $t_1 \leq t \leq t_2$, from (29) and (35) we obtain that $|\nabla h| = 0$ on Σ^n . Therefore, Σ^n must be a slice M_{t_*} for some $t_* \in [t_1, t_2]$ which is implicitly given by the condition $\zeta_c(t_*) = 0$. \square

At this point, we need quoting two other auxiliary results also due to Yau in [42].

Lemma 5.9. If u is a nonnegative smooth subharmonic function defined on (Σ^n, g) , with $u \in \mathcal{L}_g^p(\Sigma)$ for some $p > 1$, then u must be constant.

Lemma 5.10. *All complete noncompact Riemannian manifolds with nonnegative Ricci curvature have at least linear volume growth.*

These previous lemmas enable us to prove the following nonexistence result.

Theorem 5.11. *Let $\overline{M}^{-n+1} = -I \times_f M^n$ be a GRW spacetime satisfying (34), occurring the equality only at isolated points of I , and whose Riemannian fiber M^n is complete noncompact with nonnegative Ricci curvature. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{-n+1}$ with respect to $\mathcal{K} = f(t)\partial_t$ and soliton constant c , lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{-n+1}$, with $\bar{\zeta}_c(t) \geq 0$ for all $t_1 \leq t \leq t_2$, and whose height function h is such that $(f(h))^{-1} \in \mathcal{L}_g^q(\Sigma)$ for some q with $q > \gamma + 3$.*

PROOF. Supposing by contradiction the existence of such a spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{-n+1}$ and taking once more $\alpha = \gamma + 3$, from (27) we obtain that $\hat{\Delta} f^{-\alpha}(h) \geq 0$ on Σ^n . Moreover, since Σ^n is contained in a timelike bounded region and $(f(h))^{-1} \in \mathcal{L}_g^q(\Sigma)$ for some q with $q > \alpha$, it is not difficult to verify that $f^{-\alpha}(h) \in \mathcal{L}_g^p(\Sigma)$ for $p = \frac{q}{\alpha} > 1$. Thus, we can apply Lemma 5.9 to get that $f(h)$ is constant on Σ^n . Hence, since we are also supposing that the equality occurs in (34) just only at isolated points of I , returning to (27) we conclude that $|\nabla h|$ must vanish identically on Σ^n . Consequently, Σ^n is isometric (up to scaling) to M^n . So, since $f(h)$ is a positive constant, our assumption that $f(h) \in \mathcal{L}_g^q(\Sigma)$ also implies that M^n has finite volume. But, since M^n is a complete non-compact with nonnegative Ricci curvature, Lemma 5.10 leads us to a contradiction. \square

From Theorem 5.11 we obtain the following consequences.

Corollary 5.12. *Let $\overline{M}^{-n+1} = -\mathbb{R}^+ \times_{t^{\frac{2}{3}}} \mathbb{R}^n$ be the $(n + 1)$ -dimensional Einstein-de Sitter spacetime. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{-n+1}$ with respect to $\mathcal{K} = t^{\frac{2}{3}}\partial_t$ with soliton constant $c \geq 0$, lying in a timelike bounded region of \overline{M}^{-n+1} and such that its height function h satisfies $h^{-\frac{2}{3}} \in \mathcal{L}_g^q(\Sigma)$ for some q with $q > 3$.*

Corollary 5.13. *Let $\overline{M}^{-n+1} = -\mathbb{R} \times_{e^t} M^n$ be a steady state type spacetime whose Riemannian fiber M^n is complete noncompact with nonnegative Ricci curvature. There does not exist a complete spacelike mean curvature flow soliton $\psi : \Sigma^n \rightarrow \overline{M}^{-n+1}$ with respect to $\mathcal{K} = e^t \partial_t$ with soliton constant $c \geq 0$, lying in a timelike bounded region of \overline{M}^{-n+1} and such that its height function h satisfies $e^{-h} \in \mathcal{L}_g^q(\Sigma)$ for some q with $q > 3$.*

According to [38], a GRW spacetime $\overline{M}^{n+1} = I \times_f M^n$ is said to be *spatially parabolic* when its Riemannian fiber M^n is parabolic, that is, (M^n, g_M) is a non-compact complete Riemannian manifold such that the only superharmonic functions on it that are bounded from below are the constants. Analogously, \overline{M}^{n+1} is said to be *spatially parabolic covered* when its universal Lorentzian covering is spatially parabolic. For our next uniqueness result, we need of the following parabolicity criterion due to Aledo, Rubio and Salamanca (see [6, Theorem 2.2])

Lemma 5.14. *Let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a complete spacelike hypersurface immersed in a spatially parabolic covered GRW spacetime $\overline{M}^{n+1} = I \times_f M^n$. If $\sup_{\Sigma} f(h) < +\infty$ and the hyperbolic angle function Θ is bounded, then (Σ^n, \hat{g}) , endowed with the conformal metric $\hat{g} = \frac{1}{f^2(h)}g$, is parabolic.*

Using Lemma 5.14 we obtain the following result.

Theorem 5.15. *Let $\overline{M}^{n+1} = I \times_f M^n$ be a spatially parabolic covered GRW spacetime satisfying (34), holding the equality only at isolated points of I . Let $\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ be a complete spacelike mean curvature flow soliton with respect to $\mathcal{K} = f(t)\partial_t$ with soliton constant c , lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$, with $\tilde{\zeta}_c(t) \geq 0$ for all $t_1 \leq t \leq t_2$. If the hyperbolic angle function Θ is bounded, then Σ^n is a slice M_{t_*} for some $t_* \in [t_1, t_2]$ which is implicitly given by the condition $\zeta_c(t_*) = 0$.*

PROOF. First, we note that Lemma 5.14 guarantees that (Σ^n, \hat{g}) is parabolic. Moreover, it follows from (27) that $f(h)^{-\alpha}$ (where $\alpha = \gamma + 3$) is subharmonic on Σ^n . Thus, since the hypothesis that $\Sigma^n \subset \mathcal{B}_{t_1, t_2}$ implies in particular that $f(h)^{-\alpha}$ is bounded from above, it follows from the parabolicity of (Σ^n, \hat{g}) that $f(h)$ is constant on Σ^n . Consequently, since we are assuming that the equality holds in (34) only at isolated points of I , returning to (27) we conclude that $|\nabla h| = 0$ on Σ^n , which means that Σ^n is a slice. \square

We close this section quoting the following applications of Theorem 5.15.

Corollary 5.16. *Let $\overline{M}^3 = -\mathbb{R}^+ \times_{\frac{2}{t^3}} \mathbb{R}^2$ be the 3-dimensional Einstein-de Sitter spacetime. The only complete spacelike mean curvature flow soliton $\psi : \Sigma^2 \rightarrow \overline{M}^3$ with respect to $\mathcal{K} = t^{\frac{2}{3}}\partial_t$ with soliton constant $c < 0$, lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^3$ with $t_2 = (-\frac{4}{3c})^{\frac{3}{5}}$, and such that its hyperbolic angle function Θ is bounded, is the slice $\{(-\frac{4}{3c})^{\frac{3}{5}}\} \times \mathbb{R}^n$.*

Corollary 5.17. *Let $\overline{M}^{n+1} = -\mathbb{R} \times_{e^t} M^n$ be a spatially parabolic covered steady state type spacetime. The only complete spacelike mean curvature flow soliton*

$\psi : \Sigma^n \rightarrow \overline{M}^{n+1}$ with respect to $\mathcal{K} = e^t \partial_t$ with soliton constant $c < 0$, lying in a timelike bounded region $\mathcal{B}_{t_1, t_2} \subset \overline{M}^{n+1}$ with $t_2 = \log(-\frac{n}{c})$, and such that its hyperbolic angle function Θ is bounded, is the slice $\{\log(-\frac{n}{c})\} \times M^n$.

Remark 5.18. Related to Corollary 5.17 in the case $n = 2$, when the Riemannian fiber M^2 is a complete Riemannian surface having nonnegative Gaussian curvature, a classical result due to Ahlfors [2] and Blanc-Fiala-Huber [27] guarantees that M^2 has parabolic universal covering.

6. Entire spacelike mean curvature flow graphs

In this last section, we will use the theorems of the previous section in order to establish new Calabi-Bernstein type results concerning entire spacelike graphs constructed over the Riemannian fiber of a GRW spacetime. Before, we need to recall some basic facts related to these spacelike graphs.

Let $\Omega \subseteq M^n$ be a connected domain and let $u \in C^\infty(\Omega)$ be a smooth function such that $u(\Omega) \subseteq I$, then $\Sigma(u)$ will denote the (vertical) graph over Ω determined by u , that is,

$$\Sigma(u) = \{(u(x), x) : x \in \Omega\} \subset \overline{M}^{n+1} = -I \times_f M^n.$$

The graph is said to be entire if $\Omega = M^n$. Observe that $h(u(x), x) = u(x)$, for all $x \in \Omega$. Hence, h and u can be identified in a natural way. The metric induced on Ω from the Lorentzian metric of the ambient GRW spacetime via $\Sigma(u)$ is

$$g_u = -du^2 + f^2(u)g_M. \tag{36}$$

It can be easily seen from (36) that such a graph $\Sigma(u)$ is a spacelike hypersurface if and only if $|Du|_M < f(u)$, where Du stands for the gradient of u in M and $|Du|_M$ its norm, both with respect to the metric g_M . On the other hand, in the case where M^n is a simply connected manifold, from [12, Lemma 3.1] we have that every complete spacelike hypersurface $\psi : \Sigma^n \rightarrow -I \times_f M^n$ such that the warping function f is bounded on Σ^n is an entire spacelike graph over M^n . In particular, this happens for complete spacelike hypersurfaces lying in a timelike bounded region of $-I \times_f M^n$. It is also interesting to point out that, in contrast to the case of graphs in a Riemannian space, an entire spacelike graph $\Sigma(u)$ in a GRW spacetime is not necessarily complete, in the sense that the induced Riemannian metric (36) is not necessarily complete on M^n . For instance, Albuje constructed explicit examples of noncomplete entire maximal spacelike graphs (that is, whose mean curvature is identically zero) in the Lorentzian product space $-\mathbb{R} \times \mathbb{H}^2$ (see [3, Section 3]).

The future-pointing Gauss map of a spacelike graph $\Sigma(u)$ over Ω is given by the vector field

$$N = \frac{f(u)}{\sqrt{f^2(u) - |Du|_M^2}} \left(\partial_t + \frac{Du}{f^2(u)} \right). \tag{37}$$

The shape operator related to the future-pointing Gauss map (37) is given by

$$\begin{aligned}
 AX = & -\frac{1}{f(u)\sqrt{f^2(u) - |Du|_M^2}}D_X Du - \frac{f'(u)}{\sqrt{f^2(u) - |Du|_M^2}}X \\
 & + \left(\frac{-g_M(D_X Du, Du)}{f(u)(f^2(u) - |Du|_M^2)^{3/2}} + \frac{f'(u)g_M(Du, X)}{(f^2(u) - |Du|_M^2)^{3/2}} \right) Du,
 \end{aligned} \tag{38}$$

for any vector field X tangent to Ω , where D denotes the Levi-Civita connection of (M^n, g_M) . Consequently, if $\Sigma(u)$ is a spacelike graph defined over a domain $\Omega \subseteq M^n$, it is not difficult to verify from (38) that the future mean curvature function $H(u)$ of $\Sigma(u)$ is given by the following nonlinear differential equation:

$$\begin{aligned}
 H(u) = & \operatorname{div}_M \left(\frac{Du}{f(u)\sqrt{f^2(u) - |Du|_M^2}} \right) \\
 & + \frac{f'(u)}{\sqrt{f^2(u) - |Du|_M^2}} \left(n + \frac{|Du|_M^2}{f^2(u)} \right),
 \end{aligned} \tag{39}$$

where div_M stands for the divergence operator computed in the metric g_M .

Hence, from (9) and (39) we have that $\Sigma(u)$ is a spacelike mean curvature flow soliton with respect to $K = f(t)\partial_t$ with soliton constant c if, and only if, $|Du|_M < f(u)$ and u is a solution of the following nonlinear differential equation:

$$\begin{aligned}
 \operatorname{div}_M \left(\frac{Du}{f(u)\sqrt{f(u)^2 - |Du|_M^2}} \right) = & -\frac{1}{\sqrt{f(u)^2 - |Du|_M^2}} \left\{ cf(u)^2 \right. \\
 & \left. + f'(u) \left(n + \frac{|Du|_M^2}{f(u)^2} \right) \right\}.
 \end{aligned} \tag{40}$$

We say that $u \in C^\infty(M)$ has finite C^2 norm when

$$\|u\|_{C^2(M)} := \sup_{|k| \leq 2} |D^k u|_{L^\infty(M)} < +\infty.$$

In this setting, we obtain a nonparametric version of Theorem 4.4.

Theorem 6.1. *Let $\bar{M}^{-n+1} = -I \times_f M^n$ be a GRW spacetime satisfying (12) and whose Riemannian fiber M^n is complete. Suppose that c is a constant such that the modified soliton function $\bar{\zeta}_c(t)$ has strict sign in I . There does not exist a smooth function $u : M^n \rightarrow I$ with finite C^2 norm which is solution of the spacelike mean curvature flow soliton equation (40) and such that $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$.*

PROOF. Let us assume the existence of such a smooth function $u : M^n \rightarrow I$. It follows from (38) that the shape operator A of an entire spacelike graph $\Sigma(u)$ is bounded provided that u has finite C^2 norm. Note also that the finiteness of the C^2 norm of u implies, in particular, that u is bounded, which, in turn, guarantees that $\Sigma(u)$ is contained in a bounded timelike region of \overline{M}^{n+1} .

On the other hand, under the assumptions of the theorem, $\Sigma(u)$ is a complete spacelike hypersurface. Indeed, proceeding as in [9, Corollary 5.1], from (36) and the Cauchy-Schwarz inequality we get

$$\begin{aligned} g_u(X, X) &= -g_M(Du, X^*)^2 + f^2(u)g_M(X^*, X^*) \\ &\geq (f^2(u) - |Du|_M^2)g_M(X^*, X^*), \end{aligned} \tag{41}$$

for every tangent vector field X on $\Sigma(u)$, where (as before) X^* denotes the projection of X onto the Riemannian fiber M^n . Thus, since $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$, from (41) we get that

$$g_u(X, X) \geq \delta g_M(X^*, X^*), \tag{42}$$

where $\delta = (1 - \beta^2) \inf_M f^2(u)$. So, (42) implies that $L = \sqrt{\delta}L_M$, where L and L_M denote the length of a curve on $\Sigma(u)$ with respect to the Riemannian metrics g_u and g_M , respectively. As a consequence, since we are always assuming that M^n is complete, the induced metric g_u must be also complete.

Moreover, from (37) we obtain that the hyperbolic angle function Θ of $\Sigma(u)$ is given by

$$\Theta = -\frac{f(u)}{\sqrt{f^2(u) - |Du|_M^2}}. \tag{43}$$

Hence, using once more that hypothesis that $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$, from (43) we get that Θ is bounded. But, by applying Theorem 4.4 we have that $\Sigma(u)$ cannot exist. \square

From Theorem 6.1 we obtain the following applications.

Corollary 6.2. *For any constants $c \geq 0$ and $0 < \beta < 1$, there does not exist a smooth function $u : \mathbb{R}^n \rightarrow \mathbb{R}^+$ with finite C^2 norm which is a solution of the following system*

$$\begin{cases} \operatorname{div}_{\mathbb{R}^n} \left(\frac{Du}{u^{\frac{2}{3}} \sqrt{u^{\frac{4}{3}} - |Du|_{\mathbb{R}^n}^2}} \right) = -\frac{1}{\sqrt{u^{\frac{4}{3}} - |Du|_{\mathbb{R}^n}^2}} \left(cu^{\frac{4}{3}} + \frac{2n}{3u^{\frac{1}{3}}} + \frac{2|Du|_{\mathbb{R}^n}^2}{3u^{\frac{5}{3}}} \right) \\ |Du|_{\mathbb{R}^n} \leq \beta u^{\frac{2}{3}} \end{cases} \tag{44}$$

Corollary 6.3. *Let M^n be a complete Riemannian manifold with nonnegative sectional curvature. For any constants $c \geq 0$ and $0 < \beta < 1$, there does not exist*

a smooth function $u : M^n \rightarrow I$ with finite C^2 norm which is a solution of the following system

$$\begin{cases} \operatorname{div}_M \left(\frac{Du}{e^u \sqrt{e^{2u} - |Du|_M^2}} \right) = -\frac{1}{\sqrt{e^{2u} - |Du|_M^2}} \left(ce^{2u} + ne^u + \frac{|Du|_M^2}{e^u} \right) \\ |Du|_M \leq \beta e^u \end{cases} \quad (45)$$

Remark 6.4. From Examples 3.3, 3.4 and 3.5, it is not difficult to see that we can also obtain applications of Theorem 6.1 to the de Sitter and anti-de Sitter spaces similar to Corollaries 6.2 and 6.3.

Proceeding, from Theorem 5.2 we obtain the following Calabi-Bernstein type result.

Theorem 6.5. Let $\overline{M}^{\overline{n+1}} = -I \times_f M^n$ be a GRW spacetime satisfying (34), occurring the equality only at isolated points of I , and whose Riemannian fiber M^n is complete. Suppose that c is a constant such that $\zeta_c(t) \geq 0$ for all $t \in I$. If $\Sigma(u) \subset \overline{M}^{\overline{n+1}}$ is an entire spacelike graph determined by a bounded function $u \in C^\infty(M)$ which is solution of the spacelike mean curvature flow soliton equation (40) with $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$, and $|Du|_M \in \mathcal{L}_{g_M}^1(M)$, then $u \equiv t_*$ for some $t_* \in I$ which is implicitly given by the condition $\zeta_c(t_*) = 0$.

PROOF. Since we are supposing that $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$, it follows from the proof of Theorem 6.1 that $\Sigma(u)$ is a complete spacelike hypersurface.

On the other hand, reasoning once more as in [9, Corollary 5.1], we deduce from the induced metric (36) that $d\Sigma = \sqrt{|G|} dM$, where dM and $d\Sigma$ stand for the Riemannian volume elements of (M^n, g_M) and $(\Sigma(u), g_u)$, respectively, and $G = \det(g_{ij})$ with

$$g_{ij} = g_u(E_i, E_j) = f^2(u)\delta_{ij} - E_i(u)E_j(u). \quad (46)$$

Here, $\{E_1, \dots, E^n\}$ denotes a local orthonormal frame with respect to the metric g_M . So, it is not difficult to verify that

$$|G| = f^{2(n-1)}(u)(f^2(u) - |Du|_M^2). \quad (47)$$

Hence, from (46) and (47) we obtain

$$d\Sigma = f^{n-1}(u)\sqrt{f^2(u) - |Du|_M^2} dM. \quad (48)$$

Moreover, since we have that $N = N^* - \Theta \partial_t$, from (4) we get

$$|\nabla h|^2 = f^2(u)|N^*|_M^2. \quad (49)$$

Thus, from (37) and (49) we obtain

$$|\nabla h|^2 = \frac{|Du|_M^2}{f^2(u) - |Du|_M^2}. \quad (50)$$

Consequently, from (50) and (48) we get

$$|\nabla h|d\Sigma = f^{n-1}(u)|Du|_M dM. \tag{51}$$

Hence, since we are assuming that u is bounded with $|Du|_M \in \mathcal{L}^1_{g_M}(M)$, relation (51) guarantees that $|\nabla h| \in \mathcal{L}^1_g(\Sigma(u))$. Therefore, the result follows by applying Theorem 5.2. \square

From Theorem 6.5 we obtain the following applications.

Corollary 6.6. *For any constants $c < 0$ and $0 < \beta < 1$, the only bounded smooth function $u : \mathbb{R}^n \rightarrow \mathbb{R}^+$, with $u(x) \leq (-\frac{2n}{3c})^{\frac{3}{5}}$ for all $x \in \mathbb{R}^n$, $|Du|_{\mathbb{R}^n} \in \mathcal{L}^1_{g_{\mathbb{R}^n}}(\mathbb{R}^n)$ and which is solution of the system (44), is the constant $u = (-\frac{2n}{3c})^{\frac{3}{5}}$.*

Corollary 6.7. *Let M^n be a complete Riemannian manifold. For any constants $c < 0$ and $0 < \beta < 1$, the only bounded smooth function $u : M^n \rightarrow \mathbb{R}$, with $u(x) \leq \log(-\frac{n}{c})$ for all $x \in M^n$, $|Du|_M \in \mathcal{L}^1_{g_M}(M)$ and which is solution of the system (45), is the constant $u = \log(-\frac{n}{c})$.*

Taking into account once more relation (48), it is not difficult to see that from Theorem 5.11 we obtain the following nonexistence result.

Theorem 6.8. *Let $\overline{M}^{-n+1} = -I \times_f M^n$ be a GRW spacetime satisfying (34), occurring the equality only at isolated points of I , and whose Riemannian fiber M^n is complete noncompact with nonnegative Ricci curvature. Suppose that c is a constant such that $\zeta_c(t) \geq 0$ for all $t \in I$. There does not exist a bounded entire solution $u \in C^\infty(M)$ of the spacelike mean curvature flow soliton equation (39), with $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$, and such that $(f(u))^{-1} \in \mathcal{L}^q_{g_M}(M)$ for some q with $q > \gamma + 3$.*

We have the following applications of Theorem 6.8.

Corollary 6.9. *For any constants $c \geq 0$ and $0 < \beta < 1$, there does not exist a bounded smooth function $u : \mathbb{R}^n \rightarrow \mathbb{R}^+$ such that $u^{-\frac{2}{3}} \in \mathcal{L}^q_{g_{\mathbb{R}^n}}(\mathbb{R}^n)$, for some $q > 3$, and which is a solution of the system (44).*

Corollary 6.10. *Let M^n be a complete noncompact Riemannian manifold with nonnegative Ricci curvature. For any constants $c \geq 0$ and $0 < \beta < 1$, there does not exist a bounded smooth function $u : M^n \rightarrow I$ such that $e^{-u} \in \mathcal{L}^q_{g_M}(M)$, for some $q > 3$, and which is a solution of the system (45).*

Observing once more that the assumption $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$, implies that the hyperbolic angle function Θ given by (43) is bounded, Theorem 5.15 allows us to obtain the following result.

Theorem 6.11. *Let $\overline{M}^{-n+1} = -I \times_f M^n$ be a spatially parabolic covered GRW spacetime satisfying (34), holding the equality only at isolated points of I . Suppose*

that c is a constant such that $\bar{\zeta}_c(t) \geq 0$ for all $t \in I$. If $\Sigma(u) \subset \overline{M}^{n+1}$ is an entire spacelike graph determined by a bounded function $u \in C^\infty(M)$ which is solution of the spacelike mean curvature flow soliton equation (40) with $|Du|_M \leq \beta f(u)$, for some constant $0 < \beta < 1$, then $u \equiv t_*$ for some $t_* \in I$ which is implicitly given by the condition $\zeta_c(t_*) = 0$.

We finish this manuscript with the following applications of Theorem 6.11.

Corollary 6.12. *For any constants $c < 0$ and $0 < \beta < 1$, the only bounded smooth function $u : \mathbb{R}^2 \rightarrow \mathbb{R}^+$, with $u(x) \leq (-\frac{4}{3c})^{\frac{3}{5}}$ for all $x \in \mathbb{R}^2$, and which is solution of the system (44) for $n = 2$, is the constant $u = (-\frac{4}{3c})^{\frac{3}{5}}$.*

Corollary 6.13. *Let M^n be a complete Riemannian manifold with parabolic universal covering. For any constants $c < 0$ and $0 < \beta < 1$, the only bounded smooth function $u : M^n \rightarrow \mathbb{R}$, with $u(x) \leq \log(-\frac{n}{c})$ for all $x \in M^n$, and which is solution of the system (45), is the constant $u = \log(-\frac{n}{c})$.*

Acknowledgements

The authors would like to thank the referee for reading the manuscript in great detail and for his valuable suggestions and useful comments which improved the paper.

References

- [1] AARONS, MARK A. S. Mean curvature flow with a forcing term in Minkowski space. *Calc. Var. PDE* **25** (2005), 205–246. MR2188747, Zbl 1094.53060, doi: 10.1007/s00526-005-0351-8. 555
- [2] AHLFORS, LARS V. Sur le type d'une surface de Riemann. *C.R. Acad. Sc. Paris* **201** (1935), 30–32. Zbl 0011.40701. 571
- [3] ALBUJER, ALMA L. New examples of entire maximal graphs in $\mathbb{H}^2 \times \mathbb{R}_1$. *Diff. Geom. Appl.* **26** (2008), 456–462. MR2424025, Zbl 1147.53047, doi: 10.1016/j.difgeo.2007.11.035. 571
- [4] ALBUJER, ALMA L.; ALÍAS, LUIS J. Spacelike hypersurfaces with constant mean curvature in the steady state space. *Proc. Amer. Math. Soc.* **137** (2009), 711–721. MR2448594, Zbl 1162.53043, doi: 10.1090/S0002-9939-08-09546-4. 559
- [5] ALEDO, JUAN A.; ALÍAS, LUIS J.; ROMERO, ALFONSO. Integral formulas for compact space-like hypersurfaces in de Sitter space: applications to the case of constant higher order mean curvature. *J. Geom. Phys.* **31** (1999), 195–208. MR1706636, Zbl 0969.53031, doi: 10.1016/S0393-0440(99)00008-X. 559
- [6] ALEDO, JUAN A.; RUBIO, RAFAEL M.; SALAMANCA, JUAN J. Space-like hypersurfaces with functionally bounded mean curvature in Lorentzian warped products and generalized Calabi-Bernstein-type problems. *Proc. Royal Soc. Edinburgh: Section A* **149** (2019), 849–868. MR3988624, Zbl 1439.53060, doi: 10.1017/prm.2018.7. 556, 566, 570
- [7] ALÍAS, LUIS J.; BRASIL, ALDIR, JR.; GERVASIO COLARES, A. Integral formulae for spacelike hypersurfaces in conformally stationary spacetimes and applications. *Proc. Royal Soc. Edinburgh: Section A* **149** (2019), 849–868. MR1998575, Zbl 1053.53038, doi: 10.1017/S0013091502000500. 557

- [8] ALÍAS, LUIS J.; GERVASIO COLARES, A. Uniqueness of spacelike hypersurfaces with constant higher order mean curvature in generalized Robertson-Walker spacetimes. *Math. Proc. Cambridge Philos. Soc.* **143** (2007), 703–729. MR2373968, Zbl 1131.53035, doi: 10.1017/S0305004107000576. 561
- [9] ALÍAS, LUIS J.; GERVASIO COLARES, A.; DE LIMA, HENRIQUE F. Uniqueness of spacelike hypersurfaces with constant higher order mean curvature in generalized Robertson-Walker spacetimes. *Bull. Brazilian Math. Soc.* **44** (2013), 195–217. MR3077640, Zbl 1294.53054, doi: 10.1007/s00574-013-0009-7. 573, 574
- [10] ALÍAS, LUIS J.; DE LIRA, JORGE H.; RIGOLI, MARCO. Mean curvature flow solitons in the presence of conformal vector fields. *J. Geom. Anal.* **30** (2020), 1466–1529. MR4081322, Zbl 1436.53066, doi: 10.1007/s12220-019-00186-3. 555, 556, 558
- [11] ALÍAS, LUIS J.; MASTROLIA, PAOLO; RIGOLI, MARCO. Maximum principles and geometric applications. Springer Monographs in Mathematics. *Springer, Cham*, 2016. xvii+570 pp. ISBN: 978-3-319-24335-1; 978-3-319-24337-5. MR3445380, Zbl 1346.58001, doi: 10.1007/978-3-319-24337-5. 560
- [12] ALÍAS, LUIS J.; ROMERO, ALFONSO; SÁNCHEZ, MIGUEL. Uniqueness of complete spacelike hypersurfaces of constant mean curvature in generalized Robertson-Walker spacetimes. *Gen. Relat. Grav.* **27** (1995), 71–84. MR1310212, Zbl 0908.53034, doi: 10.1007/BF02105675. 556, 571
- [13] ARAÚJO, JOGLI G.; DE LIMA, HENRIQUE F.; GOMES, WALLACE F. Uniqueness and nonexistence of complete spacelike hypersurfaces, Calabi-Bernstein type results and applications to Einstein-de Sitter and steady state type spacetimes. *Rev. Mat. Complut.* **34** (2021), 653–673. MR4302236, Zbl 7404682, doi: 10.1007/s13163-020-00375-7. 556
- [14] BATISTA, MÁRCIO; DE LIMA, HENRIQUE F. Spacelike translating solitons in Lorentzian product spaces: Nonexistence, Calabi-Bernstein type results and examples. *Comm. Contemp. Math.* **24** (2022), 2150034 (20 pages). MR4492665, Zbl 7597147, doi: 10.1142/S0219199721500346. 556
- [15] BESSE, ARTHUR L. Einstein manifolds. *Ergebnisse der Mathematik und ihrer Grenzgebiete; 3. Folge, Band 10.* Springer, Berlin, Heidelberg, New York, London, Paris, Tokyo, 1987. xii+510 pp. ISBN: 978-3-540-74120-6. MR0867684, Zbl 0613.53001, doi: 10.1007/978-3-540-74311-8. 563
- [16] BONDI, H.; GOLD, T. On the generation of magnetism by fluid motion. *Monthly Not. Roy. Astr. Soc.* **108** (1948), 252–270. MR0044363, Zbl 0041.59407, doi: 10.1093/mnras/110.6.607. 559
- [17] COLOMBO, GIULIO; MARI, LUCIANO; RIGOLI, MARCO. Remarks on mean curvature flow solitons in warped products. *Discrete Contin. Dyn. Syst., Ser. S* **13** (2020), 1957–1991. MR4097628, Zbl 1440.53105, doi: 10.3934/dcdss.2020153. 555, 556, 558
- [18] ECKER, KLAUS. On mean curvature flow of spacelike hypersurfaces in asymptotically flat spacetimes. *J. Austral. Math. Soc., Ser. A* **55** (1993), 41–59. MR1231693, Zbl 0789.58020, doi: 10.1017/S1446788700031918. 555
- [19] ECKER, KLAUS. Interior estimates and longtime solutions for mean curvature flow of noncompact spacelike hypersurfaces in Minkowski space. *J. Diff. Geom.* **46** (1997), 481–498. MR1484889, Zbl 0909.53045, doi: 10.4310/jdg/1214459975. 555
- [20] ECKER, KLAUS. Mean curvature flow of spacelike hypersurfaces near null initial data. *Comm. Anal. Geom.* **11** (2003), 181–205. MR2014875, Zbl 1129.53303, doi: 10.4310/CAG.2003.v11.n2.a1. 555
- [21] ECKER, KLAUS; HUISKEN, GERHARD. Parabolic methods for the construction of spacelike slices of prescribed mean curvature in cosmological spacetimes. *Comm. Math. Phys.* **135** (1991), 595–613. MR1091580, Zbl 0721.53055, doi: 10.1007/BF02104123. 555
- [22] GAO, SHANZE; LI, GUANGHAN; WU, CHUANXI. Translating spacelike graphs by mean curvature flow with prescribed contact angle. *Arch. Math.* **103** (2014), 499–508. MR3282644, Zbl 1305.53055, doi: 10.1007/s00013-014-0699-0. 555

- [23] GERHARDT, CLAUS. Hypersurfaces of prescribed mean curvature in Lorentzian manifolds. *Math. Z.* **235** (2000), 83–97. MR1785072, Zbl 1017.53053, doi: 10.1007/s002090000124. 555
- [24] GUILFOYLE, BRENDAN; KLINGENBERG, WILHELM. Mean curvature flow of compact spacelike submanifolds in higher codimension. *Trans. Amer. Math. Soc.* **372** (2019), 6263–6281. MR4024521, Zbl 1436.53068, doi: 10.1090/tran/7766. 555
- [25] HAWKING, STEPHEN W.; ELLIS, GEORGE F. R. The Large Scale Structure of Space-Time. Cambridge Monographs on Mathematical Physics. *Cambridge University Press, London*, 1973. xx+392 pp. ISBN: 0-521-09906-4. MR0424186, Zbl 7633910, doi: 10.1017/CBO9780511524646. 559
- [26] HOYLE, F. A new model for the expanding universe. *Monthly Not. Roy. Astr. Soc.* **108** (1948), 372–382. Zbl 0031.38304, doi: 10.1093/mnras/108.5.372. 559
- [27] HUBER, ALFRED. On subharmonic functions and differential geometry in the large. *Comment. Math. Helv.* **32** (1958), 13–72. MR0094452, Zbl 0080.15001, doi: 10.1007/BF02564570. 571
- [28] HUISKEN, GERHARD; YAU, SHING-TUNG. Definition of center of mass for isolated physical system and unique foliations by stable spheres with constant curvature. *Invent. Math.* **124** (1996), 281–311. MR1369419, Zbl 0858.53071, doi: 10.1007/s002220050054. 555
- [29] JIAN, HUAI-YU. Translating solitons of mean curvature flow of noncompact spacelike hypersurfaces in Minkowski space. *J. Diff. Eq.* **220** (2006), 147–162. MR2182083, Zbl 1189.53065, doi: 10.1016/j.jde.2005.08.005. 555
- [30] KÜHNEL, WOLFGANG. Conformal transformations between Einstein spaces. *Conformal Geometry (R.S. Kulkarni and U. Pinkall, eds.), Aspects of Math., vol. E12, Braunschweig*, 105–146. *Vieweg+Teubner Verlag, Wiesbaden*, 1988. ISBN: 978-3-528-08982-5. MR0979791, Zbl 0667.53039, doi: 10.1007/978-3-322-90616-8-5. 563
- [31] LAMBERT, BEN. A note on the oblique derivative problem for graphical mean curvature flow in Minkowski space. *Abh. Math. Semin. Univ. Hambg.* **82** (2012), 115–120. MR2922727, Zbl 1276.53071, doi: 10.1007/s12188-012-0066-7. 555
- [32] LAMBERT, BEN. The perpendicular Neumann problem for mean curvature flow with a timelike cone boundary condition. *Trans. Am. Math. Soc.* **366** (2014), 3373–3388. MR3192599, Zbl 1296.53131, doi: 10.1090/S0002-9947-2014-05865-0. 555
- [33] LAMBERT, BEN; LOTAY, JASON D. Spacelike mean curvature flow. *J. Geom. Anal.* **31** (2021), 1291–1359. MR4215263, Zbl 1465.53081, doi: 10.1007/s12220-019-00266-4. 555, 558
- [34] LI, GUANGHAN; SALAVESSA, ISABEL M. C. Mean curvature flow of spacelike graphs. *Math. Z.* **269** (2011), 697–719. MR2860260, Zbl 1231.53057, doi: 10.1007/s00209-010-0768-4. 555
- [35] MONTIEL, SEBASTIÁN. Uniqueness of spacelike hypersurfaces of constant mean curvature in foliated spacetimes. *Math. Ann.* **314** (1999), 529–553. MR1704548, Zbl 0965.53043, doi: 10.1007/s002080050306. 559, 560, 561
- [36] OMORI, HIDEKI. Isometric immersions of Riemannian manifolds. *J. Math. Soc. Japan* **19** (1967), 205–214. MR0215259, Zbl 0154.21501, doi: 10.2969/jmsj/01920205. 560
- [37] O’NEILL, BARRETT. Semi-Riemannian Geometry with Applications to Relativity. *Pure and Applied Mathematics*, 103. *Academic Press, London*, 1983. xiii+468 pp. ISBN: 978-0125267403. MR0719023, Zbl 0531.53051. 556, 557, 559, 561
- [38] ROMERO, ALFONSO; RUBIO, RAFAEL M.; SALAMANCA, JUAN J. Uniqueness of complete maximal hypersurfaces in spatially parabolic generalized Robertson-Walker spacetimes. *Class. Quant. Grav.* **30** (2013), 115007. MR3055096, Zbl 1271.83011, doi: 10.1088/0264-9381/30/11/115007. 570
- [39] SCHOUTEN, J. A. Der Ricci-Kalkül. Eine Einführung in die Neueren Methoden und Probleme der Mehrdimensionalen Differentialgeometrie (Grundlehren der mathematischen

- Wissenschaften, 10) (German Edition). *Springer-Verlag, Berlin*, 1924. x, 312 S., 7 Abb. (1978) ISBN: 978-3642517983. MR516659, Zbl 0403.53007. 563
- [40] SPRUCK, JOEL; XIAO, LING. Entire downward translating solitons of the mean curvature flow in Minkowski space. *Proc. Am. Math. Soc.* **144** (2016), 3517–3526. MR3503719, Zbl 1346.53062, doi: 10.1090/proc/12969. 555
- [41] YAU, SHING-TUNG. Harmonic functions on complete Riemannian manifolds. *Comm. Pure Appl. Math.* **28** (1975), 201–228. MR0431040, Zbl 0291.31002, doi: 10.1002/cpa.3160280203. 560
- [42] YAU, SHING-TUNG. Some function-theoretic properties of complete Riemannian manifolds and their applications to geometry. *Indiana Univ. Math. J.* **25** (1976), 659–670. MR0417452, Zbl 0335.53041, doi: 10.1512/iumj.1976.25.25051. 566, 568

(Márcio Batista) CPMAT-IM, UNIVERSIDADE FEDERAL DE ALAGOAS, 57072-970 MACEIÓ, ALAGOAS, BRAZIL
mhbs@mat.ufal.br

(Giovanni Molica Bisci) DIPARTIMENTO DI SCIENZE PURE E APPLICATE (DISPEA), UNIVERSITÀ DEGLI STUDI DI URBINO CARLO BO, PIAZZA DELLA REPUBBLICA 13, 61029 URBINO (PESARO E URBINO), ITALY
giovanni.molicabisci@uniurb.it

(Henrique F. de Lima) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE CAMPINA GRANDE, 58.429-970 CAMPINA GRANDE, PARAÍBA, BRAZIL
henriquedelima74@gmail.com

(Wallace F. Gomes) DEPARTAMENTO DE MATEMÁTICA, UNIVERSIDADE FEDERAL DE CAMPINA GRANDE, 58.429-970 CAMPINA GRANDE, PARAÍBA, BRAZIL
wfgomes.uepb@gmail.com

This paper is available via <http://nyjm.albany.edu/j/2023/29-23.html>.