

Gabor Frames with Hermite Functions

Frames de Weyl-Heisenberg et Fonctions d'Hermite

Karlheinz Gröchenig^{a,1} Yurii Lyubarskii^b

^a*Faculty of Mathematics, University of Vienna, Nordbergstrasse 15, A-1090 Vienna, Austria*

^b*Department of Mathematics, Norwegian University of Science and Technology, 7491, Trondheim, Norway*

Abstract

We investigate Gabor frames based on a linear combination of Hermite functions H_n . We derive sufficient conditions on the lattice $\Lambda \subseteq \mathbb{R}^2$ such that the Gabor system $\{e^{2\pi i \lambda_2 t} H_n(t - \lambda_1) : \lambda = (\lambda_1, \lambda_2) \in \Lambda\}$ is a frame for $L^2(\mathbb{R})$. An example supports our conjecture that our conditions are sharp. The main tools are growth estimates for the Weierstrass σ -function and a new type of interpolation problem for entire functions on the Bargmann-Fock space.

Résumé

Nous étudions les propriétés de frame de l'ensemble $\{e^{2\pi i \lambda_2 t} H_n(t - \lambda_1) : \lambda = (\lambda_1, \lambda_2) \in \Lambda\}$, où H_n est une fonction de Hermite et Λ est un réseau dans \mathbb{R}^2 . Nous donnons des conditions suffisantes sur la densité de Λ pour que la propriété de frame soit satisfaite. Un contre-exemple suggère que nos conditions sont aussi nécessaires. Les outils principaux sont des estimations de croissance pour la fonction σ de Weierstrass et un nouveau type d'interpolation dans l'espace de Bargmann-Fock.

Version française abrégée

Soit $z = (x, y) \in \mathbb{R}^2$ où $z = x + yi \in \mathbb{C}$, on dénote par

$$\pi_z f(t) = e^{2\pi i y t} f(t - x) \quad x, y, t \in \mathbb{R}.$$

le décalage en temps-fréquence d'une fonction f . Un réseau $\Lambda \subset \mathbb{R}^2$ est un sous-groupe discret de la forme $\Lambda = A\mathbb{Z}^2$, où A est une matrice inversible. L'aire du rectangle fondamental se dénote par $s(\Lambda) = |\det A|$, la densité usuel de Λ est $s(\Lambda)^{-1}$.

Email addresses: karlheinz.groechenig@univie.ac.at (Karlheinz Gröchenig), yura@math.ntnu.no (Yurii Lyubarskii).

¹ K.G. was supported by the Marie-Curie Excellence Grant MEXT-CT-2004-517154, Ya. L. was partially supported by

On étudie des conditions sous lesquelles est satisfaite l'équivalence

$$A\|f\|_{L^2(\mathbb{R})}^2 \leq \sum_{\lambda \in \Lambda} |\langle f, \pi_\lambda g \rangle_{L^2(\mathbb{R})}|^2 \leq B\|f\|_{L^2(\mathbb{R})}^2, \quad \forall f \in L^2(\mathbb{R}). \quad (1)$$

Si (1) est satisfait, on dit que l'ensemble $\mathcal{G}(g, \Lambda) = \{\pi_\lambda g : \lambda \in \Lambda\}$ est un frame de Gabor (ou frame de Weyl-Heisenberg) pour $L^2(\mathbb{R})$.

En général, pour une fonction g arbitraire, la caractérisation de tous les réseaux pour lesquels $\mathcal{G}(g, \Lambda)$ est un frame, est extrêmement difficile. En fait, une telle caractérisation n'est connue que pour la fonction Gaussienne $g(t) = e^{-\pi t^2}$ [12, 13] et pour le cosinus hyperbolique $g(t) = (\cosh at)^{-1}$ [11].

Notre résultat principal traite le cas où g est une fonction de Hermite $H_n(t) = e^{\pi t^2} \left(\frac{d}{dt}\right)^n (e^{-2\pi t^2})$.

Théorème 0.1. *Si $s(\Lambda) < \frac{1}{n+1}$, l'ensemble $\mathcal{G}(H_n, \Lambda)$ est un frame pour $L^2(\mathbb{R})$.*

Dans le cas $n = 0$, où H_0 est la fonction Gaussienne, on retrouve le résultat classique de [12, 13]. En fait, si $n = 0$, la condition $s(\Lambda) < 1$ est suffisante et nécessaire. Dans le cas $n > 0$, la condition sur la croissance de la densité $s(\Lambda)^{-1}$ est tout à fait surprenante, parce que l'intuition générale disait que pour chaque fonction "raisonnable" la condition $s(\Lambda) < 1$ était suffisante pour que $\mathcal{G}(g, \Lambda)$ soit un frame.

Pour démontrer le théorème, on utilise une caractérisation abstraite des frames de Gabor par les identités de Wexler-Raz, cf. [4]. Soit $\Lambda^\circ = \frac{1}{s(\Lambda)}$ le réseau adjoint de Λ . Alors, $\mathcal{G}(g, \Lambda)$ est un frame de $L^2(\mathbb{R})$, si et seulement si il existe une fonction duale $\gamma \in L^2(\mathbb{R})$ telle que

$$\langle \gamma, \pi_\mu g \rangle = \delta_{\mu,0} \quad \forall \mu \in \Lambda^\circ, \quad (2)$$

et si $\mathcal{G}(g, \Lambda)$ est une séquence de Bessel (une condition technique qui est presque toujours satisfaite).

L'existence d'une fonction duale est déduite en transposant le problème d'analyse réelle en un problème d'analyse complexe. En utilisant la transformée de Bargmann B , on traduit les conditions (2) en un problème d'interpolation dans l'espace de Bargmann-Fock \mathcal{F} qui est constitué des fonctions entières dotées de la norme

$$\|F\|_{\mathcal{F}}^2 := \int_{\mathbb{C}} |F(\zeta)|^2 e^{-\pi|\zeta|^2} d\zeta < \infty.$$

Le lemme suivant se déduit à partir des propriétés de la transformée de Bargmann et du fait que $BH_n(z) = c_n z^n$ pour une constante $c_n \in \mathbb{R}$.

Lemme 0.2. *L'ensemble $\mathcal{G}(H_n, \Lambda)$ est un frame de Gabor pour $L^2(\mathbb{R})$ si et seulement si il existe une fonction $G \in \mathcal{F}$ telle que*

$$\sum_{k=0}^n \binom{n}{k} (-\pi \bar{\mu})^k G^{(n-k)}(\mu) = \delta_{\mu,0} \quad \forall \mu \in \Lambda^\circ. \quad (3)$$

Dans ce cas on peut prendre $\gamma = B^{-1}G$ comme fonction duale de H_n .

Remarque 1. Ces identités sont un nouveau type de problème d'interpolation avec des dérivées et ne sont pas standards, parce que les coefficients sont antiholomorphes.

La théorie classique des fonctions elliptiques fournit une fonction qui s'évanouit sur un réseau [1], la fonction σ de Weierstrass définie par

$$\sigma_{\Lambda^\circ}(u) = \sigma(u) = u \prod_{\lambda \in \Lambda^\circ \setminus \{0\}} \left(1 - \frac{u}{\lambda}\right) e^{\frac{u}{\lambda} + \frac{u^2}{2\lambda^2}}. \quad (4)$$

Alors on pose ($a \in \mathbb{C}$)

$$G(z) = n!^{-1} z^{-1} \sigma_{\Lambda^\circ}(z)^{n+1} e^{a(\Lambda^\circ)(n+1)z^2}.$$

Par construction, G satisfait les conditions $G^{(k)}(\mu) = \delta_{\mu,0} \delta_{k,n}$ for $\mu \in \Lambda^\circ$ and $0 \leq k \leq n$, et par conséquent, (4) est aussi satisfait. Il reste à démontrer que G appartient à l'espace \mathcal{F} . Dans le cas $s(\Lambda) = s(\Lambda^\circ)^{-1} < 1/(n+1)$ on peut choisir $a \in \mathbb{C}$ tel que la fonction $|G(z)|e^{-(n+1)|z|^2/s(\Lambda^\circ)}$ soit Λ° -periodique. Alors la conclusion $G \in \mathcal{F}$ découle des estimations très souples de la croissance de σ_{Λ° [9].

1. Introduction

Given a function $g \in L^2(\mathbb{R})$ and a lattice $\Lambda \subseteq \mathbb{R}^2$, we study the problem of when the set $\{e^{2\pi i \lambda_2 t} g(t - \lambda_1) : \lambda = (\lambda_1, \lambda_2) \in \Lambda\}$ is a frame for $L^2(\mathbb{R}^d)$. Precisely, write $\pi_\lambda g = e^{2\pi i \lambda_2 t} g(t - \lambda_1)$ for the time-frequency shift by $\lambda \in \mathbb{R}^2$. Then we call the set $\mathcal{G}(g, \Lambda) = \{\pi_\lambda g : \lambda \in \Lambda\}$ a *Gabor frame* or *Weyl-Heisenberg frame*, whenever there exist positive constants $A, B > 0$ such that the inequalities

$$A\|f\|_{L^2(\mathbb{R})}^2 \leq \sum_{\lambda \in \Lambda} |\langle f, \pi_\lambda g \rangle_{L^2(\mathbb{R})}|^2 \leq B\|f\|_{L^2(\mathbb{R})}^2, \quad \forall f \in L^2(\mathbb{R}) \quad (5)$$

hold. Gabor frames originate in quantum mechanics through J. von Neumann and in information theory through D. Gabor [7] and nowadays have many applications in signal processing. A large body of results describes the structure of Gabor frames and provides sufficient conditions of a qualitative nature on $\mathcal{G}(g, \Lambda)$ to form a (Gabor) frame, see [3, 8] for details and references.

The general problem of characterizing *all* lattices Λ for which $\mathcal{G}(g, \Lambda)$ is a frame seems to be extremely difficult. In fact, this problem is solved only for two classes of basis functions, namely for the Gaussians $H_0(t) = e^{-at^2}$, $a > 0$, in [12, 13] and for the hyperbolic cosine $g(t) = (\cosh at)^{-1}$ in [11]. For the Gaussian, our understanding is based on the connection between the frame property of $\mathcal{G}(g, \Lambda)$ and a classical interpolation problem in the Bargmann-Fock space of entire functions.

In this article we study the frame property of $\mathcal{G}(g, \Lambda)$ in the case when g is a Hermite function $g = H_n$ or a finite linear combination of Hermite functions. Although this case seems to be similar to the case of Gaussian Gabor frames, it is not and leads us to a striking result: the critical density of the lattice for $\mathcal{G}(H_n, \Lambda)$ to be a Gabor frame depends on n and increases with n . For $n = 0$ (the case of the Gaussian), we recover the classical results [12, 13]. An example for $n = 1$ suggests that the sufficient conditions are sharp for all n . The technical question is a new type of interpolation problem in the Bargmann-Fock space of entire functions, but this problem is no longer "purely holomorphic": the values of linear combination of a function and its derivatives are prescribed in the lattice points, the coefficients of this combination are antiholomorphic polynomials.

The next section includes the precise setting of the problem, auxiliary facts from time frequency analysis and complex analysis. In section 3 we prove the main result and remark that it is sharp for $n = 1$. In a subsequent paper we will give a more detailed analysis and applications to the Balian-Low theorem, Gabor superframes, and the time-frequency analysis of distributions.

2. Tools

Gabor Frames and Bessel sequences (see [8] for an exposition). A *lattice* is a discrete subgroup in \mathbb{R}^2 of the form $\Lambda = A\mathbb{Z}^2$, where A is an invertible 2×2 matrix. The size of Λ is defined as the area of its cell $s(\Lambda) = |\det A|$, the density Λ is $d(\Lambda) = s(\Lambda)^{-1}$, which equals the Beurling density (see, e.g., [13]). The *adjoint lattice* is $\Lambda^\circ := \frac{1}{s(\Lambda)} \Lambda$ and has size $s(\Lambda^\circ) = s(\Lambda)^{-1}$.

We say that $\mathcal{G}(g, \Lambda) = \{\pi_\lambda g : \lambda \in \Lambda\}$ is a *Bessel sequence*, if only the right-hand inequality of (5) is satisfied. The Bessel property of $\mathcal{G}(g, \Lambda)$ is a technical condition and depends both on g and Λ , but it is easy to formulate universal conditions in terms of Bargmann transform of g .

Lemma 2.1. *Assume that g satisfies the condition*

$$\|g\|_{M^1} := \int_{\mathbb{C}} \int_{\mathbb{C}} |Bg(\zeta)| e^{-\pi|\zeta|^2/2} dm_\zeta < \infty, \quad (6)$$

then the sequence $\mathcal{G}(g, \Lambda)$ is a Bessel sequence in $L^2(\mathbb{R})$ for each lattice Λ [5].

Remark 1. The M^1 -norm in (6) defines the Feichtinger algebra, which occurs in numerous applications in harmonic analysis [5].

The frame property (5) of the system $\mathcal{G}(g, \Lambda)$ is much more delicate and lies much deeper. A necessary condition is given by the following density theorem [3, 8].

Theorem 2.2. *If $\mathcal{G}(g, \Lambda)$ is a frame for $L^2(\mathbb{R})$, then $s(\Lambda) \leq 1$.*

An abstract characterization of Gabor frames is provided by the so-called Wexler-Raz identities [4, 10]

Theorem 2.3. *Let $g \in L^2(\mathbb{R})$ and $\Lambda \subset \mathbb{R}^2$ be a lattice with adjoint lattice Λ° . Then the following are equivalent:*

(i) $\mathcal{G}(g, \Lambda)$ is a Gabor frame for $L^2(\mathbb{R})$.

(ii) There exists a function $\gamma \in L^2(\mathbb{R})$ (a dual window), such that $\mathcal{G}(\gamma, \Lambda)$ is a Bessel sequence in L^2 and

$$\langle \gamma, \pi_\mu g \rangle = \delta_{\mu, 0} \quad \forall \mu \in \Lambda^\circ. \quad (7)$$

Transition to complex analysis. The Bargmann-Fock space \mathcal{F} is the Hilbert space \mathcal{F} of entire functions F such that

$$\|F\|_{\mathcal{F}}^2 := \int_{\mathbb{C}} |F(\zeta)|^2 e^{-\pi|\zeta|^2} d\zeta < \infty, \quad (8)$$

where $d\zeta$ is the Lebesgue measure on \mathbb{C} . The corresponding inner product is denoted by $\langle \cdot, \cdot \rangle_{\mathcal{F}}$.

The Bargmann transform B of a function f on \mathbb{R} is defined as

$$B : f \mapsto Bf(\zeta) = F(\zeta) = 2^{1/4} \int_{-\infty}^{\infty} f(t) e^{2\pi t \zeta - \pi t^2 - \pi \zeta^2 / 2} dt \quad z \in \mathbb{C}. \quad (9)$$

Proposition 2.4 ([6]). (i) B is a unitary mapping from $L^2(\mathbb{R})$ onto the Bargmann-Fock space \mathcal{F} .

(ii) \mathcal{F} is a reproducing kernel Hilbert space. If $e_\zeta(w) = e^{\pi \zeta w}$, then $F(\zeta) = \langle F, e_\zeta \rangle_{\mathcal{F}}$. (iii) Define a "shift" $\beta_z, z \in \mathbb{C}$, on \mathcal{F} by $\beta_z F(\zeta) = e^{i\pi x y} e^{-\pi|z|^2/2} e^{\pi \zeta \bar{z}} F(\zeta - z)$, then B intertwines the shifts π and β in the sense that $\beta_z = B\pi_z B^{-1}$ for all $z \in \mathbb{C}$. (iv) Let $H_n(t) = e^{\pi t^2} \left(\frac{d}{dt}\right)^n (e^{-2\pi t^2})$ denote the n -th Hermite function, then $BH_n(z) = \pi^{n/2} n!^{-1/2} z^n$.

Growth Estimates for Weierstrass's σ -Functions. We identify $(x, y) \in \mathbb{R}^2$ with $z = x + iy \in \mathbb{C}$. Then we may write the lattice Λ as $\Lambda = \{m_1 \lambda_1 + m_2 \lambda_2 : m_1, m_2 \in \mathbb{Z}\}$ for some basis $\lambda_1, \lambda_2 \in \mathbb{C}$ satisfying $\text{Im}(\lambda_2 / \lambda_1) > 0$. The size of Λ is then given by $s(\Lambda) = \frac{1}{2i} (\bar{\lambda}_1 \lambda_2 - \bar{\lambda}_2 \lambda_1) = |\lambda_1|^2 \text{Im}(\lambda_2 / \lambda_1) > 0$.

The Weierstrass σ -function corresponding to the lattice Λ is

$$\sigma_\Lambda(u) = \sigma(u) = u \prod_{\lambda \in \Lambda \setminus \{0\}} \left(1 - \frac{u}{\lambda}\right) e^{\frac{u}{\lambda} + \frac{u^2}{2\lambda^2}} \quad (10)$$

The theory of elliptic functions furnishes the following properties of σ_Λ , see, e.g., [1].

Proposition 2.5. (i) σ_Λ is an entire function whose zero set is precisely Λ .

(iii) Legendre's formula: $(\log \sigma(u))'$ is Λ -periodic. If $\eta_k = (\log \sigma(u + \lambda_k) - \log \sigma(u))' = \text{const.}$, then $\eta_1 \lambda_2 - \eta_2 \lambda_1 = 2\pi i$.

(iii) Multiplicative quasi-periodicity of σ : there exist $\eta_1, \eta_2 \in \mathbb{C}$ such that

$$\sigma(u + \lambda_k) = -\sigma(u) e^{\eta_k u + \frac{1}{2} \eta_k \lambda_k} \quad k = 1, 2. \quad (11)$$

In order to understand the growth properties of σ_Λ , we follow an argument of Hayman [9] who realized that, after a proper normalization and growth compensation, the absolute value of σ_Λ becomes a doubly periodic function. Let us define the numbers

$$\alpha(\Lambda) = \frac{i\pi}{\bar{\lambda}_1 \lambda_2 - \bar{\lambda}_2 \lambda_1} = \frac{\pi}{2s(\Lambda)} \quad \text{and} \quad a(\Lambda) = \frac{1}{2} \frac{\eta_1 \lambda_2 - \eta_2 \lambda_1}{\bar{\lambda}_1 \lambda_2 - \bar{\lambda}_2 \lambda_1}. \quad (12)$$

The following result can be deduced from (11) – (12) by a straightforward, but lengthy computation.

Theorem 2.6. *The function $|\sigma_\Lambda(u)e^{a(\Lambda)u^2}|e^{-\alpha(\Lambda)|u|^2}$ is periodic with periods λ_1 and λ_2 . Therefore, the modified sigma-function $\sigma_\Lambda(u)e^{a(\Lambda)u^2}$ vanishes exactly on the lattice Λ and satisfies the growth estimate*

$$|\sigma_\Lambda(u)e^{a(\Lambda)u^2}| \leq C e^{\frac{\pi}{2s(\Lambda)}|u|^2}. \quad (13)$$

3. Gabor Frames with Hermite Functions

Our main result provides a sufficient condition for the family $\mathcal{G}(H_n, \Lambda)$ to form a frame for $L^2(\mathbb{R})$. The condition involves only the size of the lattice.

Theorem 3.1. *Let $n \in \mathbb{Z}, n \geq 0$. If $s(\Lambda) < (n+1)^{-1}$, then $\mathcal{G}(H_n, \Lambda)$ is a frame for $L^2(\mathbb{R})$.*

In the first step of the proof, we translate the frame property to an interpolation problem in \mathcal{F} .

Given a (real-valued) window function g we consider the short time Fourier transform (STFT) V_g as a function of a complex variable $z = x + iy$ instead of “time” $x \in \mathbb{R}$ and “frequency” $y \in \mathbb{R}$:

$$V_g f(z) = \langle f, \pi_z g \rangle_{L^2} = \int_{-\infty}^{\infty} f(t)g(t-x)e^{-2i\pi yt} dt$$

The next proposition details a formula for the STFT with respect to a Hermite function.

Proposition 3.2. *Let $f \in L^2(\mathbb{R})$, $F = Bf$, and H_n be the n -th Hermite function. Then*

$$V_{H_n} f(z) = e^{i\pi xy} e^{-\pi|z|^2/2} \frac{1}{\sqrt{\pi^n n!}} \sum_{k=0}^n \binom{n}{k} (-\pi \bar{z})^k F^{(n-k)}(z). \quad (14)$$

Proof. With the properties of the Bargmann transform of Proposition 2.4, we obtain

$$\begin{aligned} V_{H_n} f(z) &= \langle f, \pi_z H_n \rangle_{L^2} = \langle F, \beta_z B H_n \rangle_{\mathcal{F}} = \kappa_n e^{i\pi xy} e^{-\pi|z|^2/2} \langle F(w), e^{\pi \bar{z} w} (w-z)^n \rangle_{\mathcal{F}} = \\ &= \kappa_n e^{i\pi xy} e^{-\pi|z|^2/2} \sum_{k=0}^n \binom{n}{k} (-\bar{z})^k \langle F(w), w^{n-k} e^{\pi \bar{z} w} \rangle_{\mathcal{F}}. \end{aligned}$$

The reproducing formula (??) yields $\langle F(w), w^{n-k} e^{\pi \bar{z} w} \rangle_{\mathcal{F}} = \pi^{k-n} F^{(n-k)}(z)$ and now (14) follows. \square

Combining this proposition with Theorem 2.3, we obtain the following criterion for Gabor frames with Hermite functions. We are lead to a new type of interpolation problem in the Bargmann-Fock space.

Proposition 3.3. *Let $\Lambda \subset \mathbb{R}^2$ be a lattice and $n \geq 0$. The set $\mathcal{G}(H_n, \Lambda)$ is a (Gabor) frame for $L^2(\mathbb{R})$ if and only if there exists a function $\gamma \in L^2(\mathbb{R})$ such that $\mathcal{G}(\gamma, \Lambda)$ is a Bessel sequence and the Bargmann transform $G = B\gamma$ satisfies*

$$\sum_{k=0}^n \binom{n}{k} (-\pi \bar{\mu})^k G^{(n-k)}(\mu) = \delta_{\mu,0} \quad \forall \mu \in \Lambda^\circ. \quad (15)$$

Remark. Amazingly enough, almost the same interpolation problem appears in the study of the Landau equation and its eigenspaces corresponding to high energy levels. The zero energy level is investigated in [2] and leads to a standard interpolation/uniqueness problem in the Fock space.

Sketch of the proof of Theorem 3.1. Let σ_{Λ° be the σ -function corresponding to the adjoint lattice Λ° of Λ . Denote

$$G(z) = n!^{-1} z^{-1} \sigma_{\Lambda^\circ}(z)^{n+1} e^{a(\Lambda^\circ)(n+1)z^2}. \quad (16)$$

It follows from (13) that

$$|G(z)| \leq \text{Const} e^{\pi k|z|^2/2} \quad \text{with} \quad k = \frac{n+1}{s(\Lambda^\circ)} = (n+1)s(\Lambda) < 1 \quad (17)$$

This growth estimates implies immediately that $G \in \mathcal{F}$. By construction, G solves the interpolation problem (15).

Now we choose the dual window to be $\gamma = B^{-1}G$. The growth estimate (17) yields $\gamma \in M^1$, and Lemma 2.1 implies that $\mathcal{G}(\gamma, \Lambda)$ is a Bessel sequence. By Proposition 3.3 $\mathcal{G}(H_n, \Lambda)$ is a frame. \square

Remarks. 1. It can be shown that $|\gamma(t)| \leq \text{Const } e^{-\epsilon t^2}$ and $|\hat{\gamma}(\omega)| \leq \text{Const } e^{-\epsilon \omega^2}$ for each $\epsilon < (1-k)/2$.
2. For $n = 0$ we recover the sharp result that $\mathcal{G}(H_0, \Lambda)$ is a frame if and only if $s(\Lambda) < 1$ from [12, 13].

A similar argument works for Gabor systems generated by a linear combination of Hermite functions.

Proposition 3.4. *Let $n \in \mathbb{Z}, n \geq 0$ and $g = \sum_{k=0}^n b_k H_k$, $b_n \neq 0$. If $s(\Lambda) < (n+1)^{-1}$, then $\mathcal{G}(g, \Lambda)$ is a frame for $L^2(\mathbb{R})$. The dual window γ may be chosen as in (16) for H_n .*

A Counter-Example. The function G that we constructed in the proof of Theorem 3.1 meets much stronger restrictions than just (15), namely $G^{(k)}(\mu) = \delta_{\mu,0} \delta_{k,n}$ for $\mu \in \Lambda^\circ$ and $0 \leq k \leq n$. The following statement provides some evidence that Theorem 3.1 is optimal.

Proposition 3.5. *If $\Lambda = \alpha\mathbb{Z} \times \beta\mathbb{Z}$ and $\alpha\beta = 1/2$, then $\mathcal{G}(H_{2n+1}, \Lambda)$ is not a Gabor frame.*

Proof. We use a Zak transform argument. For $\alpha > 0$ the Zak transform is defined by $Z_\alpha f(x, \xi) = \sum_{k \in \mathbb{Z}} f(x - \alpha k) e^{2i\pi \alpha k \xi}$. If $Z_\alpha g$ is continuous and $\mathcal{G}(g, \alpha\mathbb{Z} \times \frac{1}{2\alpha}\mathbb{Z})$ is a frame for $L^2(\mathbb{R})$, then the functions $Z_\alpha g(x, \xi)$ and $Z_\alpha g(x - \alpha/2, \xi)$ cannot have a common zero, see, e.g., [8, p. 159].

However, since $Z_\alpha H_{2n+1}$ is a continuous function and since H_{2n+1} is odd, we find that $Z_\alpha H_{2n+1}(0, 0) = 0$ and $Z_\alpha H_{2n+1}(\alpha/2, 0) = 0$, violating the necessary condition just described. \square

References

- [1] N.I. Akhiezer, *Elements of the theory of elliptic functions*, American Mathematical Society, Providence, RI, 1990.
- [2] G.Chistyakov, Yu. Lyubarskii, L. Pastur, *On completeness of random exponentials in the Bargmann-Fock space*, J. Math. Phys. 42 (2001), no. 8, 3754–3768
- [3] I. Daubechies. The wavelet transform, time-frequency localization and signal analysis. *IEEE Trans. Inform. Theory*, 36(5):961–1005, 1990.
- [4] I. Daubechies, H. J. Landau, and Z. Landau. Gabor time-frequency lattices and the Wexler-Raz identity. *J. Fourier Anal. Appl.*, 1(4):437–478, 1995.
- [5] H. G. Feichtinger and G. Zimmermann. A Banach space of test functions for Gabor analysis. In *Gabor analysis and algorithms*, pages 123–170. Birkhäuser Boston, Boston, MA, 1998.
- [6] G.B. Folland, *Harmonic analysis in phase space*, Princeton, NJ, 1989. x+277 pp.
- [7] D. Gabor. Theory of communication. *J. IEE (London)*, 93(III):429–457, 1946.
- [8] K. Gröchenig, *Foundations of time-frequency analysis*, Birkhäuser, Boston, 2001, xvi+359 pp.
- [9] W.K.Hayman, *The local growth of the power series: a survey of the Wiman-Valiron method*, Canad. math. Bull., 1974, 17, no.3, 317-358.
Appl. Comput. Harmon. Anal., 1(4):350–354, 1994.
- [10] A. J. E. M. Janssen. Duality and biorthogonality for Weyl-Heisenberg frames. *J. Fourier Anal. Appl.*, 1(4):403–436, 1995.
- [11] A. J. E. M. Janssen and T. Strohmer. Hyperbolic secants yield Gabor frames. *Appl. Comp. Harm. Anal.*, 2001.
- [12] Yu. Lyubarski, *Frames in the Bargmann space of entire functions*, Entire and subharmonic functions, 167–180, Adv. Soviet Math., 11, Amer. Math. Soc., Providence, RI, 1992.
- [13] K. Seip, R. Wallstén, *Density theorems for sampling and interpolation in the Bargmann-Fock space. II*, J. Reine Angew. Math. 429 (1992), 107–113.